

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF SOUTH DAKOTA



**For Control of Weeds in Soybeans in South Dakota West of Highway 281 in the Counties of Aurora, Beadle, Brown, Brule, Buffalo, Campbell, Charles Mix, Douglas, Edmunds, Faulk, Gregory, Hand, Hughes, Hyde, Jerauld, Lyman, McPherson, Potter, Spink, Sully, Tripp, Walworth**

EPA Reg. No. 5905-593

EPA SLN No. SD190007

**This label expires and must not be distributed or used in accordance with this SLN registration after December 31, 2024**

**DIRECTIONS FOR USE**

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered label.
- Failure to follow the directions for use and precautions on this label may result in poor pest control, crop injury, or illegal residues.

**Application Methods**

Apply SINISTER at 0.49 pt/A as a postemergence broadcast application to soybeans. Apply in a minimum of 15 GPA. Application restricted to ground and aerial applications only.

**Spray Additives**

Add 1% v/v methylated seed oil (MSO) plus ammonium sulfate (AMS) at a minimum of 8.5 lbs/100 gals of finished spray volume.

Weeds Controlled Weed	Maximum Number of Leaves At Application
Cocklebur, common	2
Mustard, wild	4
Nightshade, black	2
Pigweed, redroot	4
Waterhemp, common	2
Smooth Pigweed	2
Common Ragweed	4, (suppression only)
Palmer Amaranth	2
Common Lambsquarter	2 (suppression only)

<b>Rotational Crop Restrictions Rotational Crops</b>	<b>Minimum Rotational Interval after SINISTER Application (Months)</b>
Bean, Dry Potato Soybean	0 months
Small Grains, such as Wheat, Barley, Rye	9 months
Edible-podded beans and peas not otherwise specified in this label Pea, Dry	12 months
Corn, Field* Corn, Seed*	18 months
All other crops not listed above	Minimum 18 months and a successful field bioassay (refer to <b>Field Bioassay Instructions</b> section) must be performed before planting rotational crop

\* For fields with greater than 4% organic matter, a successful field bioassay (refer to **Field Bioassay Instructions** section) must be performed before planting field corn or seed corn.

### **Field Bioassay Instructions**

Plant several strips of the desired crop variety across the field which has been previously treated with SINISTER. Plant the strips perpendicular to the direction SINISTER was applied. The strips should be located so that all the different field conditions are encountered, including differences in soil texture, organic matter, pH, and drainage. If the crop does not show visible symptoms of injury and/or stand reduction, this field can be planted to this crop. If visible injury and/or stand reduction occur, do not plant the field to this crop; plant only a labeled crop.

### **Use Restrictions**

- Do not exceed 0.49 pt/A (0.176 lb ai/A) of SINISTER per acre in any one year. Do not apply to any field more than once every two years.
- Do not make a SINISTER application after June 30.
- Do not harvest soybeans within 45 days of application.
- Avoid overlapping spray swaths, as injury may occur to rotational crops.

### **24(c) Registrant:**

Helena Agri-Enterprises, LLC  
225 Schilling Boulevard, Suite 300  
Collierville, TN 38017

SINISTER is a trademark of Helena Holding Company