



Bayer CropScience LP
P.O. Box 12014
2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)

SIVANTO™ prime
EPA Reg. No. 264-1141

Reduced application rates for soil applications in Cucurbits.

PRODUCT BULLETIN

FOR DISTRIBUTION AND USE ONLY IN THE STATES OF FLORIDA AND GEORGIA

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or accepted by the U.S. Environmental Protection Agency. This product bulletin should be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

CUCURBIT VEGETABLES – SOIL

Crops of Crop Group 9 Including: Chayote (fruit), Chinese waxgourd (Chinese preserving melon), Citron melon, Cucumber, Gherkin, Gourd (edible, includes hyotan, cucuzza, hechima, Chinese okra), *Momordica* spp. (includes balsam apple, balsam pear, bitter melon, Chinese cucumber), Muskmelon (hybrids and/or cultivars of *Cucumis melo* including true cantaloupe, cantaloupe, casaba, Crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, snake melon), Pumpkin, Squash (includes summer squash types such as: crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini, and winter squash types such as acorn squash, butternut squash, calabaza, cushaw, Hubbard squash, spaghetti squash), Watermelon (includes hybrids and/or varieties of *Citrullus lanatus*)

Pests Controlled	Product Rate (fl oz/A)
Leafhoppers	14.0 -21.0
Aphids	14.0 -21.0
Whiteflies	14.0 -21.0

Soil Application Restrictions:

Pre-Harvest Interval (PHI): **21 days**

Maximum SIVANTO prime allowed per crop season: **28.0 fluid ounces/Acre** (0.365 lb AI/Acre), regardless of method of application.

Maximum crop seasons per year: **3**

Soil Application Notes:

SIVANTO prime may be applied at the specified dosage by the following methods:

1. chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment;
2. injection below the eventual seed-line prior to planting. Place SIVANTO prime 3-4 inches below seed-line;
3. potting hole drench at transplanting; or post-transplant drench following setting and covering.

For **MEDICAL** And **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577

For **PRODUCT USE** Information Call 1-866-99BAYER (1-866-992-2937)

As with any crop-protection product, always read and follow label directions.
For additional information call toll-free 1-866-99BAYER (1-866-992-2937)

Sivanto is a trademark of Bayer.