



FIFRA 2(ee) Recommendation

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF CALIFORNIA

SIVANTO® prime

EPA Reg. No. 264-1141

For the use of SIVANTO® prime by additional soil application methods at planting/transplanting in Brassica Head and Stem Vegetables and Kohlrabi, Cucurbit Vegetables, Fruiting Vegetables, and Leafy Vegetables

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its label.

BRASSICA HEAD AND STEM VEGETABLES AND KOHLRABI SOIL	
Crops of Group 5-16 - Broccoli; Brussels sprouts; Cabbage; Cabbage, Chinese, napa; Cauliflower; cultivars, varieties, and/or hybrids of these	
Pest Controlled	Product Rate (fl oz/A)
Aphids Leafhoppers Whiteflies	21.0 – 28.0
Application Information: SIVANTO prime may be applied at the specified dosage by the following methods: <ul style="list-style-type: none">• In-furrow spray directed on or below seed.• Post-seed drench, transplant-water drench or hill drench.	
Application Restrictions: <ul style="list-style-type: none">• Pre-Harvest Interval (PHI): 21 Day(s)• Do not apply more than 28 fl oz of SIVANTO prime (0.365 Pound Flupyradifurone) per Acre Per Crop Season on brassica (cole) leafy vegetables, regardless of method of application, product, or formulation.• Maximum number of crop seasons per year: 3	

CUCURBIT VEGETABLES (Group 9)
SOIL

Chayote (fruit); Chinese waxgourd (Chinese preserving melon); Citron melon; Cucumber; Gherkin; Gourd, edible (includes hyotan, cucuzza, hechima, Chinese okra); Momordica spp. (includes balsam apple, balsam pear, bitter melon, Chinese cucumber); Muskmelon (hybrids and/or cultivars of Cucumis melo including true cantaloupe, cantaloupe, casaba, Crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, and snake melon); Pumpkin; Squash, summer (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini); Squash, winter (includes acorn squash, butternut squash, calabaza, hubbard squash, spaghetti squash); Watermelon (includes hybrids and/or varieties of Citrullus lanatus)

Pest Controlled	Product Rate (fl oz/A)
Aphids	21.0 – 28.0
Leafhoppers	

Application Information:

SIVANTO prime may be applied at the specified dosage by the following methods:

- In-furrow spray directed on or below seed.
- Post-seed drench, transplant-water drench or hill drench.

A mild yellowing on leaf margins is sometimes seen following application of SIVANTO prime in cucurbits.

Application Restrictions:

- Pre-Harvest Interval (PHI): **21 Day(s)**
- Do not apply more than **28 fl oz** of SIVANTO prime (**0.365 Pound Flupyradifurone**) per **Acre Per Crop Season** on cucurbit vegetables, regardless of method of application, product, or formulation.
- Maximum number of crop seasons per year: **3**

FRUITING VEGETABLES (Group 8-10)
SOIL

African eggplant; Bush tomato; Cocona; Currant tomato; Eggplant; Garden huckleberry; Goji berry; Groundcherry; Martynia; Naranjilla; Okra; Pea eggplant; Pepino; Pepper, bell; Pepper, nonbell; Roselle; Scarlet eggplant; Sunberry; Tomatillo; Tomato; Tree tomato; cultivars, varieties and/or hybrids of these

Pest Controlled	Product Rate (fl oz/A)
Aphids	21.0 – 28.0
Leafhoppers	
Psyllids	
Whiteflies	

Application Information:

SIVANTO prime may be applied at the specified dosage by the following methods:

- In-furrow spray directed on or below seed.
- Post-seeding drench, transplant-water drench or hill drench.

Application Restrictions:

- Pre-Harvest Interval (PHI): **45 Day(s)**
- Do not apply more than **28 fl oz** of SIVANTO prime (**0.365 Pound Flupyradifurone**) per **Acre Per Crop Season** on fruiting vegetables, regardless of method of application, product or formulation.
- Maximum number of crop seasons per year: **3**

LEAFY VEGETABLE (Group 4-16)**SOIL**

Amaranth, Chinese; Amaranth, leafy; Arugula; Aster, Indian ; Blackjack ; Broccoli raab ; Broccoli, Chinese ; Cabbage, abyssinian; Cabbage, Chinese, bok choy; Cabbage, seakale; Cat's whiskers; Cham-chwi ; Cham-na-mul; Chervil, fresh leaves ; Chipilin ; Chrysanthemum, garland; Cilantro, fresh leaves; Collards; Corn salad; Cosmos ; Cress, garden; Cress, upland; Dandelion, leaves ; Dang-gwi, leaves ; Dillweed ; Dock ; Dol-nam-mul ; Ebolo ; Endive ; Escarole; Fameflower ; Feather cockscomb ; Good King Henry ; Hanover salad ; Huauzontle ; Jute, leaves ; Kale; Lettuce, bitter ; Lettuce, head; Lettuce, leaf ; Maca, leaves ; Mizuna; Mustard greens; Orach; Parsley, fresh leaves ; Plantain, buckthorn ; Primrose, English ; Purslane, garden; Purslane, winter; Radicchio ; Radish, leaves ; Rape greens; Rocket, wild ; Shepherd's purse ; Spinach; Spinach, Malabar ; Spinach, New Zealand; Spinach, tanier ; Swiss chard; Turnip greens; Violet, Chinese, leaves; cultivars, varieties, and/or hybrids of these

Pest Controlled	Product Rate (fl oz/A)
Aphids	21.0 – 28.0
Leafhoppers	
Whiteflies	

Application Information:

SIVANTO prime may be applied at the specified dosage by the following methods:

- In-furrow spray directed on or below seed.
- Post-seeding drench, transplant-water drench, or hill drench.

Application Restrictions:

- Pre-Harvest Interval (PHI): **21 Day(s)**
- Do not apply more than **28 fl oz** of SIVANTO prime (**0.365 Pound Flupyradifurone**) per **Acre Per Crop Season** on leafy vegetables, regardless of method of application, product, or formulation.
- Maximum number of crop seasons per year: 3

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937).

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937).

Bayer CropScience LP
800 N. Lindbergh Blvd.
St. Louis, MO 63167