



Bayer CropScience LP  
800 N. Lindbergh Blvd.  
St. Louis, MO 63167  
1-866-99BAYER (1-866-992-2937)

## SIVANTO® prime

Reduced rates in Pearl Millet and Sorghum  
for control of Aphids

EPA Reg. No. 264-1141

### ***FIFRA 2(ee) Recommendation***

**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE(S) OF ALABAMA,  
ARIZONA, ARKANSAS, COLORADO, FLORIDA, GEORGIA, KANSAS,  
KENTUCKY, LOUISIANA, MISSISSIPPI, NEBRASKA, NEW MEXICO, NORTH  
CAROLINA, OKLAHOMA, SOUTH CAROLINA, TENNESSEE, TEXAS AND  
VIRGINIA**

This recommendation is valid until December 31, 2025

**FIFRA Section 2(ee) Recommendation:** This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label(s) are to be followed.

#### **DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its label(s).

<b>MILLET, PEARL; SORGHUM (INCLUDING SORGHUM GROWN FOR SYRUP) – Foliar Application</b>	
<b>Pest Controlled</b>	<b>Product Rate (fl oz/A)</b>
Aphids	4.0 – 7.0
<b>Application Restrictions:</b> Pre-Harvest Interval (PHI): <b>7 Day(s)</b> – hay, forage, sorghum grown for syrup, or sweet corn Pre-Harvest Interval (PHI): <b>21 Day(s)</b> - dried grain, stover or straw Minimum interval between applications: <b>7 Day(s)</b> Minimum Application Volume: <b>10 Gallons/Acre</b>   Ground; <b>3 Gallons/Acre</b>   Aerial Do not apply more than <b>28 fl oz</b> of SIVANTO prime ( <b>0.365 Pound Flupyradifurone</b> ) per <b>Acre Per Calendar Year</b> on cereal grains, regardless of method of application, product or formulation.	

SORGHUM (INCLUDING SORGHUM GROWN FOR SYRUP) – Soil Application	
Pest Controlled	Product Rate (fl oz/A)
Aphids	6.0 – 10.0
<b>Application Information:</b> SIVANTO prime may be applied at the specified dosage by the following methods: <ul style="list-style-type: none"> <li>• Injection below the eventual seed row prior to planting. Place SIVANTO prime 3 - 4 inches below seed-line</li> <li>• In-furrow spray during planting</li> <li>• Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment</li> </ul>	
<b>Application Restrictions:</b> Pre-Harvest Interval (PHI): <b>45 Day(s)</b> Do not apply more than <b>28 fl oz</b> of SIVANTO prime ( <b>0.365 Pound Flupyradifurone</b> ) per <b>Acre Per Calendar Year</b> on Sorghum, regardless of method of application, product or formulation. Maximum number of crop seasons per year: <b>1</b>	

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.  
For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937).

*As with any crop-protection product, always read and follow label directions.  
For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*