

FIFRA Section 2(ee) Recommendation

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the U.S. Environmental Protection Agency

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF ALABAMA, TENNESSEE, LOUSIANA, FLORIDA, GEORGIA, MISSISSIPPI, VIRGINIA, NORTH CAROLINA AND SOUTH CAROLINA

SNIPER Insecticide/Miticide EPA REG NO. 34704-858

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

SOYBEANS			
Doots	DOSAGE		DEMARKS
Pests	LB AI/A	FL OZ/A	REMARKS
Kudzu bug	0.10	6.4	 Apply in a minimum of 10 gallons per acre with ground equipment or 2 gallons per acre by aircraft at a minimum of 30 days intervals. Do not apply more than 0.3 lb ai per acre per season. Do not apply within 18 days of harvest. Utilize an intensive scouting program for proper application timing. Repeat applications may be necessary.

COTTON			
Pests	DOSAGE		DEMARKS
	LB Al/A	FL OZ/A	REMARKS
Kudzu bug	0.10	6.4	 Apply in a minimum of 5 gallons per acre with ground equipment or 1 gallon per acre by aircraft. Do not apply more than 0.5 lb ai per acre per season. Do not apply within 14 days of harvest. Utilize an intensive scouting program for proper application timing. Repeat



FIFRA Section 2(ee) Recommendation

		applications may be necessary.
	•	Do not graze livestock in treated area or
		cut treated crops for feed.

Potato			
Dooto	DOSAGE		DEMARKS
Pests	LB Al/A	FL OZ/A	REMARKS
Kudzu bug	0.10	6.4	 Apply in a minimum of 10 gallons per acre with ground equipment or 3 gallons per acre by aircraft. Do not make more than 2 foliar applications per season no sooner than 21 days apart. Do not apply more than 0.5 lb ai per acre per season. Do not apply within 21 days of harvest. Utilize an intensive scouting program for proper application timing. Repeat applications may be necessary.

Succulent Peas and Beans			
Pests	DOSAGE		REMARKS
resis	LB AI/A	FL OZ/A	REIWARNS
Kudzu bug	0.10	6.4	 Apply in a minimum of 10 gallons per acre with ground equipment or 2 gallons per acre by aircraft. When applying by air, 1-2 qts of emulsified oil may be substituted for 1-2 qts of water in the finished spray. Thorough coverage is essential to achieve control. Do not apply more than 0.2 lb ai per acre per season. Do not apply within 3 days of harvest. Utilize an intensive scouting program for proper application timing. Repeat applications may be necessary.

Loveland Products, Inc P.O. Box 1286 Greeley, CO 80632