

SPECIMEN LABEL

Agri Star®

MYCLOBUTANIL GROUP 3 FUNGICIDE

Agri Star® Sonoma® 40WSP

Fungicide

ACTIVE INGREDIENT

Myclobutanil: a-butyl-a-(4-chlorophenyl)-1H-1,2,4, triazole-1-propanenitrile . . . 40%

OTHER INGREDIENTS: 60%

TOTAL: 100%

EPA Reg. No. 42750-141

EPA Est. No. 33967-NJ-001

**KEEP OUT OF REACH OF CHILDREN
WARNING AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

You may also contact CHEMTREC toll free at 1-800-424-9300 day or night, for emergency treatment information.

Manufactured For:

ALBAUGH, LLC
1525 NE 36th Street
Ankeny, Iowa 50021

See inside booklet for additional
PRECAUTIONARY STATEMENTS
and complete **DIRECTIONS FOR USE.**



PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

Causes substantial but temporary eye injury. Harmful if inhaled, swallowed or absorbed through skin. Do not get in eyes or on clothing. Wear goggles, face shield, or safety glasses. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

1. Long-sleeved shirt and long pants
2. Chemical-resistant gloves made of any waterproof material
3. Shoes plus socks
4. Protective eyewear (goggles, face shield, or safety glasses)

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/ maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENTS:

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

Water soluble packets, when used correctly, qualify as a closed mixing/loading system under the Worker Protection Standard [40 CFR 170.607(d)]. Mixers and loaders handling this product while it is enclosed in intact water soluble packets may elect to wear reduced PPE of long-sleeved shirt, long pants, shoes, socks and chemical-resistant gloves.

When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for "applicators and other handlers" and have such PPE immediately available for use in an emergency, such as a spill or equipment break-down."

USER SAFETY RECOMMENDATIONS

Users should:

1. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
2. Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
3. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible wash thoroughly and change into clean clothes.

ENVIRONMENTAL HAZARDS

Do not apply directly to water or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift or runoff from areas treated.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Restriction: Do not apply this product in greenhouses.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

1. Coveralls
2. Chemical-resistant gloves made of any waterproof material
3. Shoes plus socks
4. Protective eyewear (goggles, face shield, or safety glasses)

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool dry area above freezing. The water-soluble pouch may become brittle at storage temperatures below 32°F, but the fungicide is not affected. Do not remove the water-soluble pouches from the container except for immediate use.

PESTICIDE DISPOSAL: Wastes resulting in the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING.

Non-refillable container. Do not reuse or refill this container. Triple or pressure rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

CONTAINER HANDLING:

Water-soluble pouch: Do not open the water soluble pouches. Put the entire bag into the spray tank.

Outer bag: Non-refillable container. Do not reuse or refill this container. Offer for recycling if available, or dispose of the empty outer bag in the trash as long as water-soluble pouches are unbroken. If the outer bag is contaminated, shake all residue into the application equipment and dispose of the empty bag according to the instructions in the pesticide disposal section or by procedures allowed by federal, state and local regulations.

Steps to be Taken in Case Material is Released or Spilled: Wear eye protection. Wear protective clothing. Spray water on powder and dust. Scoop or shovel solid material into a suitable container for recovery or disposal. Keep dust to a minimum. Flush contaminated area with a large amount of water to a chemical or sanitary sewer containing a settling pit. Remove contaminated clothing promptly and wash affected skin areas with soap and water. Thoroughly launder clothing before reuse. Refer to Precautionary Statements promptly and wash affected skin areas with soap and water.

PRODUCT USE INFORMATION

Read all **Directions for Use** carefully before applying.

AGRISTAR® SONOMA® 40WSP fungicide is a systemic, protectant and curative fungicide for the control of specific diseases mentioned on this label. Optimum disease control is achieved when the fungicide is applied in a regularly scheduled preventative spray program.

RESISTANCE MANAGEMENT

MYCLOBUTANIL	GROUP	3	FUNGICIDE
--------------	-------	----------	-----------

For Resistance Management, AGRISTAR® SONOMA® 40WSP is a systemic fungicide that contains myclobutanil, a Group 3 fungicide.

Any fungal population may contain individuals naturally resistant to AGRISTAR® SONOMA® 40WSP and other Group 3 fungicide. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields.

Because resistance development cannot be predicted, the use of this product should follow appropriate resistance-management strategies. To delay fungicide resistance, use should be based on an IPM program that includes the following steps:

- Rotate the use of AGRISTAR® SONOMA® 40WSP with other fungicide Groups within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicide from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information or to report suspected resistance contact Albaugh, LLC at **1-800-247-8013**. You can also contact your pesticide distributor or university extension specialist to report resistance.

Handling

AGRISTAR® SONOMA® 40WSP is packaged in water-soluble pouches. Do not allow pouches to become wet prior to adding to the spray tank. Do not handle the pouches with wet hands or wet gloves. Always reseal overwrap bag to protect remaining unused pouches. Do not remove water-soluble pouches from overwrap except to add directly to the spray tank.

Use Rate Determination

Carefully read, understand and follow label use rates and restrictions. Under low disease conditions, minimum label use rates per application can be used while maximum label rates and shortened spray schedules are for severe or threatening disease conditions. For proper application, determine the number of acres to be treated, the specified label use rate and the gallonage to be applied per acre. Prepare only the amount of spray solution required to treat the measured acreage. Careful calibration of spray equipment is recommended prior to use.

Rotational Crops Restriction

Crops on this label may be planted immediately after the last treatment.

Do not plant any other crops not listed for use on this label within 30 days after the last application.

Application

Ground:

Thorough coverage sprays generally result in optimum disease control. To achieve good coverage use proper spray pressure, gallonage per acre, nozzles, nozzle spacing and tractor speed. Consult spray nozzle and accessory catalogues for specific information on proper equipment calibration.

Aerial:

Apply in a minimum of 10 gallons of water per acre. Avoid application under conditions when uniform coverage cannot be obtained or when excessive spray drift may occur.

CHEMIGATION USE DIRECTIONS

Sprinkler Irrigation

AGRI STAR® SONOMA® 40WSP must be applied on a regular protectant fungicide schedule, not an irrigation schedule. If irrigation cycles are less frequent than the specified application intervals for AGRI STAR® SONOMA® 40WSP, ground or aerial applications must supplement chemigation applications to achieve adequate disease control.

- Apply AGRI STAR® SONOMA® 40WSP only through sprinkler irrigation systems including center-pivot, lateral move, end tow, side (wheel) roll, traveler, solid set, or hand move irrigation systems. Do not apply product through any other type of irrigation system.
- Lack of fungicidal effectiveness or illegal pesticide residues in the crop can result from nonuniform distribution of treated water.
- If you have questions about calibration, you should contact State Extension Service specialist, equipment manufacturers or other experts.
- Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water system are in place.
- A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Before applying AGRI STAR® SONOMA® 40WSP through sprinkler irrigation equipment, the chemigation system must meet the following specifications:

- Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
- Chemigation systems connected to public water systems must contain a functional reduced-pressure zone (RPZ), backflow preventer or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

(continued)

- Systems not connected to a public water supply must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located in the irrigation pipeline to prevent water source contamination from back flow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
- The irrigation line or water pump must include a functional pressure switch that will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- Do not apply when wind speed favors drift beyond the area intended for treatment.

Center-pivot, Lateral Move, End Tow, and Traveler Irrigation Equipment (use only with electric or oil hydraulic drive systems, which provide a uniform water distribution):

- Determine size of area to be treated.
- Determine the time required to apply no more than 1/4 inch water (6,750 gallons water per acre) over the area to be treated when the system and injection equipment are operated at normal pressures recommended by the equipment manufacturer. Run system at 80 to 95% of manufacturer's rated capacity.
- Using only water, determine the injection pump output when operated at normal line pressure.
- Determine the amount of AGRI STAR® SONOMA® 40WSP required for treatment area.
- Add the required amount of AGRI STAR® SONOMA® 40WSP and sufficient water to meet the injection time requirements of the solution tank.
- Maintain constant solution tank agitation during the injection period.
- Stop injection equipment after treatment is completed. Continue to operate the system until AGRI STAR® SONOMA® 40WSP solution has cleared the sprinkler head.

Solid-set, Side (wheel) Roll, and Hand Move Irrigation Equipment:

- Determine acreage covered by sprinkler.
- Fill injector solution tank with water and adjust flow rate to use contents over a 10 to 30 minute interval.
- Determine the amount of AGRI STAR® SONOMA® 40WSP required for treatment area.
- Add the required amount of AGRI STAR® SONOMA® 40WSP into the same quantity of water used to calibrate the injection equipment.
- Maintain constant solution tank agitation during the injection period.
- Operate system at normal pressures recommended by the manufacturer of the injection equipment and used for the time interval established during calibration.
- Inject AGRI STAR® SONOMA® 40WSP at the end of the irrigation cycle or as a separate application to maximize foliar fungicide retention.
- Stop injection equipment after treatment is completed. Continue to operate the system until AGRI STAR® SONOMA® 40WSP solution has cleared the last sprinkler head.

Instructions for Using Water Soluble Packages Directly into Spray tanks:

Water Soluble Packages (WSPs) are designed to dissolve in water. Agitation may be used, if necessary, to help dissolve the WSP. Failure to follow handling and mixing instructions can increase your exposure to the pesticide products in WSPs. WSPs, when used properly, qualify as a closed mixing/loading system under the Agricultural Worker Protection Standard [40 CFR 170.607(d)].

Handling Instructions

Follow these steps when handling pesticide products in WSPs.

1. Mix in spray tank only.
2. Handle WSP(s) in a manner that protects package from breakage and/or unintended release of contents. If package is broken, put on PPE required for clean-up and then continue with mixing instructions.
3. Keep the WSP(s) in outer packaging until just before use.
4. Keep the WSP dry prior to adding to the spray tank.
5. Handle with dry gloves and according to the label instructions for PPE.
6. Keep WSP intact. Do not cut or puncture WSP.
7. Reseal the WSP outer packaging to protect any unused WSP(s).

Mixing

Always place AGRI STAR® SONOMA® 40WSP into solution prior to adding co-applied agricultural chemicals registered for use on specific crop or crops being treated. Add the required number of unopened pouches as determined by the dosage directions into the spray tank with agitation. Depending on the water temperature and the degree of agitation, the pouches should dissolve completely within approximately ten minutes from the time they are added to the water.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Mixing Instructions

Follow the steps below when mixing this product, including if tank mixed with other pesticide products. If being tank mixed, the mixing directions 1 through 9 below take precedence over the mixing directions of the other tank mix products.

WSPs may, in some cases, be mixed with other pesticide products so long as the directions for use of all mixed products do not conflict. Do not tank mix this product with products that prohibit tank mixing or have conflicting mixing directions.

1. If a basket or strainer is present in the tank hatch, remove prior to adding the WSP to the tank.
2. Fill tank with water to approximately one-third to one-half of the desired final volume of spray.
3. Stop adding water and stop any agitation.
4. Place intact/unopened WSP(s) into the tank.
5. Do not spray water from a hose or fill pipe to break or dissolve the WSP(s).
6. Start mechanical and recirculation agitation from the bottom of tank without using any overhead recirculation, if possible. If overhead recirculation cannot be turned off, close the hatch before starting agitation.
7. Dissolving the WSP(s) may take up to 5 minutes or longer, depending on water temperature, water hardness and intensity of agitation.
8. Stop agitation before tank lid is opened.
9. Open the lid to the tank, exercising caution to avoid contact with dusts or spray mix, to verify that the WSPs have fully dissolved and the contents have been thoroughly mixed into the solution.
10. Do not add other allowed products or complete filling the tank until the bags have fully dissolved and pesticide is thoroughly mixed.
11. Once the WSP have fully dissolved and any other products have been added to the tank, resume filling the tank with water to the desired level, close the tank lid, and resume agitation.
12. Use the spray solution when mixing is complete.
13. Maintain agitation of the diluted pesticide mix during transport and application.
14. It is unlawful to use any registered pesticide, including WSPs, in a manner inconsistent with its label.

Compatibility

AGRI STAR® SONOMA® 40WSP is compatible with most commonly used agricultural fungicides, insecticides, growth regulators, micronutrients and spray adjuvants. When preparing tank mixes, user should consult spray compatibility charts or State Cooperative Extension Service Specialists prior to actual use.

Note: AGRI STAR® SONOMA® 40WSP is compatible with boron and spray oils; however, the water-soluble pouches must be completely dissolved before adding spray oils or products containing boron to spray mixtures.

USE DIRECTIONS FOR TREE FRUITS

Best control of labeled diseases is achieved when AGRI STAR® SONOMA® 40WSP is applied on a 7 to 10 day application schedule.

AGRI STAR® SONOMA® 40WSP is a systemic fungicide and does not redistribute with rainfall after application. Application equipment spray nozzles should be adjusted to apply a uniform spray throughout the entire tree canopy.

The following use rates are to be used in determining the amount of AGRI STAR® SONOMA® 40WSP to be used per 100 gallons spray or per acre. Refer to specific tree fruit use directions to determine actual use rates for the control of labeled diseases.

Determination of Use Rates on an Acre Basis

The amount of AGRI STAR® SONOMA® 40WSP required per acre varies with tree size and the volume of fruit and foliage to be treated. The following summary table may be used as additional guidance for the determination of appropriate per acre use rates for AGRI STAR® SONOMA® 40WSP:

TREE HEIGHT (ft)	AGRI STAR® SONOMA® 40WSP (oz/acre)	
	Apples and Mayhaws	Stone Fruits
10 (or less)	2.5 to 5.0	2.5 to 4.0
15	3.75 to 6.0	4.0 to 6.0
20 (or more)	5.0 to 10.0	6.0

Concentrate Spray Applications: AGRI STAR® SONOMA® 40WSP may be used at the specified use rate per acre in either dilute or concentrate sprays. Use the following formula to determine the equivalent amount of product per acre in 2X, 3X, etc. spray solutions:

$$\frac{\text{Ounces AGRI STAR® SONOMA® 40WSP per acre} \times 100}{\text{Gallons of spray applied per acre}} = \text{Ounces of AGRI STAR® SONOMA® 40WSP per 100 gallons}$$

Example: An apple orchard consisting of apple trees 18 feet in height will require 5 ounces AGRI STAR® SONOMA® 40WSP for adequate apple scab control. Application equipment has been calibrated to apply 80 gallons spray per acre, therefore:

$$\frac{5 \text{ ounces AGRI STAR® SONOMA® 40WSP per acre} \times 100}{80 \text{ gallons spray applied per acre}} = 6.25 \text{ ounces per 100 gallons}$$

Dilute, Thorough Coverage Applications: Dilute thorough coverage applications are based on the amount of spray solution required to thoroughly wet trees until spray run-off. The following specific use directions for apples and mayhaws utilizes a 400 gallon per acre dilute basis and the specific use directions for stone fruits utilize a 250 gallon per acre dilute basis.

Thorough spray coverage is essential for good disease control. AGRI STAR® SONOMA® 40WSP should be applied in sufficient water to ensure adequate coverage.

USE DIRECTIONS FOR ALMONDS
(For Use in California, Arizona and Hawaii Only)

DISEASES	RATE OF AGRISTAR® SONOMA® 40WSP		APPLICATION DIRECTIONS++	RESTRICTIONS
	(oz/100 gal)+	(oz/acre)		
Blossom Blight (Monilinia spp.)	1.25 to 2.0 oz. (0.03125 – 0.05 lbs. ai)	5.0 to 8.0 oz. (0.125 – 0.20 lbs. ai)	Begin applications at pink bud stage (about 5% bloom). If conditions are favorable for disease development, apply again at full bloom and at petal fall.	Do not apply within 90 days of harvest. Do not apply more than 8 oz product (0.20 lb. a.i.) per acre per application. Do not make more than three applications or apply more than 24 oz. (0.60 lbs. ai) of AGRISTAR® SONOMA® 40WSP fungicide product per acre per year. Do not apply on an application schedule exceeding 21 days.
Shothole (Stigmina spp.)			Apply 6 ounces (.15 lbs. a.i.) of AGRISTAR® SONOMA® 40WSP product per acre. Begin application approximately 8 weeks after flowering if environmental conditions are favorable for disease development.	
Rust (Tranzschelia spp.)				
Anthrachnose (Colletotrichum spp.)			The application of AGRISTAR® SONOMA® 40WSP fungicide for control of blossom blight and shothole will suppress anthracnose.	

+Based on a standard of 400 gallons of dilute spray per acre, or the equivalent amount of product per acre.
++Best disease control is achieved in thorough coverage sprays applied on a protectant schedule, which does not exceed 10 days.

USE DIRECTIONS FOR ARTICHOKE

DISEASES	RATE OF AGRISTAR® SONOMA® 40WSP (OZ/ACRE)	APPLICATION DIRECTIONS	RESTRICTIONS
Powdery mildew (<i>Erysiphe Cichoracearum</i>)	4 oz. (0.10 lb. ai)	<p>Scout crops on a regular basis and treat when disease first appears or when conditions favor disease development.</p> <p>Apply uniformly in a spray volume that provides thorough coverage. Use a minimum of 30 gallons of spray volume per acre. Control may be reduced at low spray volumes or if spray coverage is not adequate.</p> <p>Consult your Albaugh representative, extension specialist, certified crop advisor or your state agricultural experiment station or any additional local use recommendations for your area.</p>	<p>Do not apply more than 4 oz. of AGRISTAR® SONOMA® 40WSP per acre per application or 24 oz. (0.60 lbs. ai) of AGRISTAR® SONOMA® 40WSP per acre per year. Do not make more than 6 applications per year.</p> <p>Pre-harvest Interval: Do not harvest within 3 days of application. Re-treatment Interval: 14 days</p> <p>Rotational Crop Restriction: Crops with registered uses may be planted immediately after the last treatment.</p> <p>Do not plant other crops within 30 days after the last application.</p>

USE DIRECTIONS FOR GRAPES

DISEASES	RATE OF AGRI STAR® SONOMA® 40WSP (OZ/ACRE)	APPLICATION DIRECTIONS	RESTRICTIONS
Anthracnose (Elsinoe spp.)	3 to 5 oz. (0.125 lbs. ai)	Begin application when new shoots are 1 to 3 inches in length and continue on an application schedule, which does not exceed 14 days.	Do not apply within 14 days of harvest. Do not apply more than 5 oz. product (0.125 lb. a.i.) per acre per application.
Black Rot (Guignardia spp.)		Preventative Schedule: Begin application when new shoots are 1 to 3 inches in length and continue applications on an application schedule which does not exceed 14 days. Use higher specified rate under heavy disease pressure. Post Infection Schedule: Apply within 72 hours after the beginning of an infection period.	Do not apply more than 20 ounces of AGRI STAR® SONOMA® 40WSP product (0.50 lbs. ai) per acre per year. Do not make more than 4 applications per year.
Powdery Mildew (Uncinula spp.)		Begin application at prebloom (12 to 18-inch shoots) and do not extend applications beyond a 21-day interval. Use higher specified rate or shorter spray interval on susceptible varieties or under heavy disease pressure.	

Aid in the Control of Certain Vine Diseases Following Pruning in Grapevines

To protect pruning wounds from vine diseases caused by *Botryosphaeria rhodina*, *Eutypa lata*, *Phaeoacremonium aleophilum* or *Phaeomoniella chlamydospora*, apply AGRI STAR® SONOMA® 40WSP at 5 oz per acre in 50 gallons of water per acre using power operated ground application equipment. Lower application volumes may be used only if the spray thoroughly wets all susceptible grapevine tissue and the same ratio of AGRI STAR® SONOMA® 40WSP to water is maintained: 4 oz. of AGRI STAR® SONOMA® 40WSP per acre in 42 gallons of water, or 3 oz of AGRI STAR® SONOMA® 40WSP per acre in 33 gallons of water per acre. Control may be reduced at these lower application rates. Do not apply less than 4 oz of AGRI STAR® SONOMA® 40WSP per acre per application.

Apply as a directed spray immediately after pruning (within 24 hours). Assure thorough coverage of cordons, spurs and all cut wood surfaces. A second application approximately two weeks later is recommended. A second application is necessary if rainfall occurs or if humid conditions persist.

Double pruning involves two pruning passes. Canes first are cut non-selectively to a uniform height. Later, selective pruning reduces canes to their final spur length. When conditions do not favor infections developing beyond where the final pruning cuts will be made, the first pass pruning cuts do not need to be treated. In this case, apply AGRI STAR® SONOMA® 40WSP immediately after the second pruning only. Make a second application approximately two weeks later if rainfall occurs or is expected, or if conditions favor spore dispersal and germination. If there is risk of infection moving beyond the second set of pruning cuts, applications should be made after the first and second prunings.

The addition of a labeled rate of a registered organosilicone spray adjuvant may increase penetration into cut wood surfaces. It is the responsibility of the user to assure that the organosilicone spray adjuvant is safe to the crop under the existing conditions of use. Addition of a registered spray dye to the tank mix, and visual inspection of pruning cuts after application, is recommended to assure thorough coverage of all susceptible tissue.

Specific Use Restrictions:

- Do not apply more than 24 oz. of AGRISTAR® SONOMA® 40WSP (1.50 lbs. ai) per acre per year including these applications and applications for powdery mildew control.

USE DIRECTIONS FOR HOPS
For Use and Distribution Only in the States of Alaska, Idaho, Montana, Nevada, Oregon, Utah, Washington, and Wyoming)

DISEASE	AGRISTAR® SONOMA® 40WSP (OZ/ACRE)	APPLICATION DIRECTIONS	RESTRICTIONS
Powdery Mildew (Uncinula spp.)	2 – 4 oz. (0.05 to 0.10 lbs. ai)	Apply every 7 - 10 days during emergence to training growth stage.	Do not make more than 4 applications of AGRISTAR® SONOMA® 40WSP product per acre per year. Do not apply less than 2 oz of AGRISTAR® SONOMA® 40WSP product per acre or adequate efficacy may not be achieved.
	4 – 6 oz. (0.10 – 0.15 lbs. ai)	Apply every 5 – 10 days during training to wire growth stage (prior to beginning of bloom when vines are rapidly growing).	Do not apply more than 40 oz. of AGRISTAR® SONOMA® 40WSP product (1.0 lb. a.i.) per acre per application. Pre-Harvest Interval: Do not apply within 14 days of harvest. Do not apply less than 4 oz. of AGRISTAR® SONOMA® 40WSP product per acre or adequate efficacy may not be achieved.
	6 – 10 oz. (0.15 - 0.25 lbs. ai)	Apply every 7 – 10 days during wire to 14 day pre-harvest stage.	Do not graze livestock in treated areas or harvest crops grown in treated areas for silage or hay. Do not apply less than 6 oz. of AGRISTAR® SONOMA® 40WSP product per acre or adequate efficacy may not be achieved. Chemigation: Do not apply through any type of irrigation system.

Apply AGRISTAR® SONOMA® 40WSP in sufficient water for thorough coverage using ground equipment or by air in a minimum spray volume of 10 gallons of water per acre. Thorough coverage is essential. Use the shorter spray interval on susceptible varieties or under disease pressure.

Resistance Management: AGRISTAR® SONOMA® 40WSP belongs to the sterol demethylation inhibitor (DMI) class of fungicides and is classified as a Group 3 Fungicide by EPA. Since certain fungi can develop resistance to this class of products, the use of AGRISTAR® SONOMA® 40WSP should be part of a resistance management strategy that includes alternation and/or tank mixing with fungicides of different modes of action. Consult your local or state agricultural authorities for resistance management strategies that are appropriate for your disease management program.

USE DIRECTIONS FOR PEPPERMINT AND SPEARMINT

DISEASES	RATE OF AGRI STAR® SONOMA® 40WSP (OZ/ACRE)	APPLICATION DIRECTIONS	RESTRICTIONS
Powdery Mildew (Erysiphe spp.) Rust (Puccinia spp.)	4 - 5 oz. (0.10 - 0.125 lbs. ai)	Begin application in early spring when plants break dormancy. Continue applications on a 14- to 21 -day application schedule.	Do not make more than three applications per year. Do not apply more than 5 oz. product (0.125 lb. a.i.) per acre per application. Do not apply more than 15 ounces of AGRI STAR® SONOMA® 40WSP product (0.375 lbs. ai) per acre per year. Do not apply within 30 days of harvest.

USE DIRECTIONS FOR APPLES AND MAYHAWES

DISEASES	RATE OF AGRISTAR® SONOMA® 40WSP (OZ/100 GAL) +	APPLICATION DIRECTIONS	RESTRICTIONS
Powdery Mildew (Podosphaera spp.)	1.25 to 2.5 oz. (0.03 to 0.06 lbs. ai)	Begin application at tight cluster and continue through the second cover spray. Additional sprays beyond second cover may be needed on susceptible varieties or under heavy disease pressure. Use high specified rate if powdery mildew was present in previous years.	Do not apply within 14 days of harvest. Do not make more than 4 applications of AGRISTAR® SONOMA® 40WSP product per acre per year.
Rusts (Gymnosporangium spp.)	1.25 to 2.0 oz. (0.03 to 0.05 lbs. ai)	Begin applications at pink stage and continue through the second cover spray.	Do not apply more than 2.5 oz product (0.06 lb a.i.) per acre per application.
Scab (Venturia spp.) Prebloom	1.25 to 2.0 oz. (0.03 to 0.05 lbs. ai)	Begin application at green tip or when environmental conditions become favorable for primary scab development. Apply AGRISTAR® SONOMA® 40WSP alone or tank mixed with a protectant fungicide on a 7- to 10-day schedule.	Do not apply more than 10 ounces of AGRISTAR® SONOMA® 40WSP product (0.24 lbs. ai) per acre per year.
Bloom, Postbloom	1.25 to 2.0 oz. (0.03 to 0.05 lbs. ai)	Use AGRISTAR® SONOMA® 40WSP in a tank mixture with the recommended rate of a protectant fungicide, registered for use on apples, for improved fruit scab and summer disease control	
Post Infection	2.0 oz. (0.05 lbs. ai)	AGRISTAR® SONOMA® 40WSP provides 96-hour post-infection control or curative activity. Apply as soon as possible after infection period. Follow with a standard preventative spray schedule.	

+ Based on dilute sprays with a 400-gallon per acre basis.

USE DIRECTIONS FOR BERRY AND SMALL FRUITS

DISEASES	AGRISTAR® SONOMA® 40WSP (OZ/ACRE)	APPLICATION DIRECTIONS	RESTRICTIONS
BLACKBERRY+			
Cane and Leaf Rust (Kuehneola spp.) Orange Rust (Arthuriomyces spp.) Powdery Mildew (Sphaerotheca spp.) Yellow Rust (Phragmidium spp.)	1.25 - 2.5 oz. (0.03 - 0.06 lbs. ai)	Applications should be initiated as early as bud break and should continue at 10- to 14-day intervals, depending on the disease(s) to be controlled. Use the shorter spray interval under heavy disease pressure.	Applications may be made up to the day of harvest. Do not make more than 4 applications of AGRISTAR® SONOMA® 40WSP product per acre per year. Do not apply more than 10 ounces of AGRISTAR® SONOMA® 40WSP product (0.25 lbs. ai) per acre per year.
CURRANT			
Powdery Mildew (Sphaerotheca spp.)	5.0 oz. (0.13 lbs. ai)	Make applications at pre-bloom, full bloom and 2 weeks later.	Applications may be made up to the day of harvest. Do not make more than 8 applications of AGRISTAR® SONOMA® 40WSP product per acre per year. Do not apply more than 40 ounces of AGRISTAR® SONOMA® 40WSP product (0.40 lb. ai) per acre per year.
GOOSEBERRY			
Anthracnose (Drepanopeziza spp.)	5.0 oz. (0.13 lbs. ai)	Begin applications when the first leaf has completely unfolded, then at 10- to 14-day intervals as long as environmental conditions favor continued disease development.	Applications may be made up to the day of harvest. Do not apply more than 5 oz product (0.13 lb a.i.) per acre per application. Do not make more than 8 applications of AGRISTAR® SONOMA® 40WSP product per acre per year.
Powdery Mildew (Sphaerotheca spp.)		Make applications at pre-bloom, full bloom and 2 weeks later.	Do not apply more than 40 ounces of AGRISTAR® SONOMA® 40WSP product (1.0 lb. ai) per acre per year.

(continued)

USE DIRECTIONS FOR BERRY AND SMALL FRUITS *(cont.)*

DISEASES	AGRI STAR® SONOMA® 40WSP (oz./acre)	APPLICATION DIRECTIONS	RESTRICTIONS
RASPBERRY			
Cane and Leaf Rust (Kuehneola spp.) Leaf Spot (Sphaeruiina spp.) Orange Rust (Arthuriomyces spp.) Powdery Mildew (Sphaerotheca spp.) Yellow Rust (Phragmidium spp.)	1.25 to 2.5 oz. (0.03 - 0.075 lbs. ai)	Applications should be initiated as early as bud break and should continue at 10- to 14-day intervals, depending on the disease(s) to be controlled. Use the shorter spray interval under heavy disease pressure.	Applications may be made up to the day of harvest. Do not make more than 4 applications of AGRI STAR® SONOMA® 40WSP product per acre per year. Do not apply more than 10 ounces of AGRI STAR® SONOMA® 40WSP (0.25 lbs. ai) per acre per year.
STRAWBERRY			
Powdery Mildew (Sphaerotheca spp.) Leaf Spot (Mycosphaerel/a spp.) Leaf Blight (Phomopsis spp.)	2.5 to 5.0 oz. (0.06 - 0.125 lbs. ai)	Begin applications when disease first appears or when conditions favor disease development. Repeat applications at 14- to 21 - day intervals.	Applications may be made up to the day of harvest. Do not make more than 4 applications of AGRI STAR® SONOMA® 40WSP product per acre per year. Do not apply more than 20 oz. (0.50 lbs. ai) of AGRI STAR® SONOMA® 40WSP per acre per year. Observe a 30-day plant-back interval between the last application and planting new crops at the treatment site.

+ Includes use on the following members of the berries group including varieties and/or hybrids:

bingleberry black satin berry boysenberry Cherokee blackberry Chesterberry Cheyenne blackberry Coryberry Darrowberry dewberry	Dirksen thornless berry Himalayaberry hullberry Lavacaberry lowberry Lucretiaberry mammoth blackberry marionberry Nectarberry	olallieberry Oregon evergreen berry phenomenalberry rangeberry ravenberry rossberry Shawnee blackberry youngberry
---	---	--

USE DIRECTIONS FOR STONE FRUIT

Stone Fruit - APRICOTS

DISEASES	(oz/acre)	APPLICATION DIRECTIONS	RESTRICTIONS
Brown Rot Blossom Blight (<i>Monilinia</i> spp.)	2.5 - 6.0 oz. (0.06 – 0.15 lbs. ai)	Begin application at early red bud stage before infection occurs. If conditions are favorable for disease development, apply again at full bloom and petal fall.	Do not apply more than 24 oz. of AGRI STAR® SONOMA® 40WSP product (0.60 lbs. ai) per acre per year. Do not make more than 4 applications of AGRI STAR® SONOMA® 40WSP product per acre per year. Applications may be made up to the day harvest
Brown Rot (<i>Monilinia</i> spp.)		Apply 6 ounces (2.4 oz active) per acre on a 7- to 14-day schedule. Applications should be made whenever environmental conditions favor disease development during the month prior to harvest.	
Powdery Mildew (<i>Podosphaera</i> spp.)		Follow brown rot blossom blight schedule making additional applications at 10- to 14-day intervals until terminal growth ceases.	
Shothole (<i>Stigmina</i> spp.)		Follow brown rot blossom blight schedule making additional applications at 7- to 10-day intervals as long as needed.	

Stone Fruit - CHERRY

DISEASES	(oz/acre)	APPLICATION DIRECTIONS	RESTRICTIONS
Brown Rot Blossom Blight (<i>Monilinia</i> spp)	2.5 - 6.0 oz. (0.06 – 0.15 lbs ai)	Begin application at early popcorn stage, before infection occurs. If conditions are favorable for disease development, apply again at full bloom and petal fall.	Do not make more than 4 applications of AGRI STAR® SONOMA® 40WS product per acre per year. Do not apply more than 6 oz. product (0.15 lb. a.i.) per acre per application. Do not apply more than 24 oz. (0.60 lbs. ai) of AGRI STAR® SONOMA® 40WS per acre per year.
Brown Rot (<i>Monilinia</i> spp)		Follow brown rot blossom blight schedule making additional applications at 10- to 14-day intervals until terminal growth ceases.	
Powdery Mildew (<i>Podosphaera</i> and <i>Sphaerotheca</i> spp)		Follow brown rot blossom blight schedule making additional applications at 7- to 10-day intervals as long as needed.	
Leaf Spot (<i>Blumeriella</i> spp)		Follow brown rot blossom blight schedule and continue applications at 7- to 10-day intervals. Make additional applications after harvest.	

Stone Fruit - Nectarines

DISEASES	(oz/acre)	APPLICATION DIRECTIONS	RESTRICTIONS
Brown Rot Blossom Blight (Monilinia spp.)	2.5 - 6.0 oz. (0.06 - 0.15 lbs. ai)	Begin application at early pink bud stage before infection occurs. If conditions are favorable for disease development, apply again at full bloom and petal fall.	Do not make more than 4 applications of AGRI STAR® SONOMA® 40WSP product per acre per year. Do not apply more than 6 oz product (0.15 lb a.i.) per acre per application. Do not apply more than 24 oz. (0.60 lbs. ai) of AGRI STAR® SONOMA® 40WSP per acre per year.
Brown Rot (Monilinia spp.)		Follow brown rot blossom blight schedule making additional applications at 10- to 14-day intervals until terminal growth ceases.	
Powdery Mildew (Podosphaera and Sphaerotheca spp.)		Follow brown rot blossom blight schedule making additional applications at 7- to 10-day intervals as long as needed.	
Shothole (Stigmina spp.)		Follow brown rot blossom blight schedule making additional applications at 7- to 10-day intervals as long as needed.	

Stone Fruit - Peaches

DISEASES	(oz/acre)	APPLICATION DIRECTIONS	RESTRICTIONS
Brown Rot Blossom Blight (Monilinia spp.)	2.5 - 6.0 oz. (0.06 - 0.15 lbs. ai)	Begin application at early pink bud stage before infection occurs. If conditions are favorable for disease development, apply again at full bloom and petal fall.	Do not make more than 4 applications of AGRI STAR® SONOMA® 40WSP product per acre per year. Do not apply more than 6 oz product (0.15 lb. a.i.) per acre per application. Do not apply more than 24 oz. (0.60 lbs. ai) of AGRI STAR® SONOMA® 40WSP per acre per year.
Brown Rot (Monilinia spp.)		Follow brown rot blossom blight schedule making additional applications at 10- to 14-day intervals until terminal growth ceases.	
Powdery Mildew (Podosphaera spp.)		Follow brown rot blossom blight schedule making additional applications at 7- to 10-day intervals as long as needed.	
Rust (Tranzschelia spp.)		Apply 6 ounces (2.4 oz.) per acre. Begin application approximately 8 weeks after flowering if environmental conditions are favorable for disease development.	

Stone Fruit - PLUMS, PRUNES

DISEASES	(oz/acre)	APPLICATION DIRECTIONS	RESTRICTIONS
Brown Rot Blossom Blight (Monilinia spp.)	2.5 - 6.0 oz. (0.06 - 0.15 lbs. ai)	Begin application at green tip, before infection occurs. If conditions are favorable for disease development, apply again at full bloom and petal fall.	Do not make more than 4 applications of AGRI STAR® SONOMA® 40WSP product per acre per year. Do not apply more than 6 oz product (0.15 lb a.i.) per acre per application. Do not apply more than 24 oz. (0.60 lbs. ai) of AGRI STAR® SONOMA® 40WSP per acre per year. Do not apply on an application schedule exceeding 21-days.
Brown Rot (Monilinia spp.)		Follow brown rot blossom blight schedule making additional applications at 7- to 10-day intervals as long as needed.	
Rust (Tranzschelia spp.)		Apply 6 ounces (2.4 oz active) per acre. Begin application approximately 8 weeks after flowering if environmental conditions are favorable for disease development.	

+ 250 gallon dilute spray per acre basis.

Tropical Fruits - Black Sapote, Canistel, Mamey Sapote, Mango, Papaya, Sapodilla, and Star Apple

DISEASES	AGRI STAR® SONOMA® 40WSP (OZ/ACRE)	APPLICATION DIRECTIONS	RESTRICTIONS
Powdery mildew (Oidium caricae)	10 oz. (0.25 lbs. ai)	Scout crops on a regular basis and treat when disease first appears or when conditions favor disease development. Apply uniformly in a spray volume that provides thorough coverage of the fruit and foliage. Control may be reduced at low spray volumes or if spray coverage is not adequate. Consult your Albaugh representative, extension specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.	Do not apply more than 10 oz. (0.25 lbs. ai) of AGRI STAR® SONOMA® 40WSP product per acre per application. Do not apply more than 80 oz. (2 lbs.) of AGRI STAR® SONOMA® 40WSP product per acre per year. Do not make more than 8 applications per year. Pre-harvest Interval: None Re-treatment Interval: 14 days Rotational Crop Restriction: Do not plant any crop other than those listed on the main label for 12 months following the last application of AGRI STAR® SONOMA® 40WSP Pre-harvest Interval: None Re-treatment Interval: 14 days Rotational Crop Restriction: Do not plant any crop other than those listed on the main label for 12 months following the last application of AGRI STAR® SONOMA® 40WSP product.

USE DIRECTIONS FOR VEGETABLES

Vegetables - ASPARAGUS

DISEASES	AGRISTAR® SONOMA® 40WSP (OZ/ACRE)	APPLICATION DIRECTIONS	RESTRICTIONS
Rust (Puccinia spp.)	5.0 oz. (0.125 lbs. ai)	Begin applications to the developing ferns after harvest has taken place. Repeat applications on a schedule not to exceed 14 days. Apply with a spray adjuvant, such as Latron B-1956 or Latron CS-7	<p>Do not apply to harvestable spears, and do not apply within 180 days of harvest in all states except California. In California, either a 30 or 180-day treatment to harvest interval may be used.</p> <p>When a 180-day treatment to harvest interval is used, do not make more than 6 applications of 5 oz. of AGRISTAR® SONOMA® 40WSP (0.125 lbs. ai) per acre per application, and a maximum of 30 ounces product (0.75 lbs. ai) per acre per year.</p> <p>If using a 30-day treatment to harvest interval, do not use more than 5 oz. (0.125 lbs. ai) per acre per application.</p> <p>Do not make more than 4 applications per year.</p> <p>Do not apply more than 20 oz. product (0.5 lbs. ai) per acre per year.</p>

Vegetables - CUCURBITS

Balsam apple, bitter melon, cantaloupe, casaba, Chinese cucumber, Chinese wax-gourd, citron melon, crenshaw melon, cucumber, golden pershaw melon, edible gourd, ornamental gourd, honey balls, honeydew melon, mango melon, Persian melon, pineapple melon, pumpkin, Santa Claus melon, snake melon, summer squash, winter squash, and watermelon

DISEASES	AGRI STAR® SONOMA® 40WSP (OZ/ACRE)	APPLICATION DIRECTIONS	RESTRICTIONS
Powdery Mildew (Erysiphe and Sphaerotheca spp.)	2.5 - 5 oz. (0.06 - 0.125 lbs. ai)	Begin application at first sign of disease development and continue on a 7- to 10-day application schedule. For the control of other foliar cucurbit diseases, co-applications of registered protectant fungicides should be made according to label use directions.	Do not apply more than 5 oz. of AGRI STAR® SONOMA® 40WSP product (0.125 lbs. ai) per acre per application. Do not apply more than 24 oz. product (0.6lbs. ai) per acre per year. Applications may be made up to and including the day of harvest. Observe a 30-day plant-back interval between the last application and planting new crops at the treatment site.

Vegetables – HEAD and LEAF LETTUCE

DISEASES	AGRI STAR® SONOMA® 40WSP (OZ/ACRE)	APPLICATION DIRECTIONS	RESTRICTIONS
Powdery mildew (<i>Erysiphe</i> <i>Cichoracearum</i>)	5 oz. (0.125 lbs. ai)	Scout crops on a regular basis and treat when disease first appears or when conditions favor disease development. Apply uniformly in a spray volume that provides thorough coverage. Control may be reduced at low spray volumes or if spray coverage is not adequate. Consult your Albaugh representative, extension specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.	Do not apply more than 5 oz. (0.125 lbs. ai) of AGRI STAR® SONOMA® 40WSP product per acre per application. Do not apply more than 20 oz. of AGRI STAR® SONOMA® 40WSP product (0.5 lbs. ai) per acre per year. Do not make more than 4 applications per year. Pre-harvest Interval: Do not harvest within 3 days of application. Re-treatment Interval: 14 days

Vegetables - Okra
(Not for Use In the States of Alaska, Arizona, California, Hawaii, Idaho, Montana, Nevada, Oregon, Utah, Washington, Wyoming)

DISEASE	AGRI STAR® SONOMA® 40WSP (OZ/ACRE)	APPLICATION DIRECTIONS	RESTRICTIONS
Powdery mildew (Sphaerotheca Fulginea)	2.5 – 5 oz. (0.06 - 0.125 lbs. ai)	<p>Scout crops on a regular basis and treat when disease first appears or when conditions favor disease development.</p> <p>Apply uniformly in a spray volume that provides thorough coverage of the fruit and foliage. Control may be reduced at low spray volumes or if spray coverage is not adequate.</p> <p>The lower rate and 14-day re-treatment interval may be used for small plants and under low disease pressure conditions.</p> <p>The highest specified rate and shorter re-treatment interval are recommended for larger plants and severe or threatening disease conditions. If reliable predictive modeling (risk index) systems area available, these can help to indicate disease pressure conditions.</p> <p>Consult your Albaugh representative, extension specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.</p>	<ul style="list-style-type: none"> • Do not apply more than 5 oz. (0.125 lbs. ai) of AGRI STAR® SONOMA® 40WSP product per acre per application. Do not apply more than 20 oz. of AGRI STAR® SONOMA® 40WSP product (0.5 lbs. ai) per acre per year. • Do not make more than 4 applications per year. • Pre-harvest Interval: None • Minimum Re-treatment Interval: 10 to 14 days

Vegetables – PEPPERS (Bell Pepper, Chili Pepper, Cooking Pepper, Pimento, Sweet Pepper) and EGGPLANT

(Not for Use In the States of Alaska, Arizona, California, Hawaii, Idaho, Montana, Nevada, Oregon, Utah, Washington, Wyoming)

DISEASE	AGRI STAR® SONOMA® 40WSP (OZ/ACRE)	APPLICATION DIRECTIONS	RESTRICTIONS
Powdery mildew (Leveillula taurica)	2.5 – 5 oz. (0.06 - 0.125 lbs. ai)	Scout crops on a regular basis and treat when disease first appears or when conditions favor disease development. Apply uniformly in a spray volume that provides thorough coverage of the fruit and foliage. Control may be reduced at low spray volumes or if spray coverage is not adequate. The lower rate and 14-day re-treatment interval may be used for small plants and under low disease pressure conditions. The highest specified rate and shorter re-treatment interval are recommended for larger plants and severe or threatening disease conditions. If reliable predictive modeling (risk index) systems are available, these can help to indicate disease pressure conditions. Consult your Albaugh representative, extension specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.	Do not apply more than 5 oz. (0.13 lbs. ai) of AGRI STAR® SONOMA® 40WSP product per acre per application. Do not apply more than 20 oz. of AGRI STAR® SONOMA® 40WSP product (0.5 lbs. ai) per acre per year. <ul style="list-style-type: none"> • Do not make more than 4 applications per year. • Pre-harvest Interval: None • Minimum Re-treatment Interval: 10 to 14 days

Vegetables - SNAP BEANS

DISEASES	AGRI STAR® SONOMA® 40WSP (OZ/ACRE)	APPLICATION DIRECTIONS	RESTRICTIONS
Rust (Uromyces spp.) Pod Tip Rot (Rhizoctonia spp.)	4 to 5 oz. 0.10 - 0.125 lbs. ai)	Begin applications when rust is first observed. For pod tip rot, begin applications when pods begin to develop. Continue applications on a 7- to 10-day schedule if conditions remain favorable for disease development.	Do not apply more than 5 oz (0.125 lbs. ai) of AGRI STAR® SONOMA® 40WSP product per acre per application. Do not apply more than 25 oz. of AGRI STAR® SONOMA® 40WSP product (0.65 lbs. ai) per acre per year. Observe a 30-day plant-back interval between the last application and planting new crops at the treatment site. Preharvest Interval (PHI) = 0 days Do not make more than five applications per year

Vegetables - TOMATOES

DISEASES	AGRISTAR® SONOMA® 40WSP (OZ/ACRE)	APPLICATION DIRECTIONS	RESTRICTIONS
Powdery Mildew (Leveillua spp.)	2.5 - 4.0 oz. (0.06 - 0.10 lbs. ai)	Begin applications at the first sign of disease or when environmental conditions are favorable for disease development. Apply using a minimum of 20 gallons of water per acre by ground or a minimum of 10 gallons of water by air.	Do not apply more than 4.0 oz (0.10 lbs. of ai) of AGRISTAR® SONOMA® 40WSP product per acre per application. Applications may be up to and including the day of harvest. Observe a 30-day plant-back interval between the last application and planting new crops at the treatment site. Do not exceed 21 days between application intervals. Do not make more than five applications per year. Do not apply more than 20 ounces of AGRISTAR® SONOMA® 40WSP product (0.50 lbs. ai) per acre per year.

USE DIRECTIONS FOR NON-FOOD USE

DISEASES	AGRI STAR® SONOMA® 40WSP (OZ/ACRE)	APPLICATION DIRECTIONS	RESTRICTIONS
HYBRID POPLAR (For use in nurseries or forested areas used for wood pulp production)			
Rust (<i>Melampsora</i> spp.)	4 - 6 oz. (0.10 - 0.15 lbs. ai)	Begin applications at the first sign of disease and repeat applications at 10- to 14-day intervals.	Do not apply more than 6 oz. of AGRI STAR® SONOMA® 40WSP product (0.15 lbs. ai) per acre per application. Do not make more than five applications per year. Do not apply more than 30 ounces of AGRI STAR® SONOMA® 40WSP product (0.75 lbs. ai) per acre per year.
DOUGLAS FIR (Nursery Use Only)			
Needle Rust (<i>Melampsora</i> spp.)	5 - 10 oz. (0.125 - 0.25 lbs. ai)	Begin applications in early spring. Continue applications at 2 to 3 weeks intervals until the threat of infection is past. A spray adjuvant such as Latron CS-7 or Latron B-1956 should be added to obtain good spray coverage and disease control.	Do not apply more than 6 oz. of AGRI STAR® SONOMA® 40WSP product (0.15 lbs. ai) per acre per application. Do not make more than five applications per year.
LOBLOLLY PINE (Nursery Use Only)			
Fusiform Rust (<i>Cronartium quercuum</i>)	5 - 10 oz. (0.125 - 0.25 lbs. ai)	Begin applications in early spring. Continue applications at 2 to 3 week intervals until the threat of infection is past. A spray adjuvant such as Latron CS-7 or Latron B-1956 should be added to obtain good spray coverage and disease control.	Do not apply more than 10 oz. of AGRI STAR® SONOMA® 40WSP product (0.25 lbs. ai) per acre per application. Do not make more than five applications per year. Do not apply more than 50 ounces of AGRI STAR® SONOMA® 40WSP product (1.25 lbs. ai) per acre per year.

Cotton Seed Treatment

Use AGRI STAR® SONOMA® 40WSP as a seed treatment to protect against sore shin (*Rhizoctonia solani*) and black root rot (*Thielaviopsis basicola*), diseases which impair good cotton seed germination and seedling development. AGRI STAR® SONOMA® 40WSP may be applied to dry seed with conventional slurry or mist seed treating equipment. This product can be pumped or poured directly into the cotton seed treater provided the specified amount of product is applied. For best results, the seed must be completely and uniformly covered with fungicide. AGRI STAR® SONOMA® 40WSP can be tank mixed with other properly labeled cotton seed protectants such as Apron TL for Pythium control. Do not tank mix with products whose labels contain a prohibition against tank mixing.

SEED CONTAINER LABEL REQUIREMENTS

Commercial Cotton Seed Treatment

Federal Seed Act requires that containers containing treated seeds shall be labeled with the following information:

“This seed is treated with AGRI STAR® SONOMA® 40WSP which contains the active ingredient myclobutanil. Do not use for food, feed, oil production or any other purpose except planting. Do not graze cotton plants grown from treated seed.”

Store away from food and feedstuffs.

Do not allow children, pets, or livestock to have access to treated seeds.

Wear long pants, long-sleeved shirt and protective gloves when handling treated seed.

Treated seeds exposed on the soil surface may be hazardous to wildlife.

Cover or collect treated seeds spilled during loading and planting (such as in row ends).

Dispose of all excess treated seed by burying seed away from bodies of water.

Do not contaminate bodies of water when disposing of planting equipment wash water.

Dispose of seed packaging or containers in accordance with local requirements.

Excess treated seed may be used for ethanol production if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by-products that are used in agronomic practice.

Restrictions:

In California, this product must be applied through a closed system such as WECO-RED (Returnable Evacuation Device).

Do not use for food, feed, oil production or any other purpose except planting.

Do not graze cotton plants grown from treated seed. Store away from feed and foodstuff.

Use 0.05 to 0.155 lbs. of AGRI STAR® SONOMA® 40WSP product (0.019 to 0.062 lbs. ai) per 100 lbs. of cotton seed to reduce seedling damage caused by sore shin (*Rhizoctonia solani*) and black root rot (*Thielaviopsis basicola*).

For cotton seed treatment, seed treated with this product must be visually identifiable from untreated seed by the use of an approved colorant or dye to prevent accidental use of treated seed as food for humans or feed for animals. Refer to 21 CFR, Part 2.25. Any colorant or dye added to treated seed must be cleared for use in accordance with 40 CFR, Part 153.155(c).

On Farm Cotton Seed Treatment

For mechanical treaters or hopper box treatments, dilute 0.8 to 2.5 oz of AGRI STAR® SONOMA® 40WSP product (0.32 to 1 oz ai) with 8 to 10 fl oz of water and apply per 100 lbs. of cotton seed. For on farm cotton seed treatment, loaders must wear a chemical-resistant apron in addition to the PPE listed on the label affixed to the container.

Chemigation: Do not apply this product through any type of irrigation system.

TERMS AND CONDITIONS OF USE

If terms of the following **Warranty Disclaimer**, **Inherent Risks of Use**, and **Limitation of Remedies** are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under **Warranty Disclaimer**, **Inherent Risks of Use** and **Limitations of Remedies**.

Warranty Disclaimer

Albaugh, LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. **Albaugh, LLC. MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.**

INHERENT RISKS OF USE

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Albaugh, LLC or the seller. To the extent consistent with applicable law, all such risks shall be assumed by the buyer.

LIMITATION OF REMEDIES

To the extent consistent with applicable law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Albaugh, LLC's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used

To the extent consistent with applicable law, Albaugh, LLC shall not be liable for losses or damages resulting from handling or use of this product unless Albaugh, LLC is promptly notified of such loss or damage in writing. In no case shall Albaugh, LLC be liable for consequential or incidental damages or losses.

The terms of the **Warranty Disclaimer** and **Inherent Risks of Use** above and this **Limitation of Remedies** cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Albaugh or the seller is authorized to vary or exceed the terms of the **Warranty Disclaimer** or this **Limitation of Remedies** in any manner.

