

Supplemental Labeling



EPA Reg. No.: 279-3337

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL ARE TO BE FOLLOWED.

This Supplemental labeling must be in the possession of the user at the time of pesticide application. Read the label affixed to the container for SPARTAN CHARGE Herbicide before applying. Carefully follow all precautionary statements and application use directions.

CROP ROTATIONAL INTERVALS

Shown below are the minimum intervals in months from the time of SPARTAN CHARGE application until SPARTAN CHARGE treated soil may be replanted with the crops listed. When SPARTAN CHARGE is tank mixed with other herbicide(s), refer to all those labels for re-cropping instructions, following the intervals that are the most restrictive. For crops not listed, the interval is 12 months in addition to a successful field bioassay.

The field bioassay is a test strip of the intended crop planted across the previously treated field and grown to maturity. The test strip should include low spots, knolls, and variable pH and soil types. If crop responses are not observed, the crop may be planted the following year.

CROP ROTATION INTERVALS*

CROP	INTERVAL (Months)
Alfalfa	12
Barley	4
Berries (Crop subgroup 13-07)	Anytime
Buckwheat	18
Cabbage (transplant only)	Anytime
Canola, Crambe	24
Corn, field	4
Corn, pop	12
Corn, seed	12
Corn, sweet	12
Cotton	18 or 12***
Dry Shell Peas	Anytime
Dry Shell Beans	4
Flax	Anytime
Garlic	36
Grapes	Anytime
Horseradish	Anytime
Lima Beans (Tennessee only)	Anytime
Mint	Anytime
Oats	12
Peanuts	4

Potatoes	4
Rice	10
Rye	4
Sorghum	10**
Soybeans	Anytime
Sugar Beets	24
Sugarcane	Anytime
Sunflowers	Anytime
Sweet Potatoes	12
Tobacco	Anytime
Triticale	4
Turf	Anytime
Wheat	4

* For all other crops not listed, the rotation interval is a minimum of 12 months with a successful bioassay.

** 18 month rotation for rates above 10.2 fluid ounces per acre. Crops that have rotational intervals greater than 12 months after a SPARTAN CHARGE application are the result of crop injury concerns. The crops should only be planted after a successful bioassay.

*** Cotton may be planted after 12 months where SPARTAN CHARGE was applied at rates 8 oz/acre or less and meets the following conditions:

- Medium and fine soils
- pH <7.2
- Rainfall or irrigation must exceed 15" after application of SPARTAN CHARGE to rotate to cotton



FMC Corporation
Agriculture Products Group
1735 Market Street
Philadelphia, PA 19103

FMC and Spartan -trademarks of FMC Corporation
© 2013 FMC Corporation All Rights Reserved
061813