



Bayer CropScience LP  
 P.O. Box 12014  
 2 T.W. Alexander Drive  
 Research Triangle Park, North Carolina 27709  
 1-866-99BAYER (1-866-992-2937)

## Stratego® Fungicide

For Control of yellow leaf blight (*Phyllosticta maydis*)  
 in Field and Sweet Corn.

EPA Reg. No. 264-779

# PRODUCT BULLETIN

## FOR DISTRIBUTION AND USE ONLY WITHIN THE STATES OF MINNESOTA and WISCONSIN

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. It is a violation of federal law to use this product in a manner inconsistent with its label. This product bulletin must be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

### DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its label.

### FIELD (Field Corn, Field Corn Grown for Seed and Popcorn)

Disease Controlled	Rate fluid ounces/Acre
<b>Yellow Leaf Blight</b> <i>(Phyllosticta maydis)</i>	10.0 – 12.0

**Note:** Apply Stratego Fungicide when disease first appears and continue on a 7- to 14-day interval if favorable conditions for disease development persist.

Use the higher rates and shorter intervals when disease pressure is severe.

Application of Stratego Fungicide is not recommended at times when corn is under severe environmental stress conditions.

Stratego Fungicide may be applied by ground, air or chemigation.

**Restrictions:** Do not apply more than 24 fl. oz. of Stratego Fungicide per acre per crop. Do not apply within 30 days of harvest for forage, grain and stover. Do not apply more than 2 sequential applications of Stratego Fungicide. Limit the number of Stratego Fungicide or other (Group 11) -containing fungicide application to no more than 2 per acre per crop.

SWEET CORN (Including Seed Production)

Disease Controlled	Rate fluid ounces./Acre
<b>Yellow Leaf Blight</b> <b>(<i>Phyllosticta maydis</i>)</b>	10.0
<p><b>Note:</b> Apply Stratego Fungicide when disease first appears and continue on a 7-14-day interval if favorable conditions for disease development persist. Alternate applications of Stratego another product with a different mode of action than Group 11 fungicides.</p> <p>Stratego Fungicide may be applied by ground, air or chemigation.</p>	
<p><b>Restrictions:</b> Do not apply more than 30 fl. oz. of Stratego Fungicide/acre/season. Do not apply to sweet corn within 14 days of harvest. Do not harvest sweet corn for forage within 14 days of application. In programs with Stratego Fungicide, with Group 11 tank mixes, or other pre-mixes containing a Group 11 fungicide, the number of Group 11 fungicides should be no more than ½ the total number of fungicide applications per season. Alternate every application of Stratego Fungicide with at least one application of a non Group 11 fungicide.</p>	

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information, Call 1-866-99BAYER (1-866-992-2937)

or

Visit our worldwide web site at [www.bayercropscienceus.com](http://www.bayercropscienceus.com)

*As with any crop-protection product, always read and follow label directions.*

*For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*

Stratego is a trademark of Bayer