



Bayer CropScience LP  
P.O. Box 12014  
2 T.W. Alexander Drive  
Research Triangle Park, North Carolina 27709  
1-866-99BAYER (1-866-992-2937)

**For the use of a tank mix of:**

**Stratego YLD Fungicide (EPA Reg. No. 264-1093) + Proline 480 SC Fungicide (EPA Reg. No. 264-825)**

**On Chickpea, Dry Peas, and Lentils**

## ***FIFRA 2(ee) Recommendation***

### **FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE(S) OF MONTANA AND NORTH DAKOTA**

**FIFRA Section 2(ee) Recommendation:** This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

#### **DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its label.

CHICKPEA		
Disease Controlled	Product Rates	Product Instructions
Anthracnose <i>(Colletotrichum truncatum)</i>	Stratego YLD Fungicide – 4.8 fl oz/A	Begin fungicide applications preventatively and continue as needed on a 10- to 14-day interval. Use the shorter intervals when conditions are favorable for severe disease pressure.
Ascochyta blight <i>(Ascochyta rabiei)</i>	tankmixed with Proline 480 SC Fungicide – 3.2 fl oz/A	
Grey Mold <i>(Botrytis cinerea)</i>		
White mold <i>(Sclerotinia sclerotiorum)</i>		
<b>Application Restrictions:</b> Pre-Harvest Interval (PHI): 30 day(s) If growing for animal feed, do not apply within 7 days of cutting or swathing of the crop for forage. Do not make more than 2 applications of this product combination per year. Do not exceed 0.28 lbs prothioconazole per acre per year, or 0.24 lbs trifloxystrobin per acre per year.		

DRY PEAS		
Disease Controlled	Product Rates	Product Instructions
Ascochyta leaf and pod spot ( <i>Ascochyta pisi</i> )	Stratego YLD Fungicide – 4.8 fl oz/A	Begin fungicide applications preventatively and continue as needed on a 10- to 14-day interval.  Use the shorter intervals when conditions are favorable for severe disease pressure.
Mycosphaerella blight ( <i>Mycosphaerella pinodes</i> )	tankmixed with Proline 480 SC Fungicide – 3.2 fl oz/A	
<p>Application Restrictions:</p> <p>Pre-Harvest Interval (PHI): 30 day(s)</p> <p>If growing for animal feed, do not apply within 7 days of cutting or swathing of the crop for forage.</p> <p>Do not make more than 2 applications of this product combination per year.</p> <p>Do not exceed 0.28 lbs prothioconazole per acre per year, or 0.24 lbs trifloxystrobin per acre per year.</p>		

LENTILS		
Disease Controlled	Product Rates	Product Instructions
Anthracnose ( <i>Colletotrichum truncatum</i> )	Stratego YLD Fungicide – 4.8 fl oz/A	Begin fungicide applications preventatively and continue as needed on a 10- to 14-day interval.  Use the shorter intervals when conditions are favorable for severe disease pressure.
Ascochyta blight ( <i>Ascochyta rabiei</i> )	tankmixed with Proline 480 SC Fungicide – 3.2 fl oz/A	
Grey Mold ( <i>Botrytis cinerea</i> )		
White mold ( <i>Sclerotinia sclerotiorum</i> )		
<p>Application Restrictions:</p> <p>Pre-Harvest Interval (PHI): 30 day(s)</p> <p>If growing for animal feed, do not apply within 7 days of cutting or swathing of the crop for forage.</p> <p>Do not make more than 2 applications of this product combination per year.</p> <p>Do not exceed 0.28 lbs prothioconazole per acre per year, or 0.24 lbs trifloxystrobin per acre per year.</p>		

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.  
For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937).

As with any crop-protection product, always read and follow label directions.  
For additional information call toll-free 1-866-99BAYER (1-866-992-2937).