



**Bayer CropScience**

**Bayer CropScience LP  
P.O. Box 12014  
2 T.W. Alexander Drive  
Research Triangle Park, North Carolina 27709  
1-866-99BAYER (1-866-992-2937)**

**STRATEGO® YLD Fungicide**

**EPA Reg. No.: 264-1093**

**For use on Barley and Wheat**

***Supplemental Label***

**STRATEGO® YLD Fungicide**

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read this label and the product package label before using this product. This Supplemental Label must be in possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the registered product package label for STRATEGO® YLD Fungicide

**USE DIRECTIONS**

<b>BARLEY</b>		
	<b>Disease Control</b>	<b>Rate fl. oz./acre</b>
Glume Blotch ( <i>Stagonospora nodorum</i> )	Rusts ( <i>Puccinia spp.</i> )	2.3
Leaf Blotch ( <i>Stagonospora avenae</i> )	Scald ( <i>Rhynchosporium secalis</i> )	
Net Blotch ( <i>Pyrenophora teres</i> )	Spot Blotch ( <i>Cochliobolus sativus</i> )	
Powdery Mildew ( <i>Blumeria graminis</i> )		
<p><b>Note:</b> Begin applications preventively when conditions are favorable for disease development. A second application (minimum interval of 14 days) may be made if needed. Stratego YLD Fungicide may be applied by ground, air or chemigation.</p>		
<p><b>Restrictions:</b> Do not apply more than 2 applications of Stratego YLD Fungicide per season. Do not apply more than 4.6 fl. oz. per acre per year. Do not apply after Feekes growth stage 8 (the ligule of the flag leaf emerges). Do not apply within 40 days of harvest.</p>		
<p><b>Grazing Restrictions:</b> (a). If 1 application or a total of 2.3 fl. oz of Stratego YLD Fungicide per season are applied, do not allow livestock to graze within the treated area within 30 days after application, and do not harvest the treated crop for forage within 30 days after application or for hay within 45 days after application. (b). If 2 applications or a total of 4.6 fl. oz. of Stratego YLD Fungicide per season are applied, do not allow livestock to graze within the treated area and do not harvest the treated crop for forage or hay.</p>		

WHEAT		
Disease Control		Rate fl. oz/acre
Stagonospora Blotch ( <i>Stagonospora nodorum</i> )	Rusts ( <i>Puccinia spp.</i> )	4.0
Septoria Blotch ( <i>Septoria tritici</i> )	Tan Spot ( <i>Pyrenophora tritici-repentis</i> )	
Powdery Mildew ( <i>Blumeria graminis f. sp. tritici</i> )		

**Note:** Begin applications preventatively when conditions are favorable for disease development. A second application (minimum interval of 14 days) may be made if needed.  
Stratego YLD Fungicide may be applied by ground, air or chemigation.

**Restrictions:** Do not apply more than 2 applications of Stratego YLD Fungicide per season. Do not apply after Feekes growth stage 10.5 (full head emergence). Do not apply within 35 days of harvest. Do not apply more than 2 applications of Stratego YLD Fungicide or other Group 11 -containing fungicide per acre per year without alternation with at least 2 applications of fungicide from a different (not Group 11) mode of action.

**Grazing Restrictions:** If up to a total of 4 fl. oz of Stratego YLD Fungicide per season are applied, do not allow livestock to graze within the treated area within 30 days after application, and do not harvest the treated crop for forage within 30 days after application or for hay within 45 days after application. If greater than 4 fl. oz. of Stratego YLD Fungicide are applied per season, do not allow livestock to gaze within the treated area and do not harvest the treated crop for forage or hay.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577  
For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

As with any crop-protection product, always read and follow label directions.  
For additional information call toll-free 1-866-99BAYER (1-866-992-2937).