



EPA Reg. No. 264-779-2935

**For Distribution and Use Only Within the States of Minnesota and Wisconsin**

**FOR CONTROL OF YELLOW LEAF BLIGHT (*Phyllosticta maydis*) IN FIELD AND SWEET CORN**

**ATTENTION**

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- Before using TARIAN™ in accordance with this recommendation, read and follow all applicable directions for use, restrictions, and precautions on the EPA registered label, as found on the package.
- This recommendation should be in the possession of the user at the time of application.
- This recommendation for use of this product is permitted under Section 2(ee) of FIFRA and has not been submitted to or been approved by EPA.

**DIRECTIONS FOR USE**

**FIELD (Field Corn, Field Corn Grown for Seed, and Popcorn)**

For Yellow Leaf Blight (*Phyllosticta maydis*) apply 10 to 12 fl oz per acre. Apply TARIAN when disease first appears and continue on a 7- to -14-day interval if favorable conditions for disease development persist. Use the higher rates and shorter intervals when disease pressure is severe. Application of this product is not recommended at times when corn is under severe environmental stress conditions. TARIAN may be applied by ground, air or chemigation.

**Restrictions:** Do not apply more than 24 fl oz of TARIAN per acre per crop. Do not apply within 30 days of harvest for forage, grain and stover. Do not apply more than 2 sequential applications of TARIAN. Limit the number of TARIAN or other Group 11- containing fungicide applications to no more than 2 per acre per crop.

**SWEET CORN (Including Seed Production)**

For Yellow Leaf Blight (*Phyllosticta maydis*) apply 10 fl oz per acre. Apply TARIAN when disease first appears and continue on a 7- to -14-day interval if favorable conditions for disease development persist. Alternate applications of TARIAN with another product with a different mode of action than Group 11 fungicides. TARIAN may be applied by ground, air or chemigation.

**Restrictions:** Do not apply more than 30 fl oz of TARIAN per acre per season. Do not apply to sweet corn within 14 days of harvest. Do not harvest sweet corn for forage within 14 days of application. In programs with TARIAN, with Group 11 tank mixes, or other pre-mixes containing a Group 11 fungicide, the number of Group 11 fungicides should be no more than ½ the total number of fungicide applications per season. Alternate every application of TARIAN with at least one application of a non-Group 11 fungicide.

All trademarks, service marks, trade names, trade dress, product names and logos appearing on this label are the property of their respective owners.

2022-0608

