

RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators.



ENTRA

Section 24(c) Special Local Need Label

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF GEORGIA

DICAMBA	GROUP	4	HERBICIDE
S-METOLACHLOR	GROUP	15	HERBICIDE

TAVIUM® PLUS VAPORGRIP® TECHNOLOGY

**EPA Reg. No. 100-1623
EPA SLN No. GA-190006**

**ADDITIONAL APPLICATION REQUIREMENTS FOR USE ON DICAMBA-TOLERANT
COTTON OR DICAMBA-TOLERANT SOYBEANS**

This special local needs label expires on December 20, 2020 and must not be used or distributed after this date.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label must be in the possession of the user at the time of application.
- Refer to the Tavium® Plus VaporGrip® Technology container label, EPA Reg. No. 100-1623, for complete Directions for Use and all applicable restrictions.

RESTRICTIONS/LIMITATIONS

In order to apply the Tavium Plus VaporGrip Technology to dicamba-tolerant cotton and dicamba-tolerant soybeans in Georgia the following restrictions apply:

Using Pesticides Wisely Training: The certified applicator applying this product on cotton or soybeans, in-crop, must attend Using Pesticides Wisely training at least once every two years that is approved by the Georgia Department of Agriculture. For more information on this training, contact the Georgia Department of Agriculture at 404-656-4958 or your local County Cooperative Extension Service Office.

ADDITIONAL APPLICATION METHODS

Hooded Row-Middle Spray Applications: Applications may be made with any standard spray tip for hoods as long as the system generates droplets that are coarse or larger in size (>341 microns vmd). Contact UGA Extension or your local Syngenta representative for questions.

Application Requirements:

1. Hoods must remain in contact with the soil while the herbicide application is being made.
2. A maximum of 6 mph is required to ensure hoods remain in contact with the soil.
3. Cotton or soybeans must be at least 15 inches tall at time of application.
4. Do not make more than one in-crop application (postemergence, hooded row-middle, or layby) to cotton or soybean.

Application Timing:

Dicamba-Tolerant Cotton: Do not apply later than 80 days before harvest.

Dicamba-Tolerant Soybean: Do not apply later than 90 days before harvest.

Directed Layby Applications: Applications may be made with any standard spray tip as long as the system generates droplets that are coarse or larger in size (>341 microns vmd). Contact UGA Extension or your local Syngenta representative for questions.

Application Requirements:

1. Release point for the herbicide must not be more than 10 inches from the soil and the cotton or soybean must be at least 20 inches in height. Spray tip must be angled downward to the soil making sure no spray droplets remain in the air.
2. A maximum of 6 mph is required to ensure herbicide does not lift above the release point.
3. Do not make more than one in-crop application (postemergence, hooded row-middle, or layby) to cotton or soybean.

Application Timing:

Dicamba-Tolerant Cotton: Do not apply later than 80 days before harvest.

Dicamba-Tolerant Soybean: Do not apply later than 90 days before harvest.

©2019 Syngenta

Tavium® and the Syngenta logo are trademarks of a Syngenta Group Company.

VaporGrip® is a trademark of Monsanto Technology LLC.

24(c) Registrant:
Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, NC 27419-8300

Label Code: GA1623053BA1119