



TECH-SPRAY[®]



N-SEASON 27-0-0

with 66% Slow Release Nitrogen

GUARANTEED ANALYSIS

Total Nitrogen (N).....27.0%
8.5% Urea Nitrogen
18.5% Water soluble nitrogen

Derived from: Urea, Polymethylene urea, Methylene Diurea and Monomethyl Urea.

10.50 lbs/gal

Each gallon contains 2.86 pounds of Nitrogen

TECH-SPRAY N-SEASON supplies the guaranteed nutrients in a water soluble form for use as a nutritional spray. It is designed to be combined with TECH-FLO products but may also be applied alone.

NOTICE: Do not store this product at temperatures below 40°F as precipitates may form that are difficult to dissolve. Do not expose to temperatures below 32°F.

MANUFACTURED FOR
NUTRIENT TECHNOLOGIES, INC.
1092 E. KAMM AVE., DINUBA CA 93618
(559)595-8090

TECH-SPRAY® N-SEASON

27-0-0

TS

GENERAL RECOMMENDATIONS AS A NUTRITIONAL SPRAY:

DILUTION: Apply suggested rates of Tech-Spray® in the following amounts of water per acre.
Minimum dilution rate: 1 qt per 5 gals water per suggested use rates

Field & Row Crops:

By air	3-20 gals.
By ground	10 gals. minimum.
Tree Crops:	
By air	10 gals. minimum.
By ground concentrate	25-100 gals.
By ground dilute	100-500 gals.

COMPATIBILITY: This product is designed to be compatible with most commonly used insecticides, fungicides, acaricides, and NPK fertilizers. However, it is not feasible to test all possible combinations of pesticides with which it may be mixed. A jar test is recommended. Consult your Nutrient Technologies, Inc. Representative for detailed compatibility information.

SUGGESTED USES (RATES PER ACRE):

- ALFALFA:** Apply 1-2 gals after each cutting when sufficient foliage is present.
ALMONDS: Apply 1-1½ gals at pink bud to early bloom, apply 2-2½ gals in 30 days.
Post Harvest: Apply 3-5 gals in 1 or 2 applications
APPLES: Apply 1-1½ gals at bud swell to king bloom, repeat 30 days later and 30-40 days prior to harvest for sizing.
Post harvest: Apply 3-5 gals in 1 or 2 applications.
BEANS, PEAS and RELATED LEGUMES: Apply 1-3 gals at early bloom, repeat after pods are formed.
CABBAGE, CAULIFLOWER, BROCCOLI, & OTHER COLE CROPS: 2-3 gals at early head formation and repeat 7-10 days later.
CARROTS, ONIONS, RADISHES, BEETS, & OTHER ROOT VEGETABLES:
Apply 1-2 gals. Make 3-4 applications.
CITRUS, AVOCADO: Apply 1-5 gals at pre-bloom or bloom to increase fruit set, repeat in 30 days and as extra nitrogen is needed.
CORN (FIELD): Apply up to 5 gals with seed at planting. Foliar apply up to 2 gals as extra nitrogen is needed.
COTTON: Apply 1-2 qts at 4th leaf, then 2-5 gals at early bloom, repeat as needed.
CUCUMBERS, SQUASH, & OTHER CUCURBITS: Apply 1-3 gals at early bloom, repeat approximately 4 weeks after bloom application.
GRAPES: Apply 1-2 gals when nitrogen needs are evident or an increase in canopy is required to reduce sunburn.
LETTUCE, CELERY, SPINACH, & OTHER LEAFY GREENS: 1-2 gals 10 days after transplanting. 2-3 gal. after thinning, repeat in 2-3 weeks and 2-3 weeks prior to harvest, using 7-8 gallons total over crop.
MILK & OTHER SORGHUMS: Apply up to 3 gals with seed at planting.
Foliar apply up to 2 gals any time extra nitrogen is needed.

- MINT:** (PEPPERMINT, SPEARMINT): Apply 2-5 gals as needed
OLIVES: Apply 1-3 gals pre-bloom, repeat in 30 days. Apply 4-5 gals 4-6 weeks prior to harvest
PEACHES, PLUMS, CHERRIES, & OTHER STONE FRUIT: Apply 1-1½ gals at bud swell to early bloom to increase fruit set, repeat 30 days to maintain. Apply 2-4 gals 4-6 weeks prior to harvest for improved fruit sizing, firmness and reduction in "green shoulders".
Post Harvest: apply 5-10 gals in 1 or 2 applications.
PEARS (BARTLETT): Apply 1-1½ gals at pre-bloom and repeat every 30 days until 4 weeks prior to harvest. As some varieties (i.e., D'Anjou) are more sensitive under certain conditions, use caution.
Post Harvest: apply 5-10 gals/acre in 1 or 2 applications.
PEANUTS: Apply 3-5 gals at early bloom, repeat as necessary until pods filled.
PEPPERS: Apply 2-3 gals at early bloom and repeat 14 days later.
PISTACHIOS, PECANS & OTHER DECIDUOUS TREE NUTS: Apply 1-1½ gals, repeat every 2-3 weeks as needed.
Post Harvest: Apply 3-5 gals in 1 or 2 applications
POTATOES: Apply 2 gals prior to tuberization and at early bloom. 2-3 weeks following bloom spray may be needed through fertigation system.
RASPBERRIES, BLUEBERRIES, & OTHER SMALL BERRIES: Apply 1-2 gals at early fruit set, repeat at early fruit color.
SEED CROPS: Apply 1½-2 gals at early bloom, repeat in 3-4 weeks and 30-60 days prior to harvest to fill out seed.
SOYBEANS: Apply 2-3 gals at early pod. Apply 1-2 gals as needed.
STRAWBERRIES: Apply 1-3 gals at early bloom, repeat in 14-21 days.
SWEET CORN: Apply 2-3 gals at a 12-14 inch height, at early tassel stage, and again at early silk stage.
TOMATOES: Apply 2-3 gals at early bloom, repeat at fruit set and again 15-30 days later. Apply 3-4 gals 3-4 weeks prior to harvest to strengthen canopy and to reduce sunburn.
WHEAT: Apply 2-5 gals at 6th or 7th leaf as needed. Can be applied with herbicides at 1-2 gals/acre.

RECOMMENDED DRIP FERTIGATION RATES:

- STRAWBERRIES:** 3-4 gals/acre twice a month for five to six months in CA.
TOMATOES/PEPPERS: 3-5 gals/acre twice a month for three to four months.
GRAPES: 3-5 gals/acre twice a month for three to four months
LETTUCE: 3-5 gals/acre at first irrigation and repeat as needed.
TREES & VINES: 5 gals/acre at least every 30 days in sandy soils. Apply with other drip materials at 25 and 50% ratios to improve fertilizer efficiency.

NOTE: For crops not listed on this label, consult your NUTRI-TECH representative.

NOTICE: Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>.

NOTICE: Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used according to directions under normal conditions on the crops specified. Because of widely varying use conditions, it is impossible to eliminate all risks even when the label directions are followed. Therefore, the manufacturer makes no other warranties, expressed or implied. The buyer assumes all risks associated with the use of this product. In the event of damage resulting from a breach of warranty, the buyer agrees to accept a refund of the purchase price as full discharge of the manufacturer's liability.

N.1



WARNING

Causes eye and skin irritation

Wear protective clothing, gloves, eye and face protection. Do not eat, drink or smoke when using this product. Wash hands thoroughly after handling. Remove and wash contaminated clothing before reuse. Avoid breathing vapor, mist or dust from this product. Not for human or animal consumption.

If in eyes: Rinse thoroughly with water for several minutes. Continue rinsing and get medical attention if eye irritation persists.
If on skin: Wash with plenty of soap and water.
If swallowed: Rinse mouth and get medical attention if you feel unwell.

Store in a cool, dry, well-ventilated area away from heat, combustible and incompatible materials. Dispose of unused contents, container, and other contaminated wastes in accordance with local, state, and federal law.

MADE IN USA
NUTRIENT TECHNOLOGIES, INC.
1092 E. KAMM AVE., DINUBA CA 93618
(559)595-8090