

RESTRICTED USE PESTICIDE

Due to high acute inhalation toxicity and carcinogenicity. For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

Product Bulletin

Salt Lake Holding LLC

Salt Lake Holding LLC

2211 H. H. Dow Way

Midland, MI 48674 USA

TELONE™ II

Soil Fumigant

EPA Reg. No. 95290-1

Special 2(ee) Recommendation†

For Distribution and Use Only Within the States of Alabama, Florida, Georgia, and South Carolina

Nematode Suppression in Peanuts

ATTENTION

†This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA.

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- Read the label affixed to the container for TELONE™ II Soil Fumigant before applying. Carefully follow all precautionary statements and applicable use directions.
- Except as specified in this product bulletin, use of this product is subject to all use precautions and limitations imposed by the label affixed to the container of TELONE™ II.

Directions for Use

For suppression of light to moderate nematode infestations in peanuts, apply TELONE™ II Soil Fumigant as a broadcast treatment at the rate of 6 to 9 gallons per acre. Apply when soil moisture is ideal for tillage using a moldboard plow or other suitable tillage implement capable of placing TELONE™ II at least 12 inches below the surface of the final seedbed. Seal soil immediately after application according to soil sealing directions on the label affixed to the container of TELONE™ II. Do not disturb the soil for at least one week prior to planting to ensure adequate exposure of target pests to TELONE™ II.

TELONE™ II may also be applied as an in-row treatment at 4.5 to 6 gallons per acre.

™ Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

01-2ee-A-04

Issued: 12/12/21

Replaces 01-2ee-A-03.