



SUPPLEMENTAL LABEL

TOPSIN® 4.5 FL Fungicide

For use on Soybeans to Control White Mold

EPA Reg. No. 8033-122-70506

This supplemental label expires on April 30, 2023
and must not be used or distributed after this date.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
Follow all applicable directions, restrictions and precautions on the federally-registered label.

This label must be in the possession of the user at the time of pesticide application. Read the label affixed to the container for TOPSIN 4.5 FL Fungicide before applying. Use of TOPSIN 4.5 FL Fungicide according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container of TOPSIN 4.5 FL Fungicide.

Crop	Disease	Fl. oz./A (lb a.i./A)	Application Instructions
Soybeans	White mold (Sclerotinia)	20 (0.7) for split application OR 40 (1.4) for single application	For split applications, apply initial application of 20 fl. oz of product/A (0.7 lb a.i./A) at early bloom (R-1 stage). Follow with a second application 7 to 14 days later. OR For a single application, apply 40 fl. oz. of product/A (1.4 lb a.i./A) at early bloom (R-1 stage). Thorough coverage of the flowers, stems, and branches is essential for disease control. Use a minimum of 5 gallons water/A by air.

Restrictions:

- DO NOT** apply more than 40 fl. oz./A (1.4 lb. a.i./A) per for a single application for white mold control otherwise
- DO NOT** apply more than 20 fl. oz./A (0.7 lb. a.i./A) for a single application.
- DO NOT** apply more than 40 fl. oz. of product (1.4 lbs a.i.)/A/year.
- The minimum retreatment interval is seven (7) days.
- DO NOT** apply more than two (2) applications per year.
- DO NOT** enter or allow worker entry into treated areas during the restricted entry interval (REI) of 1 day.
- Pre-harvest interval: 21 days
- DO NOT** graze or feed treated vines or hay to livestock.

MAY 2020