



SUPPLEMENTAL LABEL

TOPSIN® M WSB Fungicide

For use on Soybeans to Control White Mold

EPA Reg. No. 8033-125-70506

This supplemental label expires on April 30, 2023
and must not be used or distributed after this date.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
Follow all applicable directions, restrictions and precautions on the federally-registered label.

This label must be in the possession of the user at the time of pesticide application. Read the label affixed to the container for TOPSIN M WSB Fungicide before applying. Use of TOPSIN M WSB Fungicide according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container of TOPSIN M WSB Fungicide.

Crop	Disease	Lbs of product/A (lb a.i./A)	Application Instructions
Soybeans	White mold (Sclerotinia)	1.0 (0.7) for split application OR 2.0 (1.4) for single application	For split applications, apply initial application of 1.0 lb of product (0.7 lb a.i.)/A at early bloom (R-1 stage). Follow with a second application 7 to 14 days later. OR For a single application, apply 2.0 lbs of product (1.4 lb a.i.)/A at early bloom (R-1 stage). Thorough coverage of the flowers, stems, and branches is essential for disease control. Use a minimum of 5 gallons water/A by air.

Restrictions:

- DO NOT** apply more than 2.0 lbs of product/A (1.4 lb. a.i./A) for a single application for white mold control otherwise **DO NOT** apply more than 1.0 lb of product/A (0.7 lb a.i./A) for a single application.
- DO NOT** apply more than 2.0 lbs of product (1.4 lbs a.i.)/A/year.
- The minimum retreatment interval is seven (7) days.
- DO NOT** apply more than two (2) applications per year.
- DO NOT** enter or allow worker entry into treated areas during the restricted entry interval (REI) of 1 day.
- Pre-harvest interval: 21 days
- DO NOT** graze or feed treated vines or hay to livestock.

MAY 2020