

PRODUCT BULLETIN
2(ee) Recommendation

**THIS PRODUCT BULLETIN CONTAINS NEW OR ADDITIONAL
DIRECTIONS FOR USE
WHICH ARE RECOMMENDED BY INNVICTIS CROP CARE LLC
UNDER FIFRA SECTION 2(ee)**



EPA Reg. No. 89168-52-89391

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or accepted by the U.S. Environmental Protection Agency. It is a violation of federal law to use this product in a manner inconsistent with its label. This product bulletin must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

DIRECTIONS FOR USE - It is a violation of Federal law to use this product inconsistent with its labeling.

Field Corn, Popcorn, Corn Grown For Seed Production			
Disease	Dosage Rate		Use Directions
	Fl. Oz./A	GPA	
Tar Spot <i>(Phyllachora maydis)</i>	10.0 to 17.0	Ground minimum: 10 Aerial minimum: 2	Apply prior to disease onset when conditions favor disease development. A second application may be made no fewer than 7 days later but not to exceed 17.0 fl oz per acre per year rate. Use <i>TREVO TRZ</i> Fungicide as part of an integrated pest management program (IPM). Apply as a foliar spray or via chemigation in sufficient water to obtain thorough coverage of plants.

RESTRICTIONS AND LIMITATIONS

1. **Do not** make more than two (2) applications per year.
2. **Do not** make more than 2 sequential applications before alternating to another fungicide with a different mode of action.
3. **Do not** apply more than 17 fl oz of TREVO TRZ per acre per year.
4. **Do not** apply more than 0.09 lb ai of a tetraconazole-containing product per acre per year.
5. **Do not** apply more than 2.0 lb ai of an azoxystrobin-containing product per acre per year.
6. **Do not** apply TREVO TRZ after corn growth stage R3 (brown silk/milk).
7. **Do not** use adjuvants in sprays made between V8 (8 leaf collar) and VT (lowest branch of the tassel visible but silks have not emerged) growth stage. A compatibility agent, another fungicide, or an insecticide may be included if needed and labeled for use in corn. Refer to adjuvant product label for specific use directions and restrictions. Always follow the more restrictive label.
8. For chemigation, apply in 0.1 – 0.25 inches/A of water.
9. Do not apply within 7 days of harvest (7 day PHI).
10. Do not harvest silage within 21 days of an application.

Use Rate: 10.0 to 17.0 oz per acre

Always refer to the complete product label for full use directions and precautionary language.

Distributed By:

INNVICTIS CROP CARE, LLC
1880 Fall River Drive, Suite 100
Loveland, CO 80538

080119R1