



Bayer Environmental Science, a division of Bayer  
 CropScience  
 P.O. Box 12014  
 2 T.W. Alexander Drive  
 Research Triangle Park, North Carolina 27709  
 1-800-331-2867

## Tartan® Stressgard®

EPA Reg. No. 432-1446

For control of Take-All Root Rot,  
 Bermudagrass decline, and Warm Season  
 Turfgrass Decline (*Gaeumannomyces  
 graminis var. graminis*)

Expires: 6/14/2020

# PRODUCT BULLETIN

**FOR DISTRIBUTION AND USE ONLY IN THE STATES OF FLORIDA, TEXAS,  
 HAWAII, GEORGIA, SOUTH CAROLINA, NORTH CAROLINA, TENNESSEE,  
 ARKANSAS, OKLAHOMA, KANSAS, MISSOURI, MISSISSIPPI, ALABAMA,  
 LOUISIANA, NEW MEXICO, AND ARIZONA**

**FIFRA Section 2(ee) Recommendation:** This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. It is a violation of federal law to use this product in a manner inconsistent with its label.

**This product bulletin must be in the possession of the user at the time of application of the product.** All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

### DIRECTIONS FOR USE

For use on turfgrass on golf courses, sod farms and sports fields only.

Disease	Tartan® Stressgard®	fl oz per 1,000 sq ft	Application Interval/Timing
Take-All Root Rot,	Apply 1 to 2 preventive applications in the spring and fall, prior to conditions favorable for disease development.	2 fl oz	28 days
Bermudagrass Decline,		2 fl oz	28 days
Warm Season Turfgrass Decline ( <i>Gaeumannomyces graminis var. graminis</i> )	Apply fungicide in adequate water volume and water-in applications to root zone.	2 fl oz	28 days

For MEDICAL and TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-800-331-2867

As with any plant-protection product, always read and follow label directions.