

Product Bulletin



Dow AgroSciences

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Restricted Use Pesticide

Due to high acute inhalation toxicity and carcinogenicity.
For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

Telone* II

(EPA Reg. No. 62719-32)

Special 2(ee) Recommendation†

Cyst-Forming Nematode Suppression In Sugar Beets

(For Distribution and Use Only Within the States of Idaho, Oregon and Washington)

ATTENTION

†This recommendation is made as permitted under FIFRA Section 2(ee) and has not been reviewed or approved by EPA.

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Telone II before applying. Carefully follow all precautionary statements and applicable use directions except as specified below.
- Use of Telone II according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Telone II.

Directions for Use

For Cyst-Forming Nematode Suppression In Sugar Beets: Apply Telone* II soil fumigant as a broadcast plow sole treatment at a rate of 12 to 17 gallons per acre for suppression of light to moderate infestations of cyst-forming nematodes. Treat when soil moisture is ideal for tillage. Telone II should be placed at least 12 inches below the surface of the final seedbed. Follow instructions for soil sealing and soil fumigation interval in the "Directions for Use" section of the product label for Telone II.

*Trademark of Dow AgroSciences LLC

D06-018-016
Issued: 08/10/00
Initial printing.

Basis for Section 2(ee) Recommendation:
Application at rates for suppression of light to moderate infestations of cyst-forming nematodes in sugar beets.