

# Product Bulletin



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

## RESTRICTED USE PESTICIDE

Due to high acute inhalation toxicity and carcinogenicity. For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

## Telone<sup>®</sup> II

EPA Reg. No. 62719-32

Special 2(ee) Recommendation<sup>†</sup>

## Nematode Suppression in Corn

### ATTENTION

<sup>†</sup>This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- Read the label affixed to the container for Telone<sup>®</sup> II soil fumigant before applying. Carefully follow all precautionary statements and applicable use directions except as specified below.
- Use of Telone II according to this Section 2(ee) recommendation is subject to all use precautions and limitations imposed by the label affixed to the container for Telone II.

### Directions for Use

Apply Telone II as a row treatment at 3 to 6 gallons per acre. Use suitable ground application equipment that will insure placement of the fumigant at least 14 inches below the final planting surface. Use one chisel per row to treat only the soil where the crop is to be planted. Soil should be sealed immediately after application. Soil sealing may be accomplished using bedding or strip till equipment that properly seals the soil. After application and sealing, leave the soil undisturbed for at least a week before planting.

#### Use Precaution:

- Do not plant if the fumigant is still present at the depth of injection.

<sup>®</sup>Trademark of Dow AgroSciences LLC

R018-015

Issued: 11/19/07

Initial printing.