

# RESTRICTED USE PESTICIDE

## ACUTE TOXICITY and GROUND WATER CONTAMINATION

For retail sale to and use only by Certified Applicators or persons under the direct supervision of a Certified Applicator, and only for those uses covered by the Certified Applicator's certification.



# TEMIK<sup>®</sup> brand 15G ALDICARB PESTICIDE

For Control of Certain Insects, Mites, and Nematodes

ACTIVE INGREDIENT: Aldicarb [2-methyl-2-(methylthio) propionaldehyde O-(methylcarbamoyl)oxime] ..... 15%

INERT INGREDIENTS: ..... 85%

EPA Reg. No. 264-330

EPA Est. No. 264-GA-01



**KEEP OUT OF REACH OF CHILDREN**  
**DANGER POISON**  
**PELIGRO**



Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

For **MEDICAL And TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577  
For **PRODUCT USE** Information Call 1-866-99BAYER (1-866-992-2937)

### FIRST AID

<b>IF SWALLOWED:</b>	<ul style="list-style-type: none"><li>• Immediately call a poison control center or doctor for treatment advice.</li><li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• Do not give anything by mouth to an unconscious person.</li></ul>
<b>IF IN EYES:</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>IF ON SKIN OR CLOTHING:</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>IF INHALED:</b>	<ul style="list-style-type: none"><li>• Move person to fresh air.</li><li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li><li>• Call a poison control center or doctor for further treatment advice.</li></ul>

**For MEDICAL Emergencies Call 24 Hours A Day 1-800-334-7577.**

**Have the product container or label with you when calling a poison control center or doctor or going for treatment.**

Aldicarb is a N-methyl carbamate.

### GENERAL

Contact a physician immediately in all cases of suspected poisoning. Illness may be produced rapidly following overexposure to TEMIK® Aldicarb. If breathing stops, establish an airway and start artificial respiration, and provide oxygen. Make certain to remove all sources of continuing contamination. Remove clothing and wash skin and hair immediately with large amounts of water. Transport the patient to a physician or hospital immediately and SHOW A COPY OF THIS LABEL TO THE PHYSICIAN. If poisoning is suspected in animals, contact a veterinarian.

#### **ANTIDOTE STATEMENT**

ATROPINE SULFATE IS HIGHLY EFFECTIVE AS AN ANTIDOTE. See NOTE TO PHYSICIAN.

**NOTE TO PHYSICIAN:** TEMIK® brand aldicarb is a n-methyl carbamate insecticide which is a cholinesterase inhibitor. Overexposure to this substance may cause toxic signs and symptoms due to stimulation of the cholinergic nervous system. These effects of overexposure are spontaneously and rapidly reversible.

Gastric lavage may be used if this product has been swallowed. TEMIK® brand aldicarb poisoning may occur rapidly after ingestion and prompt removal of stomach contents is indicated.

Specific treatment consists of the administration of parenteral atropine sulfate. Caution should be exercised to prevent overatropinization. Mild cases may be given 1 to 2 mg intramuscularly every 10 minutes until full atropinization has been achieved and repeated thereafter whenever symptoms reappear. Severe cases should be given 2 to 4 mg intravenously every 10 minutes until the patient is fully atropinized, then intramuscularly every 30 to 60 minutes as needed to maintain the effect for at least 12 hours. Dosages for children should be appropriately reduced. Complete recovery from overexposure is to be expected within 24 hours.

Narcotics and other sedatives should not be used. Further, drugs such as (pyridine-2-aldoxime methiodide) are NOT recommended unless organophosphate intoxication is also suggested.

To aid in confirmation of a diagnosis, urine samples must be obtained within 24 hours of exposure and immediately frozen. Analyses will be arranged by Bayer CropScience.

Consultation on therapy can be obtained at all hours by calling the Bayer CropScience emergency number: 1-800-334-7577.

## **PRECAUTIONARY STATEMENTS**

### **HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

#### **DANGER**

**FATAL IF SWALLOWED.** Causes cholinesterase inhibition. May be fatal or harmful by skin or eye contact or by breathing dust. Rapidly absorbed through skin or eyes. Do not get on skin or in eyes. Do not breathe dust. Keep away from domestic animals.

Always stand up-wind from hopper when loading.

#### **SIGNS AND SYMPTOMS OF OVEREXPOSURE:**

Salivation, Muscle tremor, Nausea, Watery eyes, Difficult breathing, Vomiting, Pinpoint eye pupils, Excessive sweating, Diarrhea, Blurred vision, Abdominal cramps, Weakness, Headache,

In severe cases, convulsions, unconsciousness, and respiratory failure may occur.

### **PERSONAL PROTECTIVE EQUIPMENT (PPE)**

Applicators and other handlers must wear coveralls over short-sleeved shirts and short pants, waterproof gloves, chemical-resistant footwear plus socks, chemical-resistant headgear for overhead exposure, protective eye wear, chemical-resistant apron when cleaning equipment, mixing or loading and a dust/mist filtering respirator (MSHA/NIOSH D/M approval number prefix TC-21C) or a NIOSH approved respirator with any N, R, P or HE filter.

Follow manufacturer's instructions for cleaning/maintaining PPE. Keep and wash PPE separately from other laundry. Wash contaminated clothing in strong washing soda solution and rinse thoroughly before reuse. Discard clothing or other absorbent materials that have been heavily contaminated with this product. Do not reuse them.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

## User Safety Recommendations

Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

Remove clothing immediately if pesticide gets inside. Then wash body thoroughly and put on clean clothing.

Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

## ENVIRONMENTAL HAZARDS

Aldicarb residues may move into shallow ground water under certain conditions. The appended Environmental Precautions Booklet should be read and understood prior to making applications. If there are any questions, contact Bayer CropScience at 1-866-99BAYER (1-866-992-2937).

**DECOMPOSITION AND MOVEMENT IN SOIL:** This product is readily decomposed into harmless residues under most use conditions. However, a combination of permeable and acidic soil conditions, moderate to heavy irrigation and/or rainfall, use of 20 or more pounds per acre, and soil temperature below 50°F (10°C) at application time tend to reduce degradation and promote movement of residues to groundwater. If the above describes your local use conditions and groundwater in your area is used for drinking, do not use this product without first contacting Bayer CropScience at 1-866-99BAYER (1-866-992-2937).

**TOXIC TO FISH, BIRDS, AND WILDLIFE:** This product is toxic to fish, birds, aquatic invertebrates and other wildlife. Birds feeding on exposed granules in treated areas may be killed. Cover or incorporate granules in spill areas.

Run-off from treated areas may be hazardous to fish in neighboring areas. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Apply this product only as specified on this label.

**NOTICE:** Under the Endangered Species Act, it is a Federal Offense to use any pesticide in a manner that results in the death of a member of an endangered species.

This Act protects Attwater's Greater Prairie Chicken in the Texas counties of Aransas, Austin, Brazoria, Colorado, Galveston, Goliad, Harris, Refugio, and Victoria.

Prior to making applications in these counties, the user must determine that this species is not located in or immediately adjacent to the area to be treated. If the user is in doubt whether or not the above named endangered species may be affected, he should contact either the regional U.S. Fish & Wildlife Service office (Endangered Species Specialist) or personnel of the State Fish and Game office.

## DIRECTIONS FOR USE

**It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.**

**Read entire label before using this product.**

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations.

## AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

**Exception:** If the product is soil-incorporated or soil-injected, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated. Once the REI has expired, do not allow persons to come into direct contact with treated soil wet as the result of the initial irrigation or rainfall after treatment unless they are wearing the PPE specified below for early entry. PPE required for early entry to treated areas that is permitted under the Worker Protections Standard and that involves contact with anything that has been treated, such as plants, soil, or water is coveralls over short-sleeved shirt and short pants, waterproof gloves, chemical-resistant footwear plus socks, protective eyewear and chemical-resistant headgear for overhead exposure.

Notify workers of the application by warning them orally and by posting warning signs at entrances to treated areas.

## STORAGE AND DISPOSAL

### STORAGE

Store unused TEMIK® brand Aldicarb Pesticide in original container only, in well ventilated clean dry area out of reach of children and animals. Do not store in areas where temperature averages 115°F (46°C) or greater. Do not store in or around the home or home garden.

If container is broken, before cleaning up, put on long-sleeved shirt, full-length trousers, head covering and rubber gloves. Do not get dust or granules on skin or in eyes. Do not breathe dust. Always stand upwind from spill when cleaning up. Sweep up and bury any small spills or excess TEMIK® Aldicarb at least 18 inches deep in soil, isolated from water supplies and food crops.

### PESTICIDE DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide is a violation of federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or The Hazardous Waste representative at the nearest EPA regional office for guidance.

### CONTAINER DISPOSAL

Completely empty container into application equipment. Then dispose of empty container in a sanitary landfill or by incineration, or if allowed by state and local authorities, by open burning. If container is burned, stay away from and do not breathe or contact smoke.

**IN CASE OF TRANSPORTATION OR WAREHOUSE EMERGENCY INVOLVING A SPILL, FIRE, OR EXPOSURE, CALL 1-800-334-7577 TWENTY-FOUR HOURS A DAY IN THE USA.**

## GENERAL INFORMATION

### READ ENTIRE LABEL BEFORE USING THIS PRODUCT.

TEMIK® brand 15G Aldicarb Pesticide controls certain insects, mites, and nematodes. When applied into moist soil at planting and/or postemergence the active ingredient aldicarb is rapidly absorbed by roots and translocated to all parts of the plant. Rainfall or irrigation soon after application will ensure prompt uptake of aldicarb; however, if irrigation is necessary care should be taken not to over irrigate to reduce the potential for residues leaching to ground water. Control often lasts more than six weeks varying with growing conditions, rate of use, and pests. Crop yields are usually increased with treatments of TEMIK® brand 15G Aldicarb Pesticide.

## GENERAL CAUTIONS

Use TEMIK® brand 15G Aldicarb Pesticide only in accordance with label directions, warnings, and cautions. **DO NOT USE ON ANY CROP NOT LISTED ON THIS LABEL OR SUPPLEMENTAL LABELING AS ANY RESIDUES REMAINING MAY BE ILLEGAL OR HARMFUL. DO NOT STORE OR USE IN OR AROUND THE HOME OR HOME GARDEN.** Use higher rates on heavy organic or clay soils. Do not exceed the maximum label rate. Applications at higher rates or at more frequent intervals than is stated on the label may result in illegal residues in crop, meat, and milk.

Make side-dress applications close enough to plants to allow good uptake by the roots without injury to the plants from root pruning. In irrigated areas, follow application with irrigation within one week. If alternate furrows are irrigated after side-dress application, TEMIK® brand 15G Aldicarb Pesticide and water must be on the same side of the plant row.

Calibrate and adjust application equipment to insure proper rate and accurate placement. Clean application equipment thoroughly after use. For any left over material, see instructions for STORAGE AND DISPOSAL in this booklet.

Deep disc any spills at row ends immediately to prevent birds from feeding on exposed granules.

## GENERAL GROUND WATER LIMITATIONS

- Observe Environmental Hazard precautions regarding Decomposition and Movement in Soil, and carefully follow Directions For Use.
- In fields having soils with less than 15% field moisture holding capacity, special care must be taken not to over-irrigate, since substantial over-irrigation promotes the leaching of chemicals.
- Do not apply within 50 feet of any drinking water well with the exception of Florida where more restrictive setbacks apply.
- Do not wash, load or empty application equipment near any well, as this practice is a potential source of ground water contamination.
- Do not apply this product in the states of Connecticut, Maine, Massachusetts, New Hampshire, New Jersey, New York, Pennsylvania, Rhode Island, Vermont or Wisconsin, or in Del Norte or Humboldt counties in California or in Curry County, Oregon.
- For State Specific Ground Water Limitations, see the *Environmental Precautions and Soil Type Restriction Tables*.
- Apply only between March 1 and September 1 when used in California.

## ROTATIONAL CROPS RESTRICTION

Do not plant any crop not listed on this label in soil treated with TEMIK® brand 15G Aldicarb Pesticide within 10 months after the last application with the following exceptions:

### Six-month plantback restriction

- Do not plant wheat or barley within 6 months after last application.
- Do not plant bulb crops (such as onions or garlic) within 6 months after the last application.
- Do not plant brassica crops (such as broccoli or cabbage) within 6 months after the last application.

### Eight-month plantback restriction

- Do not plant corn within 8 months of last application.
- Do not plant melons within 8 months after last application.
- Do not plant other cucurbits (such as cucumbers and squash) within 8 months after last application.
- Do not plant fruiting vegetables (such as tomatoes or eggplant) within 8 months after the last application.

## MEAT AND MILK

- Do not allow livestock to graze in treated areas before harvest.

## COMPATIBILITY

Pesticidal activity of TEMIK® brand 15G Aldicarb Pesticide is not affected by normal applications of fertilizers or other pesticides. Its effectiveness may be reduced or lost if applied with alkaline materials such as lime. To minimize potential exposure hazards, do not mix TEMIK® aldicarb with other materials before application.

## DIRECTIONS FOR USE AND RECOMMENDED APPLICATIONS

To provide maximum performance and to minimize hazard to birds, work granules into or cover with soil to a depth of 2 inches or more. Deep disc any spills at row ends immediately to prevent birds from feeding on granules. When a range of rates are recommended, use the higher rate if pest infestations are expected to be severe.

---

---

### CITRUS (Oranges, Grapefruit, Lemons and Limes Only)

PESTS CONTROLLED	POUNDS PER ACRE (18 TO 25 FT ROW SPACING)	RECOMMENDED APPLICATION
Citrus Nematode (Except Florida)	33	Apply just prior to or during the spring flush of foliage growth in a band along the dripline on both sides of tree row. Spread granules uniformly and immediately work into the soil or cover with soil to a depth of 2 inches or more;  <b>OR</b> shank granules 2 to 3 inches deep using 3 to 6 shanks on 12 inch centers. The width of each band should equal 1/4 the tree row spacing.
Aphids Mites: Citrus Red Citrus Rust Texas Citrus Whiteflies	33	Apply as above, or apply in irrigation furrows using 2 shanks per furrow.

### Required Application Equipment

Application of TEMIK® brand 15G Aldicarb Pesticide to citrus must be made only with granular applicators which use Positive Displacement Metering Units such as: Pounds Motor Co., Gandy® Orbit-Air®, Gandy® Orbit-Flow®, Valmar Airflow, and Horstine Farmery Microband™, or Accu-Shot®.

### RESTRICTIONS AND PRECAUTIONS FOR CITRUS

- Not registered for use on Citrus in California.
- Apply only between January 1 and April 30 when used in Florida.
- Do not make more than one application per year.
- Do not allow livestock to graze in treated areas.
- Do not harvest lemons within 30 days of application.
- Do not exceed 33 pounds per acre for all applications to citrus.

## COTTON

Crop & Time Of Application	Pests Controlled	Pounds/Acre	Ounces/1000 Feet of Row	Recommended Application
<b>COTTON</b> At planting	Aphids, Thrips	3.5 to 5 (Except Texas, Oklahoma and New Mexico)	40" row spacing 4.5 to 6	Drill granules just below seed line or place in seed furrow and cover with soil. If seeds and TEMIK® are hill-dropped, TEMIK® rates may be reduced by one-half.
		2 to 5 (Texas, Oklahoma, and New Mexico Only)	2.5 to 6	
	Fleahoppers, Leafminers, Mites, Overwintering boll weevil (adults feeding on foliage), Plant bugs, including Lygus	4 to 7	5 to 8.5	
	Nematodes (Suppression)	4 to 7 (California Only)	5 to 8.5	Drill granules just below seed line or place in seed furrow and cover with soil.
	Nematodes	3.5 to 10 (Except Arizona and California)	4.5 to 12	Drill granules just below seed line or place in seed furrow and cover with soil. If rate exceeds 7 pounds per acre apply granules in a 4 to 6-inch band and work into the soil or cover with soil. Plant seed in or above treated zone.
	Nematodes	14 to 27 (Arizona Only)	17 to 33	Apply granules in a 4 to 6-inch band 3 to 6 inches below seed line or cover with 3 to 6 inches of soil. Plant seed above treated zone.
At first squaring	Leafhoppers, Fleahoppers, Nematodes (suppression)	7 to 14	8.5 to 17	Side-dress granules 8 to 16 inches to one side of the plant row and 2 to 6 inches deep (usually at or below bottom of water furrow).
	Mites, Boll weevil, Plant bugs, including Lygus, Nematodes (suppression)	14 to 20 (Except California)	17 to 24.5	
	Mites, Plant bugs, including Lygus, Nematodes	14 (California Only)	17	
From squaring through early bloom	Cotton leaf perforator, Nematodes (suppression)	14	17	Apply before populations reach an economic threshold. Apply for control of moderate populations. Only suppression may be expected for heavy populations of sweet potato whitefly, especially the poinsettia (B) strain (silver leaf).
	Nematodes	14 (California Only)	17	
	Whiteflies	14 to 20 (Except California)  14 (California Only)	17 to 24.5  17	

### RESTRICTIONS AND PRECAUTIONS FOR COTTON

- Do not exceed a total of 21 pounds per acre per season in California or 33 pounds per acre per season in other states for all applications to cotton.
- Apply only between March 1 and September 1 when used in California.
- Do not make more than one at planting application and one postemergence application per crop.
- Do not apply within 90 days of harvest.
- Do not feed cotton forage to livestock or allow livestock to graze in treated area.
- Treatments in excess of 7 pounds per acre made directly in the seed furrow may delay plant emergence and reduce plant stand.

## DRY BEANS

Crop & Time Of Application	Pests Controlled	Pounds/Acre	Ounces/1000 Feet of Row	Recommended Application
<b>DRY BEANS</b> At planting	Seedcorn maggot	3.5 to 5 (Michigan Only)	22" to 48" <u>row spacing</u> 4.0 to 5.5	Apply granules in seed furrow and cover with soil.
	Aphids	5 to 7	5.5 to 7.5	Drill granules 2 to 3 inches below seed line OR, 2 to 3 inches to side of seed row and 2 to 3 inches deep. <b>OR</b>
	Leafhoppers Mexican bean beetle Mites	7 to 14 (Except California)	7.5 to 15	Granules can be placed in seed furrow if the rate does not exceed 5 pounds per acre.
		7 (California Only)	7.5	
	Nematodes	7 to 14 (Except California)	7.5 to 15	Apply granules in a 6-inch band and immediately work into the soil or cover with soil. Plant seed into treated zone.
		7 (California Only)	7.5	Where furrow irrigation is employed soon after planting, drill granules 3 to 4 inches deep and 3 inches from seed row on water furrow side.

### RESTRICTIONS AND PRECAUTIONS FOR DRY BEANS

- Apply only between March 1 and September 1 when used in California.
- Do not make more than one application per crop.
- Do not harvest within 90 days of application.
- Do not feed green forage, hay or straw to livestock.
- Do not allow livestock to graze in treated areas before harvest.
- Do not use green pods as food for humans.
- Do not exceed a total of 7 pounds per acre per season in California and 14 pounds per acre per season in other states.
- Treatments in excess of 5 pounds per acre made directly to the seed furrow may delay plant emergence and reduce plant stand.

## PEANUTS

Crop & Time Of Application	Pests Controlled	Pounds/Acre	Ounces/1000 Feet of Row	Recommended Application
<b>PEANUTS</b> At planting	Thrips	7 to 14	<u>36" row spacing</u> 7.5 to 15	Apply granules in seed furrow and cover with soil. In Southwest use high rate only.
	Nematodes	14 to 20	15 to 22	Apply granules in a 6 to 12 inch band and work into the soil or cover with soil to a depth of 2 to 4 inches. Plant seed into treated zone.

### RESTRICTIONS AND PRECAUTIONS FOR PEANUTS

- Do not make more than one application per crop.
- Do not harvest within 90 days of application.
- Do not hog-off treated fields.
- Do not allow livestock to graze in treated areas before harvest.
- Do not feed hay or vines to livestock.
- Do not exceed a total of 20 pounds per acre.

## PECANS

Crop & Time Of Application	Pests Controlled	Pounds/Acre	Ounces/1000 Feet of Row	Recommended Application
<b>PECANS</b> (Southeast U.S. and Arizona)	Aphids	17 to 33 (California Only) 33 to 47 (Arizona Only)	Not Applicable	Apply as a 4 to 6-foot band along dripline on both sides of tree row by spreading granules uniformly and immediately working into soil, OR by shanking 2-3 inches into soil on 12-inch centers.
<b>Producing Trees</b> During period from bud break to nut set	Aphids Mites	33 to 67 (except California, Florida and Arizona)		
One to two weeks before bud break	Suppression of Pecan leaf phylloxera	33 (Florida only)		
<b>Newly Transplanted Trees</b> (1-5 years old) During period of bud break	Aphids Mites	Dosage Ounces <u>TEMIK®</u> <u>15G/Tree</u> 5 to 20	Not Applicable	Apply as a side-dress to individual trees by spreading the granules uniformly around the tree and immediately working into the soil to a depth of 2-3 inches.
One to two weeks before bud break	Bud moth Suppression of Pecan leaf phylloxera			

### RESTRICTIONS AND PRECAUTIONS FOR PECANS

- Do not make more than one application per year.
- Do not allow livestock to graze in treated areas.
- Do not harvest foliage or hay from treated areas.
- In California and Florida do not exceed a total of 33 pounds per acre.
- In Arizona do not exceed a total of 47 pounds per acre.
- In other states do not exceed a total of 67 pounds per acre.



## POTATOES

Crop & Time Of Application	Pests Controlled	Pounds/Acre	Ounces/1000 Feet of Row	Recommended Application
<b>POTATOES</b> <b>At planting or within one week prior to planting. Do not apply after planting.</b>  <b>FOR USE ONLY IN:</b> Florida and the Pacific Northwest <sup>1</sup> (Idaho, Montana, Oregon and Washington) and  <b>COUNTIES IN THE FOLLOWING STATES:</b>  <b>UTAH:</b> Beaver, Boxelder, Cache, Carbon, Davis, Duchesne, Iron, Millard, Plute, Salt Lake, Sanpete, Sevier, Uintah, Utah, Wasatch, Washington and Weber  <b>NEVADA:</b> Humboldt, Pershing	Flea Beetles	20	21	Apply granules with the seed pieces in the planting furrow and cover with soil.
	Aphids Colorado potato beetle Leafhoppers	14 to 20	14.5 to 21	
	Nematodes (Except Golden Nematode and Root-knot nematodes). Verticillium wilt has been suppressed where nematode populations have been controlled. Root-knot nematodes are suppressed. Despite suppression of root-knot nematode, tuber quality may not be adequately protected. If root-knot nematode is a severe economic pest, the use of other control measures should also be employed.	20	21	Apply granules with the seed pieces in the planting furrow and cover with soil  <p style="text-align: center;"><b>OR</b></p> apply granules in a 6-inch band and work into the soil or cover with soil into a depth of 4 inches. Plant seed pieces into the treated zone.

### Required Application Equipment

All applications must be made with Positive Displacement Applicators such as Horstine Microband® or Gandy PDM® or with other Positive Displacement Applicators approved under the Bayer CropScience Equipment Certification Program. The end user (i.e., applicator) must be certified through the Bayer CropScience Stewardship program.

<sup>(1)</sup> See General Use Restrictions for ground water protection.

### RESTRICTIONS AND PRECAUTIONS FOR POTATOES

- Do not harvest within 100 days of an application for potatoes grown in Florida and within 150 days of an application for potatoes grown in the Pacific Northwest (Idaho, Montana, Oregon, Washington and certain counties in Nevada and Utah).
- Do not apply after planting.
- Do not make more than one application per crop. If replanting is necessary, do not retreat the second crop with TEMIK®.
- Do not allow livestock to graze in treated areas before harvest.
- Do not exceed 20 pounds per acre.
- In the Pacific Northwest (Idaho, Montana, Oregon and Washington as well as certain counties in Utah and Nevada), do not use furrow irrigation on TEMIK® treated potatoes.
- Do not use this product on potatoes in any state other than those immediately listed above.

## SORGHUM

Crop & Time Of Application	Pests Controlled	Pounds/Acre	Ounces/1000 Feet of Row	Recommended Application
<b>GRAIN SORGHUM</b> At planting	Nematodes	7 (Except Texas and Oklahoma)	<u>36" row spacing</u> 7.5	Apply granules in seed furrow and cover with soil.
		3.5 to 7 (Texas and Oklahoma Only)	4 to 7.5	
	Greenbug Chinch Bug	7	<u>36" row spacing</u> 7.5	

### RESTRICTIONS AND PRECAUTIONS FOR SORGHUM

- Do not make more than one application per year.
- Do not harvest within 90 days of application.
- Do not feed green forage to livestock.
- Do not exceed a total of 7 pounds per acre.
- Do not allow livestock to graze in treated areas before harvest.

## SOYBEANS

Crop & Time Of Application	Pests Controlled	Pounds/Acre	Ounces/1000 Feet of Row	Recommended Application
<b>SOYBEANS</b> At planting	Mexican bean beetle Thrips Suppression of three cornered alfalfa hopper	5 to 10	30" to 48" <u>row spacing</u> 5.5 to 11	Drill granules 2 to 3 inches below seed line  <b>OR</b> 2 to 3 inches to the side of the seed row and 2 to 3 inches deep. Granules can be applied to seed furrow if rate does not exceed 5 pounds per acre.
	Nematodes	10 to 20	11 to 22	Apply granules in a 6 to 8-inch band and immediately work into the soil or cover with soil. Plant seed into treated zone.

### RESTRICTIONS AND PRECAUTIONS FOR SOYBEANS

- Do not make more than one application per crop.
- Do not harvest within 90 days of application.
- Do not allow livestock to graze in treated areas before harvest.
- Do not feed green forage, hay, or straw to livestock.
- Do not exceed a total of 20 pounds per acre.
- Treatments in excess of 5 pounds per acre made directly to the seed furrow may delay plant emergence and reduce plant stand.

## SUGAR BEETS

Crop & Time Of Application	Pests Controlled	Pounds/Acre	Ounces/1000 Feet of Row	Recommended Application
<b>SUGAR BEETS</b> At planting or within one week before planting.	Nematodes	27 to 33  (Except California)	<u>22" row spacing</u> 18 to 22	Apply granules in a 4 to 6 inch band and immediately work into the soil or cover with soil to a depth of 2 to 4 inches. Plant seed into or above treated zone;  <b>OR</b> where furrow irrigation is employed for seed germination, drill granules 3 to 4 inches deep and 3 inches from seed row on water furrow side.
		14  (California Only)	9.5	
At planting	Aphids	7 to 14	4.5 to 9.5	Drill granules 1 to 3 inches below seedline. Granules can be placed in seed furrow if rate does not exceed 7 pounds per acre.  Repeat applications will be required for continued protection against virus vectors (aphids and leafhoppers)
	Leafminers Leafhoppers	14 to 20  (Except California)	9.5 to 13.5	
		14  (California Only)	9.5	
	Sugar beet root maggot	7 to 14	4.5 to 9.5	Apply granules in a 2 to 3 inch band over seed row and immediately work into the soil or cover with soil to a depth of 2 to 4 inches;  <b>OR</b> where furrow irrigation is employed for seed germination, drill granules 2 inches deep and 2 inches from seed row on water furrow side.  For the 7 lb. rate, apply granules in a 1 to 2 inch band in front of the press wheel as the furrow is closing.
At planting plus postemergence (split applications)	Nematodes	14 to 20 at planting and 14 to 20 at postemergence  (Except California)	9.5 to 13.5 at planting and 9.5 to 13.5 at postemergence	<b>AT PLANTING:</b> Apply granules in a 4 to 6-inch band and immediately work into the soil or cover with soil to a depth of 2 to 4 inches. Plant seed into or above treated zone;  <b>OR</b> where furrow irrigation is employed for seed germination, drill granules 3 to 4 inches deep and 3 inches from seed row on water furrow side. <b>POSTEMERGENCE:</b> Apply granules to both sides of plant row and immediately work into the soil or cover with soil to a depth of 2 to 4 inches;  <b>OR</b> for furrow irrigation side-dress granules 4 to 8 inches to water furrow side of plant row at furrow depth. Irrigate soon after application. Apply within 60 days after planting.
		14 at planting and 14 at postemergence  (California Only)	9.5 at planting and 9.5 at postemergence	

## SUGAR BEETS (Continued)

Crop & Time Of Application	Pests Controlled	Pounds/Acre	Ounces/1000 Feet of Row	Recommended Application
Postemergence Do not make any post - emergence application if 27 to 33 lbs./acre were applied at planting or one week before planting. Do not make more than one at planting application and two postemergence applications per crop.	Sugar beet root maggot	7 to 14	4.5 to 9.5	Apply granules to both sides of plant row and immediately work into the soil or cover with soil to a depth of 2 to 4 inches;  <b>OR</b> for furrow irrigation side-dress granules 4 to 8 inches to water furrow side of plant row at furrow depth. Irrigate soon after application. Apply within 60 days after planting.
	Aphids	7 to 14	4.5 to 9.5	Apply as above. A repeat application may be required for continued protection against virus vectors (aphids, leafhoppers).
	Leafminers Leafhoppers	14 to 20 (Except California)	9.5 to 13.5	Apply within 60 days after planting.
		14 (California Only)	9.5	
	Nematodes	27 (Except California)	18	Apply as above. Apply within 60 days after planting. In California, apply within 30 days after planting.
		14 (California Only)	9.5	

### RESTRICTIONS AND PRECAUTIONS FOR SUGAR BEETS

- Apply only between March 1 and September 1 when used in California.
- Do not make more than one at planting application and two postemergence applications per crop.
- Do not apply within 90 days of harvest.
- If tops are to be fed to livestock, do not apply within 120 days of harvest.
- Do not use tops as food for humans.
- Do not exceed a total of 28 pounds per acre per season in California and 33 pounds per acre per season in other states for all applications to sugar beets.
- Treatments in excess of 7 pounds per acre made directly in the seed furrow may delay plant emergence and reduce plant stand.

## SUGARCANE

Crop & Time Of Application	Pests Controlled	Pounds/Acre	Ounces/1000 Feet of Row	Recommended Application
<b>SUGARCANE</b> At planting	Nematodes	14 to 20	<u>60" row spacing</u> 26 to 37	Apply granules in open row on top of newly planted cane and cover immediately with at least 6 inches of soil.

### RESTRICTIONS AND PRECAUTIONS FOR SUGARCANE

- For use in Louisiana only.
- Do not make more than one application per crop.
- Do not harvest within 120 days of application.
- Do not exceed a total of 20 pounds per acre.
- Do not feed cane forage.

## SWEET POTATOES

Crop & Time Of Application	Pests Controlled	Pounds/Acre	Ounces/1000 Feet of Row	Recommended Application
<b>SWEET POTATOES</b> At planting	Nematodes	10 to 20 (Except California)	<u>48" row spacing</u> 15 to 30	Apply granules in a 12-inch band in opened row. Cover immediately with soil by hilling up 8 to 10 inches. Plant in center of treated zone. Apply only with positive displacement applicators such as the Horstine Farmery Microband (see calibration chart at end of label), or the Gandy Orbit.
	Nematodes (suppression)	10 to 14 (California Only)		

### Required Application Equipment

All applications must be made with Positive Displacement Applicators such as Horstine Microband® or Gandy PDM® or with other Positive Displacement Applicators.

### RESTRICTIONS AND PRECAUTIONS FOR SWEET POTATOES

- Do not make more than one application per crop.
- Do not harvest within 120 days of application.
- Do not feed sweet potato vines to livestock.
- Do not exceed a total of 20 pounds per acre.

**SEE PREHARVEST AND GRAZING LIMITATIONS ELSEWHERE IN THIS LABEL.**

## TEMIK® BRAND 15G (GYPSUM) ALDICARB PESTICIDE CALIBRATION GUIDE

APPROXIMATE APPLICATOR SETTINGS\* FOR TEMIK® BRAND 15G ALDICARB PESTICIDE (GYPSUM) AT 4, 5 & 6 MPH

NOTE: This calibration chart is applicable only to TEMIK® brand 15G Aldicarb Pesticide in this container which is formulated for use on a gypsum carrier. All rates are APPROXIMATIONS and must be confirmed by using a calibration tube (See Calibration Chart provided with tube). Calibration tubes are available from your TEMIK® brand 15G Aldicarb supplier.

### TEMIK BRAND 15G (GYPSUM) ALDICARB PESTICIDE CALIBRATION GUIDE

#### TYPE OF GRANULAR APPLICATOR

#### POUNDS TEMIK® 15G PER ACRE FOR VARIOUS ROW SPACING

ROW SPACING					GANDY GAUGE SETTING			JOHN DEERE MAX EMERGE GAUGE SETTING**			JOHN DEERE MAX EMERGE II GAUGE SETTING			CASE-IH SERIES 800/900 GAUGE SETTING****		
					MPH			MPH			MPH			MPH		
22"	34"	36"	40"	48"	4	5	6	4	5	6	4	5	6	4	5	6
5.5	3.5	3.3	3	2.5	12	13	14	6***	7***	8	8	9	11	1/6.0	1/6.5	1/7.5
7.3	4.7	4.4	4	3.3	13	15	16	7***	9	10	10	12	13	1/7.0	1/7.5	1/8.5
9.1	5.9	5.5	5	4.2	14	16	18	9	11	13	12	14	15	1/8.0	1/8.5	1/9.5
10.9	7.1	6.7	6	5.0	16	18	19	11	13	15	13	15	18	1/9.0	1/9.5	2/1.0
12.7	8.2	7.8	7	5.8	17	19	21	12	15	18	15	18	21	1/9.5	2/0.5	2/2.0
14.5	9.4	8.9	8	6.7	18	20	22	14	17	20	16	20	23	2/0.0	2/2.0	2/3.5
16.4	10.6	10.0	9	7.5	19	21	23	16	19	23	18	22	27	2/1.0	2/3.0	2/5.0
18.2	11.8	11.1	10	8.3	20	22	24	17	21	25	20	25	29	2/2.0	2/4.0	2/6.5
21.8	14.1	13.3	12	10.0	22	24	26	20	25	28	23	29	35	2/3.5	2/7.0	3/0.0
25.5	16.5	15.6	14	11.7	23	26	29	24	28	30	27	35	41	2/5.5	2/8.5	3/4.5
	18.8	17.8	16	13.3	25	28	31	26	29	32	31	39	45	2/7.5	3/2.5	3/8.0
	21.2	20.0	18	15.0	26	30	33	28	31	33	36	44	48	3/0.5	3/6.0	4/2.0
	23.5	22.2	20	16.7	28	31	34	30	32	35	40	46	50	3/2.5	3/9.0	4/5.5
			24	20.0	31	34	37	32	34	37	45	49	53	3/7.5	4/6.0	5/3.0
			27	22.5	33	36	40	33	36	38	48	52	56	4/2.5	5/0.0	5/9.5

\*All settings for one hopper box outlet per row

\*\*Odd Numbered Housing (0 – 45)

\*\*\*For John Deere gauge setting below 8 there will be great variations in flow rate. A gauge setting of 6 is the lowest recommended setting.

\*\*\*\*Gate/Dial

### HORSTINE FARMERY MICROBAND\*

(Using one 3/16" rotor per outlet, part #094006)

#### POUNDS TEMIK® 15G (GYPSUM) PER ACRE FOR VARIOUS ROW SPACINGS

ROW SPACING	SPROCKET** COMBINATION					DRIVER/DRIVEN					
	19-34	22-34	25-34	28-34	25-28	22-22	28-25	34-28	34-25	34-22	34-19
22"	5.50	6.40	7.30	8.20	8.90	10.00	11.20	12.10	13.60	15.40	17.80
30"	4.00	4.60	5.30	5.90	6.50	7.30	8.10	8.80	9.90	11.20	13.00
36"	3.30	3.90	4.40	5.00	5.40	6.10	6.80	7.30	8.20	9.40	10.90
40"	3.00	3.50	4.00	4.50	4.80	5.50	6.10	6.60	7.40	8.40	9.80
48"	2.50	2.90	3.30	3.70	4.00	4.60	5.10	5.50	6.20	7.10	8.20
<b>RATIO:</b>	<b>0.550</b>	<b>0.640</b>	<b>0.730</b>	<b>0.820</b>	<b>0.890</b>	<b>1.000</b>	<b>1.120</b>	<b>1.210</b>	<b>1.360</b>	<b>1.545</b>	<b>1.789</b>

• The Microband is based on positive metering. Settings do not need to be changed with different tractor speeds.

• All settings for one hopper box outlet per row.

• For alternative dose rates, different width rotors can be fitted. For example, a 3/8" wide rotor will deliver twice the rates indicated in the above table.

\*Calibrations based upon use of a Horstine Farmery or Canaan 29.5 inch diameter Landwheel drive.

\*\*Sprocket numbers, example: 19 - 34, indicate number of teeth on the DRIVER and DRIVEN Sprockets.

# TEMIK® BRAND 15G (CORNCOB) ALDICARB PESTICIDE CALIBRATION GUIDE

APPROXIMATE APPLICATOR SETTINGS\* FOR TEMIK® BRAND 15G ALDICARB PESTICIDE (CORNCOB) AT 4, 5 & 6 MPH

NOTE: This calibration chart is applicable only to TEMIK® brand 15G Aldicarb Pesticide in this container which is formulated for use on a corncob carrier. All rates are APPROXIMATIONS and must be confirmed by using a calibration tube (See Calibration Chart provided with tube). Calibration tubes are available from your TEMIK® 15G Aldicarb supplier.

## TEMIK® BRAND 15G (CORNCOB) ALDICARB PESTICIDE CALIBRATION GUIDE

### TYPE OF GRANULAR APPLICATOR

#### POUNDS TEMIK® 15G PER ACRE FOR VARIOUS ROW SPACING

ROW SPACING					GANDY GAUGE SETTING			JOHN DEERE MAX EMERGE GAUGE SETTING**			JOHN DEERE MAX EMERGE II GAUGE SETTING			CASE-IH SERIES 800/900 GAUGE SETTING***		
					MPH			MPH			MPH			MPH		
22"	34"	36"	40"	48"	4	5	6	4	5	6	4	5	6	4	5	6
5.5	3.5	3.3	3	2.5	22	24	27	14	16	18	22	25	29	2/2.0	2/4.5	2/6.5
7.3	4.7	4.4	4	3.3	25	28	30	17	19	21	27	32	37	2/4.5	2/7.5	3/0.0
9.1	5.9	5.5	5	4.2	28	31	33	19	21	23	32	38	44	2/7.5	3/2.0	3/5.0
10.9	7.1	6.7	6	5.0	30	33	35	21	23	25	37	44	48	3/1.0	3/5.5	4/0.0
12.7	8.2	7.8	7	5.8	32	35	37	22	25	26	42	47	50	3/4.5	3/9.0	4/5.0
14.5	9.4	8.9	8	6.7	34	37	39	24	26	28	46	50	52	3/7.0	4/4.0	5/0.0
16.4	10.6	10.0	9	7.5	35	38	41	25	27	29	48	51	54	4/0.0	4/8.0	5/5.0
18.2	11.8	11.1	10	8.3	37	40	43	26	28	30	50	52	55	4/4.0	5/2.0	6/0.0
21.8	14.1	13.3	12	10.0	39	43	47	28	30	33	52	55	57	5/0.0	6/0.0	7/0.0
25.5	16.5	15.6	14	11.7	42	46	50	29	32	35	54	56	60	5/6.5	6/7.5	8/0.0
	18.8	17.8	16	13.3	44	49	54	31	34	37	56	59	62	6/4.0	7/7.5	9/0.0
	21.2	20.0	18	15.0	47	52	59	33	36	38	57	61	65	6/9.5	8/5.0	10/0.0
	23.5	22.2	20	16.7	49	56	66	34	37	40	59	63		7/7.5	9/2.5	11/2.5
			24	20.0	54	66		37	40	44	62			9/0.0	11/0.5	
			27	22.5	59			38	42		65			10/1.0		

\*All settings for one hopper box outlet per row

\*\*Odd Numbered Housing (0 – 45)

\*\*\*Gate/Dial

### HORSTINE FARMERY MICROBAND\*

(Using two 3/16" rotors per outlet, part #094006)

#### POUNDS TEMIK® 15G (CORNCOB) PER ACRE FOR VARIOUS ROW SPACINGS

ROW SPACING	SPROCKET** COMBINATION						DRIVER/DRIVEN				
	19-34	22-34	25-34	28-34	25-28	22-22	28-25	34-28	34-25	34-22	34-19
22"	5.50	6.40	7.30	8.20	8.90	10.00	11.20	12.10	13.60	15.40	17.80
30"	4.00	4.60	5.30	5.90	6.50	7.30	8.10	8.80	9.90	11.20	13.00
36"	3.30	3.90	4.40	5.00	5.40	6.10	6.80	7.30	8.20	9.40	10.90
40"	3.00	3.50	4.00	4.50	4.80	5.50	6.10	6.60	7.40	8.40	9.80
48"	2.50	2.90	3.30	3.70	4.00	4.60	5.10	5.50	6.20	7.10	8.20
<b>RATIO:</b>	<b>0.550</b>	<b>0.640</b>	<b>0.730</b>	<b>0.820</b>	<b>0.890</b>	<b>1.000</b>	<b>1.120</b>	<b>1.210</b>	<b>1.360</b>	<b>1.545</b>	<b>1.789</b>

• The Microband is based on positive metering. Settings do not need to be changed with different tractor speeds.

• All settings for one hopper box outlet per row.

• For alternative dose rates, different width rotors can be fitted. For example, a 3/4" wide rotor will deliver twice the rates indicated in the above table.

\*Calibrations based upon use of a Horstine Farmery or Canaan 29.5 inch diameter Landwheel drive.

\*\*Sprocket numbers, example: 19 - 34, indicate number of teeth on the DRIVER and DRIVEN Sprockets.

## IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

**CONDITIONS:** The directions for use of this product are believed to be adequate and should be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience. All such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** BAYER CROPSCIENCE MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. BAYER CROPSCIENCE DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

**LIMITATIONS OF LIABILITY:** THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE'S ELECTION, THE REPLACEMENT OF PRODUCT.

TEMIK is a registered trademark of Bayer.



**Bayer CropScience LP**  
P.O. Box 12014, 2 T.W. Alexander Drive  
Research Triangle Park, North Carolina 27709  
1-866-99BAYER (1-866-992-2937)  
<http://www.bayercropscience.us>

12/16/03



## **RESTRICTED USE PESTICIDE**

### **ACUTE TOXICITY AND GROUND WATER CONTAMINATION**

For retail sale to and use only by Certified Applicators or persons under the direct supervision of a Certified Applicator, and only for those uses covered by the Certified Applicator's certification.

# **TEMIK<sup>®</sup> brand 15G Aldicarb Pesticide**

EPA Reg. No. 264-330

## **Environmental Precautions and Soil Type Restriction Tables**



Refer to the container label for additional use precautions and directions.



**Bayer CropScience**

Bayer CropScience LP  
P.O. Box 12014, 2 T.W. Alexander Drive  
Research Triangle Park, North Carolina 27709  
1-866-99BAYER (1-866-992-2937)  
<http://www.bayercropscience.us>

TEMIK is a registered trademark of Bayer.

Contents	Page
----------	------

ENVIRONMENTAL PRECAUTIONS & GENERAL USE RESTRICTIONS .....	1
SPECIFIC GROUNDWATER LIMITATIONS FOR FLORIDA .....	1

**SOIL TYPE TABLES BY STATE**

ALABAMA .....	2
COLORADO .....	2
DELAWARE .....	2
GEORGIA .....	2
ILLINOIS .....	3
INDIANA .....	3
IOWA .....	3
KANSAS .....	3
KENTUCKY .....	3
LOUISIANA .....	3
MARYLAND .....	4
MICHIGAN .....	4
MINNESOTA .....	4
MISSISSIPPI .....	5
MISSOURI .....	5
MONTANA .....	5
NEBRASKA .....	5
NORTH CAROLINA .....	6
NORTH DAKOTA .....	6
OHIO .....	6
SOUTH CAROLINA .....	6
SOUTH DAKOTA .....	7
TENNESSEE .....	7
VIRGINIA .....	7
WEST VIRGINIA .....	7
WYOMING .....	8

## ENVIRONMENTAL PRECAUTIONS GENERAL USE RESTRICTIONS

**AGRICULTURAL CHEMICALS HAVE THE POTENTIAL TO MOVE INTO SHALLOW GROUND WATER.  
THE FOLLOWING RESTRICTIONS HAVE BEEN DEVELOPED TO PROTECT DRINKING WATER SUPPLIES.**

**DO NOT APPLY WITHIN 50 FEET OF ANY DRINKING WATER WELL TO MINIMIZE CONTAMINATION BY SURFACE RUNOFF.  
MORE STRINGENT RESTRICTIONS MAY BE REQUIRED, AS DISCUSSED BELOW.**

Do not wash, load, or empty application equipment near any well, as this practice is a potential source of ground water contamination . In fields having soils with less than 15% field moisture holding capacity, special care must be taken not to over-irrigate, since substantial over-irrigation promotes the leaching of chemicals.

Contact your state pesticide regulatory authority for further information on state requirements for the use of this product.

Some states have or may be developing more restrictive regulations regarding the use and application of TEMIK®. Follow all state regulations restricting the use and application of this product, including limitations on applications near drinking water sources. In all cases, the more restrictive requirements must be followed. It is the responsibility of the applicator to document the construction of wells claimed not to be shallow.

### STATE SPECIFIC GROUND WATER LIMITATIONS

**Do not apply this product in the states of Connecticut, Maine, Massachusetts, New Hampshire, New Jersey, New York, Pennsylvania, Rhode Island, Vermont or Wisconsin, or in Del Norte or Humboldt counties in California or in Curry County, Oregon.**

**FOLLOW THE LISTED "ADDITIONAL RESTRICTIONS" if the following conditions are present:**

STATE	SOILS FOR WHICH RESTRICTIONS APPLY	ADDITIONAL RESTRICTIONS															
AL, CO, DE, GA, KS, KY, LA, MD, MO, MS, NC, SC, TN, VA, WV	Loamy sand or sand surface soils <u>and</u> subsoils <u>with an</u> average organic matter in the upper 12 inches of less than 2% by weight.  See <i>SOIL TYPE RESTRICTION TABLES</i> for specific soil types.	If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within <u>300 feet</u> of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table.															
IA, IL, IN, MI, MN, MT, ND, NE, OH, SD, WY	Sandy loam, loamy sand, or sand surface soils, <u>and</u> loamy sand or sand subsoils, <u>with an</u> average organic matter in the upper 12 inches of less than 2% by weight.  See <i>SOIL TYPE RESTRICTION TABLES</i> for specific soil types.	If a vulnerable soil is present and the water table is less than 25 feet, do not apply within <u>500 feet</u> of a drinking water well unless it is known or reasonably believed based upon an authoritative source that such wells are either cased to 100 feet below the ground level or a minimum of 30 feet below the water table.															
FL	All Soils	<p>In Florida, do not exceed a total of 33 pounds per acre. State regulations require that TEMIK® not be applied within <u>300 feet</u> of any drinking water well. In addition, TEMIK® aldicarb cannot be used on Florida citrus within 1000 feet of a drinking water well regardless of depth of water table, when soils (such as those listed below) have a permeability rate greater than 20 inches per hour with an available water capacity less than 0.06 in all layers to a depth of 80 inches as identified by the U.S.D.A. Soil Conservation Service, unless it is known or reasonably believed based on authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. The U.S.D.A. Soil Conservation Service which serves your county can tell you if the soils in your grove(s) fall within this category.</p> <table><tr><td>• Adamsville</td><td>• Candler</td><td>• Neilhurst</td><td>• Paola</td><td>• St. Lucie</td></tr><tr><td>• Archbold</td><td>• Cassia</td><td>• Orsino</td><td>• Satellite</td><td>• Tavares</td></tr><tr><td>• Astatula</td><td>• Lake</td><td>• Palm beach</td><td></td><td></td></tr></table> <p>In Florida, it is the responsibility of the applicator to document the construction of wells claimed not to be shallow. This must consist of (a) a copy of the well construction permit issued by the appropriate water management district; or (b) a statement certified as to accuracy by a licensed well contractor.</p>	• Adamsville	• Candler	• Neilhurst	• Paola	• St. Lucie	• Archbold	• Cassia	• Orsino	• Satellite	• Tavares	• Astatula	• Lake	• Palm beach		
• Adamsville	• Candler	• Neilhurst	• Paola	• St. Lucie													
• Archbold	• Cassia	• Orsino	• Satellite	• Tavares													
• Astatula	• Lake	• Palm beach															

**SOIL TYPE RESTRICTION TABLE****FOLLOW THE LISTED ADDITIONAL RESTRICTIONS IF THE SOIL TYPES ARE PRESENT IN YOUR STATE:****ALABAMA**

If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within **300 feet** of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table.

alaga	blanton	corolla	foxworth	lakeland	pactolus
alpin	bruno	crevasse	fripp	latonia	plummer
americus	buncombe	duckston	garcon	leon	saffell
bassfield	chipleay	eustis	gorgas	nugent	scranton
bigbee	chipola	flomaton	jones	osier	tarboro

**COLORADO**

If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within **300 feet** of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table.

bangston	cotopaxi	grieves	littlebear	peetz	stecum
bankard	coyet	grimm	lonetree	pescar	tassel
barcus	crestman	grimstone	luning	pinequest	thoroughfare
baroid	crustown	gunbarrel	mathis	platte	tipper
batterson	dailey	hiwan	maybell	redcreek	tipperary
bijou	dix	inavale	medano	resort	tivoli
blakeland	dunday	ipage	mespun	roswell	tomah
boel	dunul	ironsprings	mido	ruedloff	tomichi
brad	dwyer	juget	mirror lake	ruhe	trail
breece	eachuston	julesburg	moenkopie	ryark	tricera
bresser	ecklund	kandaly	moosed	san isabel	tullock
canlodore	eghelm	kassler	mosca	sawcreek	valent
cascajo	ellicott	kerber	munjor	schamber	valentine
chappell	els	kettle	mysten	schooner	wigton
chaseville	elsmere	kippen	nakai	sheppard	willwood
clark fork	falcon	laird	nesda	siebert	yetull
columbine	farb	laney	newcomb	southace	zeona
comad	farisita	las animas	osgood	space city	
corlena	gilcrest	layoint	ouray	sphinx	
corlett	gracot	legault	pando	spool	
costilla	gretdivid	lincoln	patna	stapleton	

**DELAWARE**

If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within **300 feet** of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table.

downer	galestown	hurlock	osier	rumford
evesboro	hammonton	kenansville	plummer	
fort mott	hooksan	klej	pocomoke	

**GEORGIA**

If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within **300 feet** of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table.

americus	duckston	kershaw	osier	valdosta
bigbee	echaw	kingsferry	ousley	wateree
blanton	eustis	kureb	penney	
boulogne	flomaton	lakeland	plummer	
buncombe	foxworth	leon	pottsburg	
cainhoy	fripp	lynn haven	ridgeland	
centenary	gorgas	mandarin	ridgewood	
chipleay	hurricane	meadowbrook	sapelo	
chipola	johns	molena	scranton	

**SOIL TYPE RESTRICTION TABLE (continued)****ILLINOIS**

If a vulnerable soil is present and the water table is less than 25 feet, do not apply within **500 feet** of a drinking water well unless it is known or reasonably believed based upon an authoritative source that such wells are either cased to 100 feet below the ground level or a minimum of 30 feet below the water table.

alганsee	carmi	eleva	landes	plainfield	stonelick
alvin	carr	flagler	lanier	princeton	thetford
beavercreek	casco	grable	lorenzo	robby	watseka
billett	chelsea	granby	matherton	rodman	zumbro
bloomfield	chute	hodge	minneiska	saffell	
boone	coloma	homer	morocco	sarpy	
boyer	dickinson	hononegah	oakville	seafield	
brems	disco	hoopeston	oshtemo	sparta	
burkhardt	el dara	lamont	plainbo	spinks	

**INDIANA**

If a vulnerable soil is present and the water table is less than 25 feet, do not apply within **500 feet** of a drinking water well unless it is known or reasonably believed based upon an authoritative source that such wells are either cased to 100 feet below the ground level or a minimum of 30 feet below the water table.

abscota	bruno	granby	morocco	riverdale	tedrow
alганsee	carmi	gravelton	moundhaven	robby	thetford
alvin	casco	hanna	nesius	rodman	tyner
billett	chatterton	homer	newton	saugatuck	wasepi
bloomfield	chelsea	hononegah	oakville	seafield	watseka
bobtown	coloma	hoopeston	ormas	shipshe	zaborosky
boyer	conotton	junius	oshtemo	sisson	zadog
brady	conrad	kosciusko	ouiatenon	sparta	
branch	desker	landes	plainfield	spinks	
brems	dickinson	leoni	princeton	stockland	
bronson	elston	maumee	prochaska	stonelick	

**IOWA**

If a vulnerable soil is present and the water table is less than 25 feet, do not apply within **500 feet** of a drinking water well unless it is known or reasonably believed based upon an authoritative source that such wells are either cased to 100 feet below the ground level or a minimum of 30 feet below the water table.

backbone	carr	estherville	hoopeston	oesterle	watseka
billett	chelsea	finchford	klum	perks	zenor
boone	chute	flagler	lamont	salida	
brady	dickinson	fruitfield	landes	sarpy	
buckney	dickman	granby	lilah	sparta	
burkhardt	elrick	hawick	montieth	toolesboro	

**KANSAS**

If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within **300 feet** of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table.

aline	dwyer	goodnight	kanza	pratt	thurman
bankard	els	gracemore	las animas	sarpy	tivoli
dillwyn	elsmere	haxtun	likes	schamber	valent
dix	goltry	inavale	lincoln	simeon	valentine

**KENTUCKY**

If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within **300 feet** of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table.

bruno	flomaton	lakin	potomac	smithdale
crevasse	iuka	molena	saffell	

**LOUISIANA**

If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within **300 feet** of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table.

alaga	bigbee	darden	hackberry	lakeland	osier
bassfield	bruno	eustis	haggerty	latonia	palm beach
betis	cheniere	felicity	hainesville	lotus	peveto
bienville	crevasse	flo	kenney	nugent	

**SOIL TYPE RESTRICTION TABLE (continued)****MARYLAND**

If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within **300 feet** of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table.

aura	duckston	galestown	klej	newhan	runclint
carteret	evesboro	hammonton	lakeland	osier	
colts neck	fisherman	hooksan	leetonia	plummer	
downer	fort mott	hurlock	leon	pocomoke	
dragston	fripp	ingleside	molena	remlik	

**MICHIGAN**

If a vulnerable soil is present and the water table is less than 25 feet, do not apply within **500 feet** of a drinking water well unless it is known or reasonably believed based upon an authoritative source that such wells are either cased to 100 feet below the ground level or a minimum of 30 feet below the water table.

abbaye	chesaning	granby	mahtomedi	padus	spinks
abscota	coloma	grattan	mancelona	pelkie	springlake
adams	covert	graycalm	manitowish	pemene	st. ignace
alcona	croswell	grayling	matherton	pence	steuben
alганsee	cunard	guardlake	maumee	pequaming	summersville
allouez	deer park	halfaday	mecosta	perrin	summersville
alpena	deerton	hodenpyl	menahga	peshekee	sundog
alvin	desker	ingalls	millersburg	pipestone	tedrow
arcadian	dickinson	ionia	monico	plainfield	tekenink
arkport	dixboro	ishpeming	montcalm	rapson	thetford
au gres	duel	kalkaska	morocco	richter	trimountain
banat	dunbridge	karlin	nadeau	ridgeville	tyre
battlefield	east lake	kawbawgam	newton	riverdale	velvet
bixler	eastport	keweenaw	nottawa	rodman	vilas
blue lake	eleva	kingsville	oakville	rousseau	wainola
bohemian	elston	kiva	oconto	rubicon	waiksa
boyer	ensign	klacking	ocqueoc	sarona	wallace
brady	esau	koontz	omega	saugatuck	wasepi
branch	evart	kosciusko	onota	sayner	watseka
brems	fabius	landes	ormas	seafield	winterfield
bronson	finch	leelanau	oshtemo	shawano	zeba
channing	freesoil	leoni	otisco	shelldrake	zimmerman
chatham	gilchrist	liminga	ottokee	sisson	
chelsea	gladwin	lode	paavola	sparta	

**MINNESOTA**

If a vulnerable soil is present and the water table is less than 25 feet, do not apply within **500 feet** of a drinking water well unless it is known or reasonably believed based upon an authoritative source that such wells are either cased to 100 feet below the ground level or a minimum of 30 feet below the water table.

abscota	copaston	hiwood	nymore	sugarbush
alban	cormant	hoopeston	oesterle	torning
alганsee	cowhorn	insula	omega	ulen
alvin	cromwell	kanaranzi	osakis	venlo
andrusia	deer park	karlstad	pierz	verndale
anoka	dickinson	kost	plainbo	wamduska
arvilla	dickman	lamont	plainfield	watseka
beavercreek	egeland	langola	poppleton	waupaca
becker	eleva	lasa	quetico	wawina
belchester	emmert	leola	redby	winterfield
bergkeller	evart	lilah	renshaw	zenor
billet	faunce	lino	rosholt	zimmerman
boone	finchford	lohnes	rosy	zippel
burkhardt	flagler	maddock	salida	zumbro
burnsville	fossum	mahtomedi	sanburn	
carmi	friendship	marquette	sartell	
caryville	gotham	meehan	serden	
chelsea	granby	menahga	shawano	
chetek	graycalm	mesaba	sioux	
claire	hamar	minneiska	soderville	
cloquet	hawick	nemadji	sparta	
conic	hecla	northfield	spinks	

**SOIL TYPE RESTRICTION TABLE (continued)****MISSISSIPPI**

If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within **300 feet** of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table.

alaga	bigbee	eustis	latonia	pactolus	st. lucie
bassfield	bruno	flomaton	nugent	plummer	
beulah	crevasse	lakeland	osier	saffell	

**MISSOURI**

If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within **300 feet** of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table.

alvin	carr	eustis	maldeen	scotco	
beulah	clana	finchford	plainfield	shelldrake	
bruno	crevasse	hodge	sandbur	sparta	
canalou	diehlstadt	landes	sarpy	wideman	

**MONTANA**

If a vulnerable soil is present and the water table is less than 25 feet, do not apply within **500 feet** of a drinking water well unless it is known or reasonably believed based upon an authoritative source that such wells are either cased to 100 feet below the ground level or a minimum of 30 feet below the water table.

alberton	calicott	glendive	mirror	sachett	trey
ambrant	castner	hanly	moiese	sawcreek	tullock
apeldorn	cheadle	haskill	nelson	scravo	turnercrest
ashbon	cheyenne	haverdad	nemote	selle	tusler
ashuelot	chinook	haverly	nesda	selway	twilight
assinniboine	clark fork	havrelon	nippt	seroco	upsata
bangston	cohagen	hiwan	oceanet	sheege	usine
bankard	colburn	jugson	oraid	shingle	utica
banks	comad	kalsted	ovando	shook	valentine
baxendale	como	kirby	pend orielle	shurley	vebar
bearmouth	cowood	krause	poin	splitro	victor
beaverton	cozberg	laporte	redcreek	stecum	virgelle
beenom	dast	larim	rencot	tally	wabek
beisigl	dayschool	lihen	rentsac	tamely	wilsonville
belain	dominic	lisk	rhame	tassel	woodgulch
biglake	dwyer	lolopeak	ridge	terry	woodside
blackhall	elkner	lone rock	riedel	tinsley	worland
blanchard	elmira	lowercreek	riverside	tipperary	yellowbay
bonner	ervide	macfarlane	rivra	tomichi	yetull
brandenburg	flasher	manning	rochester	travessilla	zeona
branham	fleak	mccaffery	ryell	travson	
busby	glaciercreek	mcilwaine	ryorp	treble	
cabba	glenberg	mires	sacheen	trembles	

**NEBRASKA**

If a vulnerable soil is present and the water table is less than 25 feet, do not apply within **500 feet** of a drinking water well unless it is known or reasonably believed based upon an authoritative source that such wells are either cased to 100 feet below the ground level or a minimum of 30 feet below the water table.

alda	chappell	gannett	meadin	talmo
almeria	cheyenne	glenberg	munjor	tassel
anselmo	craft	gothenburg	o'Neill	thurman
bankard	dailey	haverson	ord	trelona
barney	darr	hedville	orpha	treon
bigwinder	dickinson	hersh	ortello	tryon
blendon	dix	hoffland	orwet	tuthill
boel	doger	holt	phiferson	valent
bolent	draknab	inavale	pivot	valentine
boone	duda	ipage	platte	whitelake
brunswick	dunday	laird	ronson	wildhorse
busher	dwyer	las animas	sarpy	
calamus	eckley	lute	schamber	
canyon	els	manter	selia	
carr	elsmere	marlake	shingle	
cass	fonner	mckelvie	simeon	

## SOIL TYPE RESTRICTION TABLE (continued)

### NORTH CAROLINA

If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within **300 feet** of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table.

alaga	centenary	galestown	mandarin	pocomoke	tomahawk
alpin	charleston	immokalee	molena	potomac	valhalla
baymeade	chiple	johns	nansemond	pottsborg	wake
biltmore	conetoe	kenansville	newhan	remlik	wakulla
blanton	corolla	klej	norfolk	rimini	wando
bojac	dragston	kureb	ona	rumford	wateree
buncombe	duckston	lakeland	osier	seabrook	
butters	echaw	leon	ostin	st. lucie	
cainhoy	eustis	lillington	ousley	suncook	
candor	foreston	louisburg	pactolus	tarboro	
carteret	fripp	lynn haven	plummer	toisnot	

### NORTH DAKOTA

If a vulnerable soil is present and the water table is less than 25 feet, do not apply within **500 feet** of a drinking water well unless it is known or reasonably believed based upon an authoritative source that such wells are either cased to 100 feet below the ground level or a minimum of 30 feet below the water table.

appam	chinook	fossum	lohnes	seroco	verendrye
arvilla	claire	glendive	maddock	sioux	virgelle
aylmer	coe	hamar	manning	stirum	wabek
banks	cohagen	hanly	metigoshe	telfer	wamduska
bantry	cormant	haskill	minnewaukan	tinsley	yecross
beisigl	dast	havrelon	osakis	trembles	yetull
binford	desart	hecla	poppleton	trey	zeona
blackhall	egeland	inkster	renshaw	tusler	
blanchard	ekalaka	karlsruhe	rhame	twilight	
brandenburg	falsen	ladner	ruso	ulen	
breien	flasher	lemert	schaller	vebar	
cabba	fleak	lihen	serden	venlo	

### OHIO

If a vulnerable soil is present and the water table is less than 25 feet, do not apply within **500 feet** of a drinking water well unless it is known or reasonably believed based upon an authoritative source that such wells are either cased to 100 feet below the ground level or a minimum of 30 feet below the water table.

abscota	bronson	elnora	lakin	perrin	spinks
alганsee	casco	galen	landes	pinegrove	stafford
alvin	coloma	granby	lanier	plainfield	steinsburg
arkport	colonie	hazleton	maumee	princeton	stonelick
barkcamp	conotton	homer	oakville	rodman	tedrow
bixler	dekalb	hoopeston	oshtemo	schaffenaker	tyner
boyer	dixboro	ionia	otisville	sisson	watertown
brady	dunbridge	kingsville	ottokee	sparta	

### SOUTH CAROLINA

If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within **300 feet** of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table.

alaga	carteret	eustis	louisburg	pactolus	tomahawk
alpin	centenary	foreston	lynn haven	plummer	wakulla
baratari	charleston	foxworth	molena	pocomoke	wando
baymeade	chiple	fripp	mouzon	ridgeland	wateree
blanton	corolla	johns	murad	rimini	witherbee
brogdon	crevasse	kenansville	nansemond	rosedhu	
buncombe	dragston	kershaw	newhan	scranton	
butters	duckston	kureb	olanta	seabrook	
cainhoy	echaw	lakeland	osier	seewee	
candor	elloree	leon	ousley	tarboro	



## SOIL TYPE RESTRICTION TABLE (continued)

### SOUTH DAKOTA

If a vulnerable soil is present and the water table is less than 25 feet, do not apply within **500 feet** of a drinking water well unless it is known or reasonably believed based upon an authoritative source that such wells are either cased to 100 feet below the ground level or a minimum of 30 feet below the water table.

almeria	chinook	flasher	lute	rockoa	tryon
alwilda	cohagen	fleak	maddock	ronson	tusler
anselmo	claire	fossum	manning	sarpy	tuthill
arvilla	craft	gannett	manter	schamber	twilight
assinniboine	dailey	glenberg	marlake	serden	ulen
aylmer	dix	glendive	mawer	seroco	valent
bankard	doger	hamar	mccaffery	shingle	valentine
banks	dogiecreek	hanly	meadin	simeon	vebar
bantry	duda	hecla	minnewaukan	sioux	venlo
beisigl	dunday	henkin	munjor	stirum	wabek
blackhall	dwyer	holt	murdo	storla	whitelake
blendon	eckley	hopdraw	o'Neill	talmo	yecross
boel	egeland	inavale	orton	tassel	zeona
butche	ekalaka	ipage	orwet	telfer	
cabba	els	kirby	platte	thurman	
canyon	elsmere	ladner	renshaw	travessilla	
cass	evridge	las animas	reva	trembles	
chappell	fedora	lihen	rhame	trey	

### TENNESSEE

If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within **300 feet** of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table.

barkcamp	bruno	crevasse	nugent
bigbee	buncombe	eustis	potomac

### VIRGINIA

If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within **300 feet** of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table.

alaga	catpoint	fripp	lakin	osier	seabrook
assateague	chiple	galestown	leetonia	ousley	spessard
aura	conetoe	johns	leon	pactolus	tarboro
biltmore	corolla	kenansville	magotha	plummer	wakulla
bojac	dragston	klej	millrock	pocomoke	wando
buncombe	drall	kureb	molena	remlik	wateree
camocca	duckston	lakehurst	nansemond	rumford	
carteret	fisherman	lakeland	newhan	schaffenaker	

### WEST VIRGINIA

If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within **300 feet** of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table.

drall	lakin	landes	leetonia	potomac	schaffenaker
-------	-------	--------	----------	---------	--------------

## SOIL TYPE RESTRICTION TABLE (continued)

### WYOMING

If a vulnerable soil is present and the water table is less than 25 feet, do not apply within **500 feet** of a drinking water well unless it is known or reasonably believed based upon an authoritative source that such wells are either cased to 100 feet below the ground level or a minimum of 30 feet below the water table.

ashbon	chinook	glenton	meadowlake	rentsac	tipper
adilis	clarkelen	grenoble	means	rivra	tipperary
alcova	comad	grieves	millburne	roget	tolman
alflack	conpeak	grimm	milok	rohonda	travessilla
anasazi	cordes	grimstone	mishak	roxal	travson
anselmo	corlett	gunbarrel	mosroc	ryark	trekona
apeldorn	cotha	handran	motoqua	ryorp	trembles
assinniboine	cotopaxi	hanks	mudray	sachett	treoff
bankard	cowestglen	hanly	nathale	sawcreek	treon
banks	cowood	hapjack	nelson	schamber	trook
barcus	coyet	haterton	nesda	schooner	troutville
barnum	cragosen	haverdad	newfork	scravo	tullock
baroid	crestman	haverly	nizina	seedskadee	turnback
barrett	crownest	havermom	norriston	sharland	turnercrest
bearmouth	cryluha	hazton	norte	sheege	twilight
beaverton	curabith	hechtman	oceanet	shingle	valent
bigwin	dailey	henrieville	onason	shoshone	valentine
bigwinder	dast	herbman	orpha	shurley	vible
bilycreek	derrick	hiland	otterson	sobson	vonason
blackhall	devore	hiwan	ouray	southace	walcott
bobtail	dix	hoot	pahlow	southfork	walden
bodorumpe	draknab	huguston	peetz	space city	wall
bosler	dunday	irigul	pendergrass	splitro	wardboro
bottle	dunul	julesburg	pepton	spool	wendover
boyle	dwyer	kandaly	pescar	stecum	wetmore
branham	eckley	koonich	phiferson	sudworth	whaley
breece	edlin	labou	pilotpeak	sunup	wiggleton
brownsto	elk mountain	lamarsh	pinequest	taluce	willwood
bruja	ellicott	lambman	poin	tassel	wilsonville
busher	emblem	laney	preatorson	tasselman	wint
butche	enos	laporte	pugsley	teagulf	worf
byrnie	ethelman	lariat	quealman	teewinot	workfa
calicott	eyre	larim	quealy	tellman	worfman
cambarge	farisita	las animas	randman	tenorio	worfstone
canwall	farson	lazeare	ratake	terada	worland
carbol	feltner	littlebear	redbank	terro	yetull
cass	fleak	littsan	redcreek	terry	youjay
castner	folavar	lone rock	redfeather	tetonville	zeomont
cathedral	frontier	macfarlane	rekop	theedle	zeona
cestnik	gas creek	manter	relsob	thermopolis	
chappell	glenberg	mathers	rencot	tieside	
cheadle	glendive	mccaffery	reno	tine	

TEMIK is a registered trademark of Bayer.



**Bayer CropScience**

**Bayer CropScience LP**  
**P.O. Box 12014, 2 T.W. Alexander Drive**  
**Research Triangle Park, North Carolina 27709**  
**1-866-99BAYER (1-866-992-2937)**  
**<http://www.bayercropscience.us>**

12/16/03