



Bayer Environmental Science
A Division of Bayer CropScience, LP
5000 CentreGreen Way, Suite 400
Cary, North Carolina 27513
1-866-99BAYER (1-866-992-2937)

TEMPO® SC Ultra Insecticide

For control of Exterior Fleas.

EPA Reg. No. 432-1363

FIFRA 2(ee) Recommendation

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE(S) OF ALABAMA, ALASKA, ARIZONA, ARKANSAS, CALIFORNIA, COLORADO, CONNECTICUT, DELAWARE, DISTRICT OF COLUMBIA, FLORIDA, GEORGIA, IDAHO, ILLINOIS, INDIANA, IOWA, KANSAS, KENTUCKY, LOUISIANA, MAINE, MICHIGAN, MINNESOTA, MISSISSIPPI, MISSOURI, MONTANA, NEBRASKA, NEVADA, NEW JERSEY, NORTH CAROLINA, NORTH DAKOTA, OHIO, OKLAHOMA, PENNSYLVANIA, RHODE ISLAND, SOUTH CAROLINA, SOUTH DAKOTA, TENNESSEE, UTAH, VERMONT, VIRGINIA, WASHINGTON, WEST VIRGINIA, WISCONSIN, AND WYOMING.

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its label.

APPLICATION FOR OUTDOOR TREATMENT OF FLEAS

Use this product by application as a perimeter treatment, spray or broadcast application to exterior surfaces of buildings, lawns, grounds, and ornamental plantings. This product controls designated pests on landscape and recreational turf. Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes. Not for use on golf course turf.

USE RATE • 8 to 16 ml (0.27 to 0.54 fl oz) per 1,000 sq ft - See Table 1 for mixing directions. • Use high rate for severe infestations.

HOW TO APPLY (Also see APPLICATIONS FOR OUTDOOR/ PERIMETER PESTS)

- Perimeter Treatment: Treat area adjacent buildings in a band up to 10 feet wide. Also treat the building foundation to a height of 2 to 3 feet in areas where pests are active or can find entrance.
- Apply just enough dilution to adequately cover the area without excessive dripping or runoff. Volume can vary depending on the surface type treated. For example, mulch requires more spray than bare soil. If mulch is heavy, rake it back to expose soil before treating.
- Avoid contact with treated surfaces until dry.

WHEN TO APPLY

- When pests are seen or found • Re-apply every 7 to 10 days if needed.

Apply as a mist or coarse low pressure spray at the rate of 8 to 16 ml (0.27 to 0.54 fl oz) of this product per 1,000 sq ft of surfaces. Apply in a sufficient volume of water to provide adequate coverage but do not allow run-off to occur.

APPLICATION RESTRICTIONS

Do not water the treated area to the point of run-off.

Do not make applications during rain.

Do not apply directly to impervious horizontal surfaces such as sidewalks, driveways, and patios except as a spot, pin stream, or crack and crevice treatment.

During application, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters, or surface waters.

All outdoor applications, if permitted elsewhere on the product label, must be limited to spot, pin stream, or crack and crevice treatments only, except for the following permitted uses:

1. Applications to soil or vegetation, as listed on this label, around structures;
2. Applications to lawns, turf, and other vegetation, as listed on this label;
3. Applications to the side of a building, up to a maximum height of 3 feet above grade;
4. Applications to underside of eaves, soffits, doors, or windows permanently protected from rainfall by a covering, overhang, awning, or other structure;
5. Applications around potential pest entry points into buildings, when limited to a surface band not to exceed one inch in width;
6. Applications made through the use of a coarse, low pressure spray to only those portions of surfaces that are directly above bare soil, lawn, turf, mulch or other vegetation, as listed on this label, and not over an impervious surface, drainage or other condition that could result in runoff into storm drains, drainage ditches, gutters, or surface waters, in order to kill occasional invaders or aggregating pests. Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to fresh water and estuarine fish and invertebrates. Use with care when applying in areas adjacent to any body of water. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters, or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

This pesticide is highly toxic to bees exposed to direct treatment. Do not apply or allow it to drift to crops or weeds on which bees are actively foraging.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.
For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937).

*As with any crop-protection product, always read and follow label directions.
For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*