

SUPPLEMENTAL LABELING

RESTRICTED USE PESTICIDE

Toxic to fish and aquatic organisms. For retail sale to and use only by certified applicators, or persons under their direct supervision and only for those uses covered by the certified applicator's certification.

FLUOXASTROBIN	GROUP	11	FUNGICIDE
BIFENTHRIN	GROUP	3	INSECTICIDE

TEPERA® PLUS FUNGICIDE / INSECTICIDE

Supplemental Label to Add DIRECTIONS FOR USE on CANOLA

This supplemental label expires March 31, 2021 and must not be used or distributed after this date.

For Agricultural Uses

ACTIVE INGREDIENTS:	% BY WT.
Fluoxastrobin: [(1E)-[2-[[6-(2-Chlorophenoxy)-5-fluoro-4-pyrimidinyl]oxy]phenyl] (5,6-dihydro-1,4,2-dioxazin-3-yl) methanone-O-methyloxime].....	9.26%
Bifenthrin: *	9.07%
OTHER INGREDIENTS:	81.67%
TOTAL:	100.0%

This product contains 0.83 pounds of fluoxastrobin per gallon

This product contains 0.82 pounds of bifenthrin per gallon

*Cis isomers 97% minimum, trans isomers 3% maximum.

KEEP OUT OF REACH OF CHILDREN CAUTION / PRECAUCIÓN

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION. READ THE LABEL AFFIXED TO THE CONTAINER FOR **TEPERA® PLUS FUNGICIDE** BEFORE APPLYING. USE OF **TEPERA PLUS FUNGICIDE** ACCORDING TO THIS LABELING IS SUBJECT TO THE USE PRECAUTIONS, RESTRICTIONS, AND LIMITATIONS IMPOSED BY THE LABEL AFFIXED TO THE CONTAINER FOR **TEPERA PLUS FUNGICIDE (EPA Reg. No. 66330-431)**.

073118V005SU

CANOLA, Crambe, and Rapeseed		
Pest Control	Product Use Rate fl oz/A (lb ai/A)	Application Directions
At Plant Soilborne and Seedling Disease Control		
Rhizoctonia Root Rot	9.2 – 12.5 (0.06 – 0.08 fluoxastrobin + 0.06- 0.08 bifenthrin)	See IN-FURROW AND BANDING APPLICATION RATES table for corresponding use rate per 1000 row feet based on crop row spacing. Apply as a 5- to 7-inch band (T-band) over an open furrow or in-furrow with the seed.
At Plant Insect Control		
Army cutworm Armyworm species Cutworm species Grape colaspis Grubs Root aphids Seed corn beetle Seed corn maggot Stalkborer Sugarcane beetle True armyworm Wireworm	9.2 (0.06 fluoxastrobin + 0.06 bifenthrin)	See IN-FURROW AND BANDING APPLICATION RATES table for corresponding use rate per 1000 row feet based on crop row spacing. Apply as a 5- to 7-inch band (T-band) over an open furrow or in-furrow with the seed.
Foliar Disease Control		
Alternaria Blackspot Blackleg*	12.5 (0.08 fluoxastrobin + 0.08 bifenthrin)	For optimum results, begin applications preventively and continue as needed on a 7- to 14-day interval. Use the higher specified rates and shorter interval when disease pressure is high.
Foliar Insect Control		
Aphids Armyworms Cutworms Diamondback Moth Loopers Other Lepidopterous Larvae Flea Beetle Flea Hopper Grasshopper Plant Bug Stink Bugs Seedpod Weevil Thrips Whitefly	5.1 – 6.2 (0.033 - 0.04 fluoxastrobin + 0.033 - 0.04 bifenthrin)	Apply in a minimum of 2 gallons finished spray per acre by air or in a minimum of 10 gallons per acre with ground equipment. When applying by air, 1 to 2 quarts of emulsified oil may be substituted for 1 to 2 quarts of water in the finished spray. Thorough coverage is essential to achieve control.
RESTRICTIONS:		
<ul style="list-style-type: none"> Do not apply more than 12.5 fl oz (0.08 lb ai fluoxastrobin + 0.08 lb ai bifenthrin) of product per acre per single application. Do not apply more than 0.30 lb ai of fluoxastrobin per acre per year including at plant and foliar applications of all fluoxastrobin-containing products. Do not apply more than 0.08 lb ai per season including at-plant and foliar applications of all bifenthrin containing products. There is a maximum number of 2 applications per acre per year and a minimum interval of 14 days between applications. Do not apply within 35 days of harvest. 		
*Not registered for use in California.		

Produced For:
ARYSTA LIFESCIENCE NORTH AMERICA, LLC
15401 Weston Parkway, Suite 150
Cary, NC 27513

Arysta LifeScience and the Arysta LifeScience logo are registered trademarks of Arysta LifeScience Corporation. TEPERA is a registered trademark of an Arysta LifeScience Group Company.
©2018 Arysta LifeScience Group Company.