

SUPPLEMENTAL LABELING

FLUTRIAFOL	Group	3	Fungicide
------------	-------	---	-----------



**TOPGUARD® Fungicide**  
EPA Reg. No. 279-3557

For barley, sweet corn, and rice as a rotational crop.

***THIS SUPPLEMENTAL LABEL EXPIRES ON January 31, 2023 AND MUST NOT BE USED OR DISTRIBUTED AFTER THIS DATE***

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Read the entire label. This label must be in the possession of the user at the time of application. Read the label affixed to the container for TOPGUARD Fungicide before applying. Use of TOPGUARD Fungicide according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the TOPGUARD Fungicide container.

**Crop Rotation:** Fields treated with TOPGUARD Fungicide may be planted to the following crops and plant back intervals:

Crop	Plant back Interval
Alfalfa, barley, field corn; field corn grown for seed; popcorn; sweet corn; cotton; cucurbits; fruiting vegetables; grapes; hops; Brassica leafy vegetables crop group 5; leafy vegetables, except Brassica, crop group 4; peanut; pome fruit; grain sorghum; soybean; stone fruit; strawberry; sugar beet; tomato; tree nuts; triticale; or wheat	Immediately
Rice	30 days
All other crops	Prohibited

SUPPLEMENTAL LABELING

**CROP USE DIRECTIONS**

**Use Directions for Sweet Corn (Foliar Applications)**

<b>Diseases</b>	<b>Product Rate (fl oz /A)</b>	<b>Application Directions</b>	<b>Application Interval (Days)</b>
<p>Gray Leaf Spot (<i>Cercospora zeae-maydis</i>)</p> <p>Southern Corn Leaf Blight (<i>Cochliobolus heterostrophus</i>)</p> <p>Northern Corn Leaf Blight (<i>Setosphaeria turcica</i>)</p> <p>Rust, Common (<i>Puccinia sorghi</i>)</p> <p>Southern Rust (<i>Puccinia polysora</i>)</p> <p>Eyespot (<i>Aureobasidium zeae</i>)</p> <p>Anthracnose Leaf Blight (<i>Colletotrichum graminicola</i>)</p> <p>Northern Corn Leaf Spot (<i>Cochliobolus carbonum</i>)</p>	<p style="text-align: center;">7 - 14</p> <p style="text-align: center;">(0.057-0.114 lb ai/A)</p>	<p>Apply preventatively or when conditions are favorable for disease development. One repeat application may be made.</p> <p>Apply no later than growth stage R4 (early dough stage).</p> <p>For aerial application, use a minimum of 2 gallons of spray solution per acre.</p>	<p style="text-align: center;">7</p>
<p><b>Restrictions:</b></p> <ul style="list-style-type: none"> <li>• No single application may exceed 14 fl oz of product (0.114 lb ai) /A.</li> <li>• Do not apply more than 2 applications per year.</li> <li>• Do not apply more than 28 fl oz of product (0.228 lb ai)/A per year.</li> <li>• Do not apply more than 0.228 lb ai of flutriafol/A per year.</li> <li>• An adjuvant may be used with TOPGUARD Fungicide prior to the V8 corn growth stage and after the VT corn growth stage.</li> <li>• Restricted Entry Interval (REI): The REI for sweet corn is 3 days.</li> <li>• Preharvest Interval: The PHI for sweet corn is 7 days.</li> </ul>			

SUPPLEMENTAL LABELING

**Use Directions for Barley**

<b>Diseases</b>	<b>Product Rate (fl oz/A)</b>	<b>Application Directions</b>	<b>Application Interval (Days)</b>
Rust <i>(Puccinia spp.)</i>  Powdery Mildew <i>(Blumeria graminis)</i>  Leaf and Glume Blotch <i>(Septoria spp.)</i>	10 - 14  (0.081-0.114 lb ai/A)	Apply preventatively or when conditions are favorable for disease development. Repeat as necessary if conditions are favorable for disease development.  For aerial application, use a minimum of 2 gallons of spray solution per acre.	7
<b>Suppression:</b> Fusarium Head Blight <i>(Fusarium spp.)</i>	10 - 14  (0.081-0.114 lb ai/A)	Apply at anthesis (Feekes Growth Stage 10.51). Spray equipment nozzles should be properly set up to ensure good coverage of the head.  For aerial application, use a minimum of 2 gallons of spray solution per acre.	7
<b>Restrictions:</b> <ul style="list-style-type: none"> <li>• No single application may exceed 14 fl oz of product (0.114 lb ai)/A.</li> <li>• Do not apply more than 2 applications per year.</li> <li>• Do not apply more than 28 fl oz of product (0.228 lb ai)/A per year.</li> <li>• Do not apply more than 0.228 lb ai of flutriafol/A per year.</li> <li>• Restricted Entry Interval (REI) is 12 hours.</li> <li>• Preharvest Interval: Do not apply within 30 days of harvest when grown for grain, 15 days of harvest grown for hay, zero (0) day PHI when grown for forage.</li> </ul>			

Label code: R-4368 042920 02-21-20

Sold By



FMC Corporation

2929 Walnut Street

Philadelphia, PA 19104