

**THIS RECOMMENDATION IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA.**



**2(ee) Recommendation**

**FOR DISTRIBUTION AND USE ONLY  
IN THE STATE OF NEW YORK**

**It is a violation of Federal law to use this product  
in a manner inconsistent with its labeling.**

**EPA Reg. No.: 279-3596**

**For early season disease protection in spring and winter wheat.**

*THIS 2(ee) RECOMMENDATION EXPIRES ON JUNE 30, 2023 AND MUST NOT BE USED AFTER THIS DATE*

| <b>CROP</b>                         | <b>PEST</b>  | <b>RATE OF APPLICATION</b> |
|-------------------------------------|--|----------------------------|
| <b>Wheat</b><br>(Spring and Winter) | <b>Leaf, Stem and Stripe Rusts</b> ( <i>Puccinia</i> spp.)<br><b>Powdery Mildew</b> ( <i>Blumeria graminis tritici</i> )<br><b>Septoria Leaf and Glume Blotch</b> ( <i>Septoria</i> spp.)<br><b>Tan Spot</b> ( <i>Pyrenophora tritici-repentis</i> ) | <b>3 - 4 fl. oz./A</b>     |

**DIRECTIONS FOR USE:**

Apply Topguard® EQ fungicide in early spring from tillering to jointing growth stages (Feekes stages 3-7) to provide protection against early season leaf diseases. Topguard® EQ fungicide may be tank-mixed with adjuvants, fertilizer, insecticides (such as **Mustang® MAXX** insecticide) or herbicides (such as **Harmony® Extra SG herbicide with TotalSol®** soluble granules) to implement disease, insect and weed management. An additional application of Topguard® EQ fungicide may be required at flag leaf (Feekes 8-9) for control of foliar diseases as specified by product label. Apply in a minimum of 2 gallons per acre for aerial application and 10 gallons per acre by ground application.

**RESTRICTIONS:**

- Do not apply more than 2 applications per year.
- Do not apply more than 0.228 lb. ai of flutriafol/A per year.
- Do not apply more than 0.4 lb. ai of azoxystrobin per acre per year.
- Restricted Entry Interval (REI) is 12 hours.
- Preharvest Intervals (PHI): Do not apply within 30 days of harvest when grown for grain, 15 days of harvest when grown for hay, 7 days of harvest when grown for forage.

**ALL APPLICABLE DIRECTIONS, RESTRICTIONS, WORKER PROTECTION STANDARDS REQUIREMENTS AND PRECAUTIONS ON THE EPA REGISTERED TOPGUARD EQ FUNGICIDE, EPA REG.NO. 279-3596, LABEL MUST BE FOLLOWED. THIS LABEL MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.**