

SUPPLEMENTAL LABELING

FLUTRIAFOL	Group	3	Fungicide
AZOXYSTROBIN	Group	11	Fungicide



TOPGUARD® EQ Fungicide

EPA Reg. No. 279-3596

For Use on Barley, Sweet Corn and Rice as a Rotational Crop.

THIS SUPPLEMENTAL LABEL EXPIRES ON January 31, 2023 AND MUST NOT BE USED OR DISTRIBUTED AFTER THIS DATE

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Read the entire label. This label must be in the possession of the user at the time of application. Read the label affixed to the container for Topguard EQ Fungicide before applying. Use of Topguard EQ Fungicide according to this labeling is subject to the use precautions, restrictions, and limitations imposed by the label affixed to the Topguard EQ Fungicide container.

Crop Rotation Interval: Treated areas may be rotated to the following crops based on plant back intervals outlined in the table below. Rotation to any other crop not listed in the table is not allowed.

Crop	Plant Back Interval
Alfalfa, barley, Brassica leafy vegetables, corn (field, field corn grown for seed, popcorn, and sweet corn), cotton, cucurbits, fruiting vegetables, grain sorghum, grapes, leafy vegetables, peanuts, pome fruit, soybeans, stone fruit, strawberry, sugar beets, tree nuts, triticale and wheat.	0 Days
Rice	30 days

SUPPLEMENTAL LABELING

Use Directions for Sweet Corn

Not for use in CA

Target Diseases	Product Rate (fl oz/A)	Application Directions
Common Rust <i>(Puccinia sorghi)</i> Southern Rust <i>(Puccinia polysora)</i> Anthracnose Leaf Blight <i>(Colletotrichum graminicola)</i> Gray Leaf Spot <i>(Cercospora sorghi)</i> Northern Corn Leaf Blight <i>(Setosphaeria turcica)</i> Northern Corn Leaf Spot <i>(Cochliobolus carbonum)</i> Southern Corn Leaf Blight <i>(Cochliobolus heterostrophus)</i> Eye Spot <i>(Aureobasidium zeae)</i>	5.0 – 7.0	For optimum results begin applications when disease first appears and continue as needed on a 7 to 10 day interval. Use high end of the use rate when disease pressure is high and conditions are favorable for disease development. Apply no later than growth stage R4 (early dough stage). For multiple applications refer to the guidelines under Resistance Management.
<p>Restrictions:</p> <ul style="list-style-type: none"> • Do not exceed 7 fl oz of product/A for a single application. • Do not apply more than 2 applications of product per growing season. • Do not apply more than 14 fl oz/A of product per acre per year. • Do not apply more than 0.228 lb ai of flutriafol per acre per year. • Do not apply more than 2.0 lb ai of azoxystrobin per acre per year. • An adjuvant may be used with Topguard EQ prior to the V8 corn growth stage and after the VT corn growth stage. • Restricted Entry Interval (REI): The REI for sweet corn is 3 days. • Preharvest Interval: Do not apply within 7 days of harvest. 		

SUPPLEMENTAL LABELING

Use Directions for Barley

Not for use in CA

Target Diseases	Product Rate (fl oz/A)	Application Directions
Leaf Rust (<i>Puccinia</i> spp.) Powdery Mildew (<i>Blumeria graminis tritici</i>) Septoria Leaf and Glume Blotch (<i>Septoria</i> spp.) Stem Rust (<i>Puccinia graminis</i>) Stripe Rust (<i>Puccinia striiformis</i>) Tan Spot (<i>Pyrenophora tritici-repentis</i>)	4.0 – 7.0	Apply preventatively or when conditions are favorable for disease development. Repeat as necessary if conditions are favorable for disease development. Do not apply past Feekes 10.54. An adjuvant may be added at recommended rates. For multiple applications refer to the guidelines under Resistance Management.
<p>Restrictions:</p> <ul style="list-style-type: none"> • No single application may exceed 7 fl oz of product/A. • Do not apply more than 2 applications per year. • Do not apply more than 0.228 lb ai of flutriafol per acre per year. • Do not apply more than 0.40 lb ai of azoxystrobin per acre per year. • Restricted Entry Interval (REI): The REI is 12 hours. • Do not apply within 30 days of harvest when grown for grain, 15 days of harvest when grown for hay, 7 days when grown for forage. 		

Label Code: R-4374 042920 02-21-20

Sold By



FMC Corporation
 2929 Walnut Street
 Philadelphia, PA 19104