

Supplemental Labeling

GROUP

3 & 11

FUNGICIDES

TOPGUARD[®]

EQ

FUNGICIDE

For use on almond, brassica leafy vegetables crop group 5, celery and Chinese celery, corn, cotton, cucurbit vegetables, grain sorghum, grapes, leafy vegetables, except brassica, crop group 4, peanuts, pecan, soybeans, stone fruit, strawberry, sugar beets, tomatoes, tree nuts, vegetable, fruiting, group 8-10, wheat, and triticale.

This supplemental label expires on October 1, 2018 and must not be used or distributed after this date.

EPA Reg. No. 279-3596

EPA Est. No. 82978-FRA-001

ACTIVE INGREDIENTS:	By Wt.
Azoxystrobin: methyl (E)-2-{2-[6-(2-cyanophenoxy) pyrimidin-4-yloxy]phenyl}-3-methoxyacrylate*	25.30%
Flutriafol	18.63%
OTHER INGREDIENTS:	56.07%
TOTAL:	100.00%

Contains 1.82 lb. of flutriafol and 2.47 lb. of azoxystrobin active ingredient per gallon.
Suspension Concentrate.
*IUPAC

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Sold By



FMC Corporation
2929 Walnut Street
Philadelphia, PA 19104

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read the entire label. This label must be in the possession of the user at the time of application. Please refer to the TOPGUARD EQ FUNGICIDE container label for additional PRECAUTIONARY STATEMENTS. In addition to the restrictions listed here, the user must follow all application directions, restrictions and precautions otherwise listed on the TOPGUARD EQ FUNGICIDE container label.

Crop Rotation Interval: Treated areas may be rotated to the following crops based on plant back intervals outlined in the table below. Rotation to any other crop not listed in the table is not allowed.

Crop	Plant Back Interval
Brassica leafy vegetables, corn (field, field corn grown for seed, and popcorn), cotton, cucurbits, fruiting vegetables, grain sorghum, grapes, leafy vegetables, peanuts, pome fruit, soybeans, stone fruit, strawberry, sugar beets, tree nuts, triticale and wheat.	0 Days
Sweet corn	180 Days

Use Directions for Cucurbit Vegetables. See below for a list of crops. Not for use in CA

Target Diseases	Product Rate (fl oz/A)	Application Directions
Anthraxnose (<i>Colletotrichum lagenarium</i>) Belly Rot (<i>Rhizoctonia solani</i>) Downy Mildew (<i>Pseudoperonospora cubensis</i>) Gummy Stem Blight (<i>Didymella bryoniae</i>) Leaf Spots (<i>Alternaria</i> spp., <i>Cercospora</i> spp.) Myrothecium Canker (<i>Myrothecium roridum</i>) Plectosporium Blight (<i>Plectosporium tabacinum</i>) Powdery Mildew (<i>Sphaerotheca fuliginea</i> , <i>Erysiphe cichoracearum</i>) Target Spot (<i>Corynespora cas-siicola</i>) Ulocladium Leaf Spot (<i>Ulocladium cucurbitae</i>)	5.0 – 8.0	For Downy Mildew make preventative applications on a 5 – 7 day schedule. For Belly Rot control, the first application should be made at the 1 - 3 leaf crop stage with a second application just prior to vine tip over or 10 - 14 days later whichever occurs first. For all other diseases, apply preventatively or when conditions are favorable for disease development and continue throughout the season at 7 - 14 day intervals. For multiple applications refer to the guidelines under Resistance Management. An adjuvant may be added at recommended rates. Do not tank mix with COC, MSO or silicon adjuvants. Do not tank mix with Malathion, Kelthane®, Thiodan®, Phaser®, Lannate®, Lorsban®, MPede®, Nufos® 4E or Botran®.

List of Cucurbit Vegetables crops: Cantaloupe, Chayote, Chinese-Waxgourd, Cucumber, Gourds, Honeydew Melons, Momordica spp. (bitter melon, balsam apple), Muskmelon, Watermelon, Pumpkin, Squash, Zucchini. Including cultivars and/or hybrids of these crops.

Restrictions:

- No single application may exceed 8 fl. oz. of product/A.
- Do not apply more than 4 applications per year.
- Do not apply more than 0.456 lb a.i. of flutriafol per acre per year.
- Do not apply more than 1.5 lb a.i. of azoxystrobin per acre per year.
- **Restricted Entry Interval (REI):** The REI is 12 hours.
- **Preharvest Interval:** The PHI is 1 day.

WARRANTY DISCLAIMER

FMC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, FMC MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

INHERENT RISKS OF USE

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of FMC or the Seller. All such risks shall be assumed by Buyer and User. Buyer and User agree to hold FMC and the Seller harmless for any claims related to such factors.

LIMITATION OF REMEDIES

To the extent consistent with applicable law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to one of the following, at FMC's election:

- (1) Refund of purchase price paid by buyer or user for product bought, or
- (2) Replacement of amount of product used.

To the extent consistent with applicable law, FMC shall not be liable for consequential, incidental, or special damages or losses.

The terms of the Warranty Disclaimer above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of FMC or the Seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

TOPGUARD and Nufos are registered trademarks of FMC

Aliette and Phaser are registered trademarks of Bayer CropScience.

Ambush and Warrior with Zeon Technology are registered trademarks of Syngenta Group Company.

Lorsban and Kelthane are registered trademarks of Dow AgroSciences, LLC

Lannate is a registered trademark of DuPont Crop Production

MPede is a registered trademark of Mycogen Corporation.

Thiodan is a registered trademark of Universal Crop Protection Alliance, LLC

Botran is a registered trademark of Gowan.