



FIFRA 2(ee) Recommendation

EPA Reg. No. 279-3557

**FOR DISTRIBUTION AND USE ONLY IN:
Texas**

THIS RECOMMENDATION, WHICH CONTAINS ADDITIONAL DIRECTIONS FOR USE, IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA.

This recommendation for TOPGUARD fungicide is valid until June 30, 2023, or until withdrawn, canceled or suspended.

**FOR DISEASE PROTECTION OF EARLY VEGETATIVE STAGE AND GRAIN FILL
STAGE WINTER WHEAT**

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.

THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

CROP	PEST	RATE OF APPLICATION
Winter Wheat	Common and stripe rust (<i>Puccinia spp.</i>) Powdery mildew (<i>Blumeria graminis tritici</i>) Leaf and Glume blotches (<i>Septoria spp.</i>) Tan Spot (<i>P. tritici-repentis</i>)	5 fl. oz./A - greenup Followed by: 5 – 14 fl. oz./A - flagleaf
<p>APPLICATION INSTRUCTIONS AND RESTRICTIONS:</p> <ul style="list-style-type: none"> • Apply in early spring tillering to early joint (feekes stages 3-7) with a sequential application occurring at flagleaf (feekes stages 8-10) • The minimum application interval is 7 days. • Do not apply more than 0.228 lb ai of flutriafol/A per year. • Do not make more than 2 applications of Topguard fungicide per year. • Do not apply more than 28 fl. oz. of product/A per year. • Restricted entry interval (REI) is 12 hours. • Preharvest Intervals (PHI): Do not apply within 30 days of harvest when grown for grain, 15 days of harvest when grown for hay, zero (0) day PHI when grown for forage. 		
<p>This two-application program is intended for protection of both early vegetative stage wheat as well as grain-fill stage wheat.</p> <p>To achieve extended crop protection in susceptible varieties/hybrids, or under heavy disease infestations, and to provide protection through critical pollination and grain-fill periods, use full rates of Topguard fungicide (10-14 fl. oz./A), Topguard EQ fungicide (4-7 fl oz./A) or Lucento fungicide (3-5.5 fl. oz./A) applied at flagleaf.</p> <p>Topguard fungicide may be mixed with adjuvants, insecticides (such as Mustang® Maxx insecticide) or herbicides (such as Finesse® Cereal and Fallow herbicide) to provide insect or weed control and disease management in one application.</p>		
<p>ALL APPLICABLE DIRECTIONS, RESTRICTIONS, WORKER PROTECTION STANDARDS REQUIREMENTS AND PRECAUTIONS ON THE EPA REGISTERED TOPGUARD FUNGICIDE, EPA REG.NO. 279-3557, LABEL MUST BE FOLLOWED. THIS LABEL MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.</p>		

R-4414 031820