

**THIS RECOMMENDATION IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA**



**EPA Reg. No. 279-3557**

<b>2(ee) Recommendation</b>
FOR DISTRIBUTION AND USE ONLY IN WASHINGTON, OREGON, & IDAHO
THIS PRODUCT BULLETIN CONTAINS NEW OR ADDITIONAL DIRECTIONS FOR USE WHICH ARE RECOMMENDED BY FMC CORPORATION UNDER FIFRA SECTION 2(ee)
<b>IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH IT LABELING</b>

**DIRECTIONS FOR USE: HOPS, GRAPES AND STONE FRUIT**

When applied as directed through drip irrigation and micro sprinklers, Topguard Fungicide Specialty Crops will provide control of powdery mildew on the foliage of hops, grapes and stone fruit.

<b>Crop</b>	<b>Pest</b>	<b>Rate (fl. oz/A)</b>	<b>Application Directions</b>	<b>Retreatment Interval (Days)</b>
<b>Hops</b>	Powdery Mildew ( <i>Podosphaera macularis</i> , <i>Sphaerotheca macularis</i> )	14	Apply preventatively or when conditions are favorable for disease development. Repeat as necessary if conditions are favorable for disease development.	14
<b>Grapes</b> (fresh, table, wine, raisin)	Powdery Mildew ( <i>Uncinula necator</i> )	10	Apply preventatively or when conditions are favorable for disease development. Repeat as necessary if conditions are favorable for disease development.	14
<b>Stone Fruit</b> (See label for list of stone fruit)	Powdery Mildew ( <i>Podosphaera clandestina</i> )	14	Start applications at petal fall. Repeat as necessary if conditions are favorable for disease development.	7

**ALL APPLICABLE DIRECTIONS, RESTRICTIONS, AND PRECAUTIONS ON THE TOPGUARD FUNGICIDE SPECIALTY CROPS LABEL (EPA REG. NO. 279-3557) MUST BE FOLLOWED. THIS RECOMMENDATION SHOULD BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.**