

## SUPPLEMENTAL LABELING

FLUTRIAFOL	Group	3	Fungicide
------------	-------	---	-----------



EPA Reg. No. 279-3594

**For Use on Alfalfa and Rice as a Rotational Crop.**

**THIS SUPPLEMENTAL LABEL EXPIRES ON January 31, 2023 AND MUST NOT BE USED OR DISTRIBUTED AFTER THIS DATE**

### **DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Read the entire label. This label must be in the possession of the user at the time of application. Read the label affixed to the container for Topguard® Terra before applying. Use of Topguard® Terra according to this labeling is subject to the use precautions, restrictions, and limitations imposed by the label affixed to the Topguard® Terra container.

**Crop Rotation:** Fields treated with Topguard® Terra may be planted to the following crops and plant back intervals:

Crop	Plant back Interval
Alfalfa, barley, field corn; field corn grown for seed; popcorn; sweet corn; cotton; cucurbits; fruiting vegetables; grapes; hops; Brassica leafy vegetables crop group 5; leafy vegetables, except Brassica, crop group 4; peanut; pome fruit; grain sorghum; soybean; stone fruit; strawberry; sugar beet; tomato; tree nuts; triticale; or wheat	Immediately
Rice	30 days
All other crops	Prohibited

## SUPPLEMENTAL LABELING

### Use Directions for ALFALFA

Topguard® Terra works by forming a protective barrier around the alfalfa root at the point of disease infection. Precipitation in the form of rainfall or irrigation is required to move the product into the infection zone. The amount of water required will vary with soil type, soil moisture, and rainfall or irrigation intensity.

#### Application Method:

Apply Topguard® Terra as a broadcast spray application at a minimum spray volume of 10 gallons per acre or 5 gallons by air, or through sprinkler irrigations systems designed as described in the Chemigation section of this label. For continuous-move irrigation systems, apply the labeled rate in ½ acre-inch or more per acre. For stationary or non-continuous moving systems, inject Topguard® Terra spray mixture during the first 20-30 minutes of the set to insure fungicide reaches the soil surface. Applications should be made postemergence to alfalfa at the timing(s) outlined below.

Disease	Product Rate	Application Directions
Texas Root Rot / Cotton Root Rot (Phymatotrichopsis omnivora)	7.5 – 12 fl oz/A (0.24 – 0.39 lbs ai/A)	<p>Single Application: Apply in early spring prior to three (3) inches of new growth on the alfalfa OR after first cut of alfalfa when alfalfa is between two (2) and three (3) inches regrowth.</p> <p>Use the highest specified product rate when field is known to have high levels of cotton root rot disease pressure.</p>
	<p>First application: 7.5 – 12 fl oz/A (0.24 – 0.39 lbs ai/A)</p> <p>Second application: 6 – 7.5 fl oz/A (0.2 – 0.24 lbs ai/A)</p>	<p>Split Application: Apply first application in early spring prior to three (3) inches of new growth on the alfalfa</p> <p>Apply second application after second or third cutting of alfalfa when alfalfa is between two (2) and three (3) inches regrowth.</p> <p>Use split application under the following situations:</p> <ul style="list-style-type: none"> <li>• High intensity production practices</li> <li>• 2nd year of established plantings and older</li> </ul> <p>Use the highest specified product rate when field is known to have high levels of cotton root rot disease pressure.</p>

## SUPPLEMENTAL LABELING

### Restrictions:

- No single application may exceed 12 fl oz/A of product (0.39 lb ai)/A
- Make only one application per cutting
- Do not apply more than two applications per year.
- Do not exceed 19.5 fl oz (0.63 lb ai)/A per year
- Preharvest Interval (PHI): Do not apply within 14 days of harvest
- Restricted Entry Interval (REI) is 12 hours

Label Code: R-4372 032320 02-21-20