

SUPPLEMENTAL LABELING

TOLFENPYRAD	GROUP	21A	INSECTICIDE
	GROUP	39	FUNGICIDE



EPA Reg. No. 71711-31

Crop: Potato

States: States East of Mississippi River

This supplemental labeling expires 06/18/2021 and must not be used or distributed after this date.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This labeling and the EPA approved container label must be in the possession of the user at the time of application.

Read the label affixed to the container for Torac Insecticide before applying. Use of Torac Insecticide according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for Torac Insecticide.

New use directions appear on this supplemental labeling that may be different from those that appear on the container label

This supplemental labeling extends the use of Torac Insecticide to states located east of the Mississippi River.

APPLICATION RATE CHART

Potato		
Pest	Rate/Acre	Use Directions
Colorado potato beetle Leafhoppers	14.0 to 21.0 fl oz/acre (0.14 to 0.21 lb ai/acre)	USE RESTRICTIONS <ul style="list-style-type: none"> Apply by ground using a minimum of 20 gallons of water per acre. Apply by air using a minimum of 5 gallons of water per acre. See Chemigation statement in Application Directions. Do not apply more than 42.0 fluid ounces (0.42 lb ai) per acre per crop cycle.
Aphids Potato psyllid Thrips	17.0 to 21.0 fl oz/acre (0.17 to 0.21 lb ai/acre)	<ul style="list-style-type: none"> Do not make more than 2 applications per crop cycle. Allow at least 14 days between applications. Preharvest Interval (PHI): 14 days

©2018 Nichino America, Inc. TORAC and Nichino America logo are registered trademarks of Nichino America, Inc.

NICHINO
AMERICA[®]

Nichino America, Inc.
4550 Linden Hill Road, Suite 501
Wilmington, DE 19808
888-740-7700