

# SUPPLEMENTAL LABELING

TOLFENPYRAD	GROUP	<b>21A</b>	INSECTICIDE
	GROUP	<b>39</b>	FUNGICIDE



**EPA Reg. No. 71711-31**

**Cucurbit Vegetables Group (Crop Group 9)**

chayote (fruit); Chinese waxgourd (Chinese preserving melon); citron melon; cucumber; gherkin; gourd, edible ( includes hyotan, cucuzza, hechima, Chinese okra); *Momordica* spp (includes balsam apple, balsam pear, bitter melon, Chinese cucumber); muskmelon (hybrids and/or cultivars of *Cucumis melo*) (includes true cantaloupe, cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, and snake melon); pumpkin; squash, summer (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini); squash, winter (includes butternut squash, calabaza, hubbard squash, acorn squash, spaghetti squash); watermelon (includes hybrids and/or varieties of *Citrullus lanatus*)

This supplemental labeling expires 06/18/2021 and must not be used or distributed after this date.

**DIRECTIONS FOR USE**

**It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This labeling and the EPA approved container label must be in the possession of the user at the time of application. Read the label affixed to the container for TORAC insecticide before applying. Use of TORAC insecticide according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for TORAC insecticide.**

**New use directions appear on this supplemental labeling that may be different from those that appear on the container label.**

**APPLICATION RATE CHART FOR TORAC INSECTICIDE**

Pest	Rate/Acre	Use Directions
Leafhoppers  Aphids Flea beetles  Thrips Powdery mildew  Armyworm* Cucumber beetle* Melonworm* Pickleworm* Whitefly* Downy mildew*  *suppression only	14.0 to 21.0 fl oz/acre (0.14 to 0.21 lb ai/acre)  17.0 to 21.0 fl oz/acre (0.17 to 0.21 lb ai/acre)  21.0 fl oz/acre (0.21 lb ai/acre)	<p><b>USE RESTRICTIONS</b></p> <ul style="list-style-type: none"> <li>• Apply by ground <b>only</b>, using a minimum of 20 gallons of water per acre.</li> <li>• <b>Do not apply by air on cucurbits.</b></li> <li>• Do not apply more than 42.0 fluid ounces (0.42 lb ai) per acre per crop cycle.</li> <li>• Do not make more than 2 applications per crop cycle.</li> <li>• Do not make more than 4 applications per year.</li> <li>• Allow at least 14 days between applications.</li> <li>• Preharvest interval (PHI): 1 day</li> </ul>

©2018 Nichino America, Inc. TORAC and Nichino America logo are registered trademarks of Nichino America, Inc.

**NICHINO**  
**AMERICA**  
 Nichino America, Inc.  
 4550 Linden Hill Road, Suite 501  
 Wilmington, DE 19808  
 888-740-7700