SUPPLEMENTAL LABELING

TOLFENPYRAD GROUP 21A INSECTICIDE GROUP 39 FUNGICIDE



EPA Reg. No. 71711-31

For Use On: Brassica Leafy Greens (Crop Subgroup 4-16B)

arugula; broccoli, Chinese; broccoli raab; cabbage, abyssinian; cabbage, Chinese, bok choy; cabbage, seakale; collards; cress, garden; cress, upland; hanover salad; kale; maca, leaves; mizuna; mustard greens; radish, leaves; rape greens; realish, brocket, wild, shapped a purpositive greens; waterspress; sultivers, veriation, and bubridg of these commodities.

rocket, wild; shepherd's-purse; turnip greens; watercress; cultivars, varieties, and hybrids of these commodities

Use Pattern: Aerial Application

This supplemental labeling expires 07/15/2025 and must not be used or distributed after this date.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. This labeling and the EPA approved container label must be in the possession of the user at the time of application.

Read the label affixed to the container for TORAC[®] Insecticide before applying. Use of TORAC Insecticide according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for TORAC Insecticide.

New use directions appear on this supplemental labeling that may be different from those that appear on the container label.

APPLICATION RATE CHART FOR TORAC INSECTICIDE

Brassica Leafy Greens (Crop Subgroup 4-16B)		
Pest	Rate/Acre	Use Directions
Leafhoppers Planthoppers	14.0 to 21.0 fl oz/acre (0.14 to 0.21 lb ai/acre)	Aerial Application: Use a minimum of 5 gallons of water per acre.
Aphids Diamondback moth Flea beetle Garden symphylan Imported cabbageworm Cabbage maggot Thrips Alfalfa caterpillar* Bagrada bug* Armyworms*	17.0 to 21.0 fl oz/acre (0.17 to 0.21 lb ai/acre) 21.0 fl oz/acre (0.21 lb ai/acre)	 acre. Ground Application: Use a minimum of 20 gallons of water per acre. USE RESTRICTIONS Do not apply more than 42.0 fluid ounces (0.42 lb ai) per acre per crop cycle. Do not make more than 2 applications per crop cycle. Allow at least 14 days between applications. Do not make more than 4 applications per year. Preharvest Interval (PHI): 1 day
Cabbage webworm* Cloverworm, green* Corn earworm* Cross-striped cabbageworm* Cutworm species* Garden webworm* Saltmarsh caterpillar* Southern cabbageworm* Tobacco budworm* Tomato hornworm* Whitefly*		

©2022 Nichino America, Inc. Torac is a registered trademark of Nichino America, Inc. The Fujiichi-mark is a trademark of Nihon Nohyaku Co., Ltd.

