

Supplemental Labeling



Dow AgroSciences

Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

Transline[®]

EPA Reg. No. 62719-259

Rangeland and Permanent Grass Pastures

(For Use or Distribution in Arizona, New Mexico, Oklahoma, and Texas ONLY)

This supplemental label expires on December 31, 2015, and must not be used or distributed after this date.

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Transline[®] herbicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Transline according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Transline.

Directions for Use

Use Transline to control susceptible broadleaf weeds on rangeland areas, shelterbelts, Conservation Reserve Program acres, or established forage grasses in permanent grass pastures. Best results on most weeds are obtained when weeds are small and actively growing (see specific information below) and application is made in 10 gallons or more per acre of water using ground equipment.

There are no grazing or haying restrictions following Transline applications when used at labeled rates.

Application Rates

Apply Transline at a rate of 1/3 to 1 1/3 pint per acre when weeds are young and actively growing. Apply Transline as described below for control of spotted and diffuse knapweed, Canada thistle, musk thistle, yellow starthistle and suppression of Russian knapweed. Use the lower labeled application rate for young, actively growing weeds. Use the higher rate under less favorable growing conditions or on dense weed stands and/or larger weeds. Transline may also be tank mixed with 2,4-D at 1/2 to 1 lb ae per acre where weed species present are susceptible to 2,4-D.

Weed Species	Application Rate (pint/acre)	Application Timing
thistle, musk	1/3 - 1 ¹	Apply from rosette to early bolt growth stage.
thistle, artichoke thistle, Italian	1/3 - 2/3	Apply at the rosette growth stage.
starthistle, yellow	1/2 - 1	Apply from rosette to mid-bolt growth stage
knapweed, diffuse knapweed, spotted	2/3 - 1	Apply any time plants are actively growing, including fall regrowth. Optimum time is from mid bolt to late bud stage of growth.
thistle, artichoke thistle, Italian		Apply during the bolting growth stage.
thistle, Canada	2/3 - 1 1/3	Apply after the majority of basal leaves have emerged through the beginning of the bud stage. Application may also be made to fall regrowth
knapweed, Russian (suppression)	1 - 1 1/3	Apply from bud to mid-flower growth stage or treat fall regrowth.

¹Transline may be applied to musk thistle in the rosette stage at 1/3 pint per acre only when applied in tank mixture with 2,4-D at 1/2 to 1 lb ae per acre. Otherwise, apply Transline to musk thistle at 2/3 to 1 pint per acre.

®Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow
R113-014
Accepted: 03/25/15
Initial Printing