

---

## SUPPLEMENTAL LABELING

---

**Syngenta Crop Protection, LLC**  
P. O. Box 18300  
Greensboro, North Carolina 27419-8300

**SCP 1648B-S2 0223**

PYDIFLUMETOFEN	GROUP	<b>7</b>	FUNGICIDE
----------------	-------	----------	-----------

**Trebuset™**

Fungicide

ADEPIDYN® Technology\*

**This supplemental label expires on 08/23/2024 and must not be used or distributed after this date.**

Active Ingredient(s):

Pydiflumetofen** .....	41.7%
------------------------	-------

Other Ingredients:	58.3%
--------------------	-------

Total:	100.0%
--------	--------

\*Technology denotes the active ingredient Pydiflumetofen.

\*\*CAS No. 1228284-64-7

Trebuset is a flowable concentrate for seed treatment containing 4.17 lb pydiflumetofen per gallon.

**KEEP OUT OF REACH OF CHILDREN.**

## CAUTION

**EPA Reg. No. 100-1648**

All applicable directions, restrictions, and precautions on the EPA-registered label are to be followed. Before using Trebuset as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA-registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and/or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of federal law to use this product in a manner inconsistent with its labeling.

**syngenta.**

# Seed Treatment Directions

## Legume Vegetables (Succulent or Dried) – Crop Group 6

Crops			
Bean ( <i>Lupinus</i> spp.) Grain Lupin, Sweet Lupin, White Lupin, White Sweet Lupin Bean ( <i>Phaseolus</i> spp.) Field Bean, Kidney Bean, Lima Bean, Navy Bean, Pinto Bean, Runner Bean, Snap Bean, Tepary Bean, Wax Bean	Bean ( <i>Vigna</i> spp.) Adzuki Bean, Asparagus Bean, Blackeyed Pea, Catjang, Chinese Longbean, Cowpea, Crowder Pea, Moth Bean, Mung Bean, Rice Bean, Southern Pea, Urd Bean, Yardlong Bean	Pea ( <i>Pisum</i> spp.) Dwarf Pea, Edible-pod Pea, English Pea, Field Pea, Garden Pea, Green Pea, Snow Pea, Sugar Snap Pea	Broad Bean (fava bean) ( <i>Vicia faba</i> ) Chickpea (garbanzo bean) ( <i>Cicer arietinum</i> ) Guar ( <i>Cyamopsis tetragonoloba</i> ) Jackbean ( <i>Canavalia ensiformis</i> ) Lablab Bean (hyacinth bean) ( <i>Lablab purpureus</i> ) Lentil ( <i>Lens esculenta</i> ) Pigeon Pea ( <i>Cajanus cajan</i> ) Soybean, (immature seed) (edamame) ( <i>Glycine max</i> ) Sword Bean ( <i>Canavalia gladiata</i> )
Target Disease/Pest	Use Rate		
	fl oz/100 lb seed	mg ai/1,000 seeds*	
Ascochyta Blight caused by seed-borne <i>Ascochyta</i> spp.  Black root rot caused by <i>Fusarium solani</i>  Fusarium root rot caused by <i>Fusarium</i> spp.	0.31-0.77	Beans <sup>1</sup> : 36-91  Peas <sup>2</sup> : 24-59  Chickpea <sup>3</sup> : 51-126  Cowpea <sup>4</sup> : 14-35  Lentil <sup>5</sup> : 4-11	
*Based on an average seed weight of: 1. 1,250 beans/lb 2. 1,930 peas/lb 3. 900 chickpeas/lb 4. 3,200 cowpeas/lb 5. 10,600 lentils/lb			
<b>Resistance Management:</b> <ul style="list-style-type: none"> <li>Refer to Section 3.1.</li> </ul>			
USE RESTRICTIONS			
<ul style="list-style-type: none"> <li><b>DO NOT</b> feed or harvest cowpea forage and hay.</li> <li><b>DO NOT</b> exceed 0.357 lb ai/A/year of pydiflumetofen-containing products through any combination of seed and foliar applications.</li> <li><b>Sections 4.0, 4.1, 5.0, and 6.0</b> contain additional product use restrictions.</li> </ul>			