

**PRODUCT BULLETIN**  
**2(ee) Recommendation**

**THIS PRODUCT BULLETIN CONTAINS NEW OR ADDITIONAL DIRECTIONS  
FOR USE WHICH ARE RECOMMENDED BY INNICTIS CROP CARE LLC  
UNDER FIFRA SECTION 2(ee)**



**EPA Reg. No. 89167-103-89391**

**FIFRA Section 2(ee) Recommendation:** This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or accepted by the U.S. Environmental Protection Agency. It is a violation of federal law to use this product in a manner inconsistent with its label. This product bulletin must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATES OF FLORIDA, LOUISIANA & TEXAS**

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product inconsistent with its labeling.

**Addition of sweet orange scab (*Elsinoe australis*) as a disease controlled in citrus**

<b>CROP</b>	<b>TARGET DISEASE</b>	<b>USE RATE fl oz Product/A</b>	<b>Applications Instructions</b>
Citrus  Grapefruit Lemon Lime Orange (sour and sweet) Tangerine  Including all cultivars and/or hybrids of these	Sweet Orange Scab ( <i>Elsinoe australis</i> )	10 - 15.4	TREVO DCZ applications should begin prior to disease development and continue throughout the season on 7-21 day intervals following the resistance management guidelines. Applications may be made by ground or air. An adjuvant may be added at recommended rates. A horticultural spray oil should be used to improve control of greasy spot.  If disease pressure is high, use the shortest interval and highest rate.  Make no more than 2 sequential applications before alternating to another fungicide with a non-QoI (Group 11) different mode of action. <b>DO NOT</b> make more than 4 applications of TREVO DCZ or other Group 11 fungicides per season.

**Application:** For best results, sufficient water volume must be used to provide thorough coverage. TREVO DCZ can be applied by either ground or aerial application. A minimum of 15 gal/A for ground applications is recommended. For aerial applications a minimum of 10 gal/A of water is recommended.

**Complete list of citrus fruits:** Calamondin; citron; citrus hybrids (includes chironja, tangelo, tangor); grapefruit; kumquat; lemon; lime; mandarin (tangerine); orange, sour; orange, sweet; pummelo; Satsuma mandarin

**Specific Use Restrictions:**

- **DO NOT** use TREVO DCZ in citrus plant propagation nurseries.
- **DO NOT** apply more than 61.5 fl oz/A/year of TREVO DCZ.
- Single Maximum Application Rate of TREVO DCZ: 15.4 fl oz/A.
- **DO NOT** apply more than 0.5 lb ai/A/year of difenoconazole-containing products.
- **DO NOT** apply more than 1.5 lb ai/A/year of azoxystrobin-containing products.
- May be applied the day of harvest (0-day PHI).
- Re-treatment Interval: 7 days

**Always refer to the complete product label for full use directions and precautionary language.**

**Distributed By:**

INNICTIS CROP CARE, LLC  
1880 Fall River Drive, Suite 100  
Loveland, CO 80538

031023