

PRODUCT BULLETIN
2(ee) Recommendation

**THIS PRODUCT BULLETIN CONTAINS NEW OR ADDITIONAL
DIRECTIONS FOR USE
WHICH ARE RECOMMENDED BY INNVICTIS CROP CARE LLC
UNDER FIFRA SECTION 2(ee)**



EPA Reg. No. 89168-100-89391

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or accepted by the U.S. Environmental Protection Agency. It is a violation of federal law to use this product in a manner inconsistent with its label. This product bulletin must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

DIRECTIONS FOR USE - It is a violation of Federal law to use this product inconsistent with its labeling.

Field Corn, Popcorn, Corn Grown For Seed Production			
Disease	Dosage Rate		Use Directions
	Fl. Oz./A	GPA	
Tar Spot (<i>Phyllachora</i> <i>Maydis</i>)	13 to 22.7	Ground minimum: 10 Aerial minimum: 2	Apply prior to disease onset when conditions favor disease development. A second application may be made no fewer than 7 days later but not to exceed 22.7 fl oz per acre per year rate. Use <i>TREVO TRZ DRV</i> Fungicide as part of an integrated pest management program (IPM). Apply as a foliar spray or via chemigation in sufficient water to obtain thorough coverage of plants.

RESTRICTIONS

- **DO NOT** make more than two (2) applications per year when applied at reduced rates.
- **DO NOT** make more than 2 sequential applications before alternating to another fungicide with a different mode of action.
- **DO NOT** apply more than 22.7 fl oz (0.11 lb azoxystrobin & 0.09 lb tetraconazole) per acre per application.
- **DO NOT** apply more than 22.7 fl oz (0.11 lb azoxystrobin & 0.09 lb tetraconazole) of *TREVO TRZ DRV* per acre per year.
- **DO NOT** apply more than 0.09 lb ai of a tetraconazole-containing product per acre per year.
- **DO NOT** apply more than 2.0 lb ai of an azoxystrobin-containing product per acre per year.
- **DO NOT** apply *TREVO TRZ DRV* after corn growth stage R3 (brown silk/milk).
- **DO NOT** apply within 7 days of harvest (7 day PHI).
- **DO NOT** harvest silage within 21 days of an application.
- Minimum Retreatment Interval = 7 days

Use Rate: 13 to 22.7 oz per acre

Always refer to the complete product label for full use directions and precautionary language.

Distributed By:

INNVICTIS CROP CARE, LLC
1880 Fall River Drive, Suite 100
Loveland, CO 80538

120121R1