

---

## SUPPLEMENTAL LABEL

---



**Syngenta Crop Protection, Inc.**  
P. O. Box 18300  
Greensboro, North Carolina 27419-8300  
[www.syngenta-us.com](http://www.syngenta-us.com)

**Trimmit® 2SC**

**Plant Growth Regulator**

**For Turfgrass**

EPA Reg. No. 100-1014

Supplemental for *Poa annua* Suppression

*Active Ingredient:*

Paclobutrazol

(±)-(R\*,R\*)-beta-[(4-chlorophenyl)methyl]-alpha-

(1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethanol ..... 22.3%

*Other Ingredients:* ..... 77.7%

Total: ..... 100.0%

Contains 2 lbs. active ingredient per gallon.

**KEEP OUT OF REACH OF CHILDREN.**

**CAUTION**

**All applicable directions, restrictions and precautions on the EPA-registered label are to be followed.**

**Before using Trimmit 2SC as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling supersedes the Directions for Use on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.**

SCP 1014B-S1 1006

## DIRECTIONS FOR USE

### When to Use for *Poa annua* Suppression

Apply when the *Poa annua* is actively growing. In moderate climates the application timing may be fall through late spring. In northern climates, spring and fall timings are recommended. In climates with a prolonged winter dormancy, fall applications can be made up to one month prior to anticipated first killing frost. Repeat application should be made as part of a *Poa annua* suppression program.

### When to Use for Color and Quality Enhancement of Winter Overseeded Turf and *Poa annua* Suppression

Apply anytime after overseeded turf has successfully established itself (cool-season turf species should be near 85% of turf composition). Do not apply after March 15<sup>th</sup> to putting greens to avoid delay in bermudagrass green-up. Moderate soil moisture conditions should be present before and after application to achieve best regulating effect.

#### Specific Rates

Warm Season Grasses	Cool Season Grasses <sup>1</sup> (except putting greens)	Bentgrass Putting Greens & Overseeded Bermuda	Overseeded Bermudagrass
0.50 lbs. a.i./A (sandy soils)	0.25 to 0.50 lbs. a.i./A	0.10 to 0.25 lbs. a.i./A	0.10 to 0.16 lbs. a.i./A
0.75 lbs. a.i./A (clay soils)			

<sup>1</sup>Reduce application rate by 50% if *Poa annua* is a significant portion of turfgrass population (greater than 50%).

### Program Scheduling

#### For *Poa annua* Suppression

Trimmit 2SC will suppress the growth and competitive ability of *Poa annua*. Apply when *Poa annua* is actively growing at 16 to 32 ounce per acre. Lower rates and more frequent applications can be made if *Poa annua* discoloration cannot be tolerated. However, only moderate *Poa annua* suppression may be achieved. Do not apply more than 4 quarts per acre per year (2 lbs. a.i./A)

#### For Color, Quality Enhancement, and *Poa annua* Suppression in Overseedings

On bentgrass greens and overseeded bermudagrass, repeat applications can be made as long as the turf is actively growing. Do not apply more than 4 quarts per acre per year (2 lbs. a.i./A). Apply to overseeded turf when seed has established and turf composition is near 85% of the cool season species.

---

Trimmit® and the Syngenta logo are trademarks of a Syngenta Group Company

©2007 Syngenta

Syngenta Crop Protection, Inc.  
P.O. Box 18300  
Greensboro, North Carolina 27419-8300  
[www.syngenta-us.com](http://www.syngenta-us.com)

SCP 1014B-S1 1006