
SUPPLEMENTAL LABELING

Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, North Carolina 27419-8300
SCP 1613A-S1 0718

PROPICONAZOLE	GROUP	3	FUNGICIDE
BENZOVINDIFLUPYR	GROUP	7	FUNGICIDE
AZOXYSTROBIN	GROUP	11	FUNGICIDE

Trivapro® Fungicide

This supplemental label expires on 4/25/2021 and must not be used or distributed after this date.

Active Ingredients:

Benzovindiflupyr*.....	2.9%
Azoxystrobin**.....	10.5%
Propiconazole***.....	11.9%
<hr/>	
Other Ingredients:	74.7%
Total:	100.0%

*CAS No. 1072957-71-1

**CAS No. 131860-33-8

***CAS No. 60207-90-1

Trivapro Fungicide is formulated as a suspo-emulsion and contains 0.25 lb of benzovindiflupyr, 0.92 lb of azoxystrobin, and 1.04 lb of propiconazole active ingredients per gallon.

KEEP OUT OF REACH OF CHILDREN.

WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See additional precautionary statements and directions for use inside booklet.
See First Aid statement inside booklet and on container label.

EPA Reg. 100-1613

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed. Before using Trivapro Fungicide as permitted according to this Supplemental Labeling, read and follow all applicable directions, restrictions, and precautions on the EPA-registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and/or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

DIRECTIONS FOR USE

Bulb Vegetable Group 3-07; Dry Bulb Group 3-07A; Onions, green Group 3-07B

Crops (including all cultivars, varieties, and/or hybrids of these)			
Dry Bulb Group 3-07A: Daylily, bulb Fritillaria, bulb Garlic, bulb Garlic, great-headed Garlic, serpent, bulb Lily, bulb Onion, bulb Onion, Chinese, bulb Onion, pearl	Potato, bulb Shallot, bulb Green Onion Group 3-07B: Chive, Chinese fresh leaves Chive, fresh leaves Elegans hosta Fritillaria, leaves Kurrat Lady's leek Leek	Leek, wild Onion, Beltsville bunching Onion, fresh Onion, green Onion, marcostem Onion, tree, tops Onion, Welsh, tops Shallot, fresh leaves	
Target Disease	Rate (fl oz/A)	Application Timing	Use Directions
Cladosporium leaf blotch <i>(C. allii)</i> Powdery Mildew <i>(Leveillula taurica)</i> Purple Blotch <i>(Alternaria porri)</i> Rust <i>(Puccinia allii)</i> Stemphyllium leaf blight and stalk rot <i>(S. vesicarium)</i>	16.0 – 27.7	Apply prior to disease development. Continue applications through season on a 7 to 14-day interval, following the resistance management guidelines. No more than two applications of Trivapro Fungicide may be applied on a 7-day interval.	Apply by ground, air, or chemigation. An adjuvant may be added at recommended rates. If disease pressure is high, use the shortest interval and highest rate.
Resistance Management:			
<ul style="list-style-type: none"> • Do not make more than two applications of Trivapro Fungicide or other Group 7 or 11 fungicides before alternation with a fungicide that is not in Group 7 or 11. 			
USE RESTRICTIONS			
<ol style="list-style-type: none"> 1) Refer to Section 6.1 for additional product use restrictions. 2) Maximum Single Application Rate: 27.7 fl oz/A 3) Minimum Application Interval: 7 days 4) Maximum Annual Rate: 55.4 fl oz product/A/year <ol style="list-style-type: none"> a. Do not apply more than 0.45 lb ai/A/year of propiconazole-containing products. b. Do not apply more than 1.5 lb ai/A/year of azoxystrobin-containing products. c. Do not apply more than 0.272 lb ai/A/year of benzovindiflupyr-containing products. 5) Do not exceed 4 applications per year. 6) Pre-harvest Interval (PHI): <ol style="list-style-type: none"> a. Green Onion Group 3-07B: 7 days b. Dry Bulb Group 3-07A: 14 days 7) Aerial application is prohibited in New York State. 			

Grasses Grown for Seed

Crops (including all cultivars and/or varieties of these)			
Bluegrass Bromegrass Fescue		Orchardgrass Ryegrass	
Target Disease	Rate (fl oz/A)	Application Timing	Use Directions
Ergot Stem Diseases Powdery Mildew (<i>Erysiphe graminis</i>) Rusts (<i>Puccinia</i> spp.) Selenophoma Stem Eyespot (<i>Selenophoma</i> spp.)	16.0 – 27.4	Applications should begin prior to disease development and continue throughout the season on a 14-28 day schedule.	Apply by ground, air, or chemigation. The addition of a spreading/penetrating type adjuvant such as organo-silicon blends with either non-ionic surfactants (NIS) or vegetable based crop oils (COC); or vegetable based COC (not mineral); or NIS with at least 90% concentration is recommended. If disease pressure is high, use the shortest interval and highest rate.
Resistance Management:			
<ul style="list-style-type: none"> Do not make more than two applications of Trivapro Fungicide or other Group 7 or 11 fungicides before alternation with a fungicide that is not in Group 7 or 11. 			
USE RESTRICTIONS			
<ol style="list-style-type: none"> Refer to Section 6.1 for additional product use restrictions. Maximum Single Application Rate: 27.4 fl oz/A Minimum Application Interval: 14 days Maximum Annual Rate: 54.8 fl oz/A/year <ol style="list-style-type: none"> Do not apply more than 0.9 lb ai/A/year of propiconazole-containing products. Do not apply more than 0.8 lb ai/A/year of azoxystrobin-containing products. Do not apply more than 0.11 lb ai/A/year of benzovindiflupyr-containing products. Do not exceed 2 applications per year. Use limited to Idaho, Minnesota, Nebraska, Oregon, and Washington. Only for use on the following cool season grasses: bluegrass, bromegrass, fescue, orchardgrass, and ryegrass. Aerial application is prohibited in New York State. Do not apply to Bermuda grass grown for seed. Do not feed treated straw, seed, or screenings to livestock. Do not feed hay cut within 20 days of last application. Do not graze treated areas within 140 days of the last application. Pre-harvest Interval (PHI): 20 days 			

Sugarcane

Crops (including all cultivars, varieties, and/or hybrids of these)			
Sugarcane			
Target Disease	Rate (fl oz/A)	Application Timing	Use Directions
Brown Rust (<i>Puccinia melanocephala</i>)	16.0 - 20.7	Apply prior to disease development.	Apply by ground, air, or chemigation.
Orange Rust (<i>Puccinia kuehnii</i>)		Continue applications through season on a 14 to 28-day interval, following the resistance management guidelines.	An adjuvant may be added at recommended rates. If disease pressure is high, use the shortest interval.
Resistance Management:			
<ul style="list-style-type: none"> Do not make more than two applications of Trivapro Fungicide or other Group 7 or 11 fungicides before alternation with a fungicide that is not in Group 7 or 11. 			
USE RESTRICTIONS			
<ol style="list-style-type: none"> Refer to Section 6.1 for additional product use restrictions. Maximum Single Application Rate: 20.7 fl oz/A Minimum Application Interval: 14 days Maximum Annual Rate: 62.1 fl oz/A/year <ol style="list-style-type: none"> Do not apply more than 0.675 lb ai/A/year of propiconazole-containing products. Do not apply more than 0.80 lb ai/A/year of azoxystrobin-containing products. Do not apply more than 0.204 lb ai/A/year of benzovindiflupyr-containing products. Do not exceed 3 applications per year. Pre-harvest Interval (PHI): 30 days Aerial application is prohibited in New York State. 			

Trivapro® and the Syngenta logo are trademarks of a Syngenta Group Company

©2018 Syngenta