



S-METOLACHLOR GROUP 15 HERBICIDE

HERBICIDE

FOR WEED CONTROL IN BEANS, PEAS, AND LENTILS; CORN; COTTON; GRASSES GROWN FOR SEED; HORSERADISH; PEANUTS; POTATOES; PUMPKIN; RHUBARB; SAFFLOWERS; SWEET, GRAIN, OR FORAGE SORGHUM; SOYBEANS; SOYBEANS, IMMATURE SEED; SUGAR BEETS; SUNFLOWERS; AND TOMATOES

ACTIVE INGREDIENT:

S-metolachlor ..... % BY WT. 62.0%

OTHER INGREDIENTS: ..... 38.0%

TOTAL: ..... 100.0%

This product is formulated as an emulsifiable concentrate (EC) and contains 5.69 pounds of active ingredient per gallon

KEEP OUT OF REACH OF CHILDREN  
CAUTION / PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail).

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300

SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS.

Not for Sale, Sale into, Distribution and/or Use in Nassau and Suffolk Counties of New York State.

EPA Reg. No.: 89168-99-89391



HERBICIDE



Distributed By:  
INNICTIS® CROP CARE, LLC  
1880 Fall River Drive, Suite 100  
Loveland, CO 80538

110320RD062121

FIRST AID	
<b>IF IN EYES:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF ON SKIN OR CLOTHING:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF SWALLOWED:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• <b>DO NOT</b> induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• <b>DO NOT</b> give anything by mouth to an unconscious person.</li> </ul>
<b>IF INHALED:</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give the person artificial respiration, preferably mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>HOTLINE NUMBER</b> Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency information concerning this product, call the National Pesticides Information Center (NPIC) at <b>1-800-858-7378</b> or your poison control center at <b>1-800-222-1222</b> . For Chemical Spill, Leak, Fire or Exposure, call CHEMTREC <b>800-424-9300</b> .	

**PRECAUTIONARY STATEMENTS**  
**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**  
**CAUTION**

Harmful if swallowed. Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

- PERSONAL PROTECTIVE EQUIPMENT (PPE)**  
**Applicators and other handlers must wear:**
- Long-sleeved shirt and long pants
  - Chemical-resistant gloves made of barrier laminate or Viton® ≥14 mils
  - Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

**Engineering Control Statements**

Mixers and loaders supporting aerial applications are required to use closed systems. The closed system must be used in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.607(d-e)]. When using the closed system, the mixers' and loaders' PPE requirements may be reduced or modified as specified in the WPS.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.607(d-e)], the handler PPE requirements may be reduced or modified as specified in the WPS.

**USER SAFETY RECOMMENDATIONS**

**Users should:**

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**ENVIRONMENTAL HAZARDS**

**DO NOT** apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment wash water or rinsate.

**Groundwater Advisory**

S-metolachlor is known to leach through soil into ground water under certain conditions as a result of label use. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.

**Surface Water Advisory**

S-metolachlor has the potential to contaminate surface water through ground spray drift. Under some conditions, it may also have a high potential for runoff into surface water (primarily via dissolution in runoff water) for several months post-application. These include poorly drained or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlaying extremely shallow groundwater, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas overlaying tile drainage systems that drain to surface water.

**Mixing/Loading Instructions**

Care must be taken when using this product to prevent back-siphoning into wells, spills, or improper disposal of excess pesticide, spray mixtures, or rinsates. Check-valves or anti-siphoning devices must be used on all mixing and/or irrigation equipment.

This product may not be mixed or loaded within 50 feet of perennial or intermittent streams and rivers, natural or impounded lakes, and reservoirs. This product may not be mixed/loaded or used within 50 feet of all wells, including abandoned wells, drainage wells, and sink holes. Operations that involve mixing, loading, rinsing, or washing of this product into or from pesticide handling or application equipment or containers within 50 feet of any well are prohibited, unless conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be positioned on or moved across the pad. Such a pad shall be designed and maintained to contain any product spills or equipment leaks, container or equipment rinse or wash water, and rain water that may fall on the pad. Surface water shall not be allowed to either flow over or from the pad, which means the pad must be self-contained. The pad shall be sloped to facilitate material removal. An unroofed pad shall be of sufficient capacity to contain at a minimum 110% of the capacity of the largest pesticide container or application equipment on the pad. A pad that is covered by a roof of sufficient size to completely exclude precipitation from contact with the pad shall have a minimum containment capacity of 100% of the capacity of the largest pesticide container or application equipment on the pad. Containment capacities as described above shall be maintained at all times. The above-specified minimum containment capacities **DO NOT** apply to vehicles when delivering pesticide shipments to the mixing/loading site.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

This product must be used only in accordance with instructions on this label.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.**

**Exception:** If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: coveralls, chemical-resistant gloves made of barrier laminate or Viton  $\geq$  14 mils and shoes plus socks.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR WEED CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

To avoid spray drift, **DO NOT** apply under windy conditions. Avoid spray overlap, as crop injury may result.

**Not for Sale, Sale into, Distribution and/or Use in Nassau and Suffolk Counties of New York State**

### PRODUCT INFORMATION

*VISOR II XRT* is a selective herbicide for use as a preplant surface-applied, preplant incorporated, or preemergence treatment in water or fluid fertilizer for control of most annual grasses and certain broadleaf weeds in beans, peas, and lentils; corn (all types); cotton; grasses grown for seed; peanuts; potatoes; sowfollers; sweet, grain, or forage sorghum; soybeans; soybean, immature seed; sugar beets; sunflowers; and tomatoes.

Dry weather following preemergence application of *VISOR II XRT* or a tank mixture may reduce effectiveness. Cultivate if weeds develop. Where reference is made to weeds partially controlled, partial control can either mean erratic control from good to poor, or consistent control at a level below that generally considered acceptable for commercial weed control.

### Tank Mixtures

Observe all precautions and limitations on the labels of each product used in tank mixtures. Tank mixtures are permitted only in those states where the tank mix partner is registered. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

### Precautions

- Injury may occur following the use of this product under abnormally high soil moisture conditions during early development of the crop.

### Restrictions

- **DO NOT** use in nurseries, turf, or landscape plantings.
- **DO NOT** apply under conditions which favor runoff or wind erosion of soil containing this product to nontarget areas.
- If this product is incorporated, any supplemental tillage before planting must not exceed the depth of incorporation.
- Certain states may have established rate limitations for atrazine within specific geographical areas. Consult your state lead pesticide control agency for additional information when tank mixing this product with atrazine-containing products. It is a violation of this label to deviate from state use regulations.

### To prevent off-site movement due to runoff or wind erosion:

1. Avoid treating powdery dry or light sand soils when conditions are favorable for wind erosion. Under these conditions, the soil surface should first be settled by rainfall or irrigation.
2. **DO NOT** apply to impervious substrates, such as paved or highly compacted surfaces.

3. **DO NOT** use tailwater from the first flood or furrow irrigation of treated fields to treat nontarget crops, unless at least 1/2 inch of rainfall has occurred between application and the first irrigation.

### RESISTANCE-MANAGEMENT RECOMMENDATIONS

S-metolachlor, the active ingredient in this product, is a Group 15 herbicide. Any weed population may contain or develop plants naturally resistant to this product and other Group 15 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance-management strategies should be followed.

### Weed Management

To delay herbicide resistance, take one or more of the following steps:

- Rotate the use of this product or other Group 15 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in the field.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation, and that considers tillage (or other mechanical control methods), cultural (e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Scout before and after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method such as hoeing or tillage. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or certified crop advisors for additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact INNVICTIS CROP CARE, LLC at 855-466-8428.

### Management of Resistant Biotypes

Since the occurrence of resistant weeds cannot be determined until after product use and scientific confirmation, manufacturer is not responsible for any losses that may result from the failure of this product to control resistant weed biotypes.

The following good agronomic practices are recommended to reduce the spread of resistant biotypes:

- If a naturally occurring resistant biotype is present in your application site, this product should be tank mixed or applied sequentially with an appropriately labeled herbicide with a different mode of action to achieve control.
- Cultural and mechanical control practices (e.g. crop rotation or tillage) may also be used as appropriate.
- Scout treated application site after herbicide applications and control escaping weeds including resistant biotypes before they get set.
- Thoroughly clean equipment before leaving fields known to contain resistant biotypes.
- Contact your local sales representative, crop advisor, or extension agent to find out if suspected resistant weeds to this Mode of Action have been found in your region. If resistant biotypes of target weeds have been reported, use the application rates of this product specified for your local conditions. Tank mix products so that there are multiple effective mechanisms of actions for each target weed.

## Integrated Pest (Weed) Management

This product may be integrated into an overall weed pest management strategy whenever the use of an herbicide is required. Practices known to reduce weed development (tillage, crop competition) and herbicide use (weed scouting, proper application timing, banding) should be followed wherever possible. Consult local agricultural and weed authorities for additional IPM strategies established for your area.

## SOIL TEXTURES AND HERBICIDE RATES

Where rates are based on coarse-, medium-, or fine-textured soils, it is understood that soil textural classes are generally categorized as follows:

COURSE	MEDIUM	FINE
Sand	Loam	Sandy Clay Loam
Loamy Sand	Silt Loam	Silty Clay Loam
Sandy Loam	Silt	Clay Loam
		Sandy Clay
		Silty Clay
		Clay

Within rate ranges in the rate tables and elsewhere on this label, use the lower rate on soils relatively coarse-textured or low in organic matter; use the higher rate on soils relatively fine-textured or high in organic matter.

This product may be applied preemergence alone, or in combination with tank mix partners specified on this label, following preplant incorporated herbicides when used according to their label requirements, provided that such use is not prohibited on the respective labels.

Thoroughly clean sprayer or other application device before using. Dispose of cleaning solution in a responsible manner. **DO NOT** use a sprayer or applicator contaminated with any other materials, or crop damage or clogging of the application device may result.

## VISOR II XRT APPLIED ALONE

### Weeds Controlled

**VISOR II XRT** is taken up by the shoots and/or roots of emerging weeds. This uptake results in the inhibition of shoot and root tissue growth soon after weed germination. Because of this, this product will not control emerged weeds and should be applied prior to weed emergence.

If **VISOR II XRT** is incorporated, **DO NOT** exceed a 2- to 3-inch depth. Any tillage after the incorporation of this product and before planting should not exceed 2 to 3 inches.

Dry weather following application of **VISOR II XRT** may reduce weed control. Cultivate if weeds develop.

Where reference is made to weeds partially controlled, partial control can either mean erratic control from good to poor or consistent control at a level below that generally considered acceptable for commercial weed control. Control of these weeds can be erratic, due partially to variable weather conditions. The following procedures may improve the control of weeds listed as partially controlled in **Table 1**:

1. Thoroughly till soil to destroy germinating and emerged weeds.
2. Plant crop into moist soil immediately after tillage. If this product is to be used preemergence, apply at planting or immediately after planting.
3. If available, sprinkle irrigate within 2 days after application. Apply 1/2 to 1 inch of water. Use lower water volume (1/2 inch) on coarse-textured soils and higher volume (1 inch) on fine-textured soils. Also, refer to the section on **Center Pivot Irrigation** for application of this method of applying this product.
4. If irrigation is not possible and rain does not occur within 2 days after planting and application, weed control may be decreased. Under these conditions, a uniform, shallow cultivation is recommended as soon as weeds emerge.

**Table 1: Weeds Controlled or Partially Controlled by VISOR II XRT Applied Prior to Weed Emergence**

Common Name	Scientific Name	Control (C) Partial Control(PC)
<b>GRASSES</b>		
Barnyardgrass	<i>Echinochloa crus-galli</i>	C
Crabgrass, large	<i>Digitaria ischaemum</i>	C

Common Name	Scientific Name	Control (C) Partial Control(PC)
<b>GRASSES</b>		
Crabgrass, smooth	<i>Digitaria sanguinalis</i>	C
Crowfootgrass	<i>Dactyloctenium aegyptium</i>	C
Cupgrass, Prairie	<i>Eriochloa contracta</i>	C
Cupgrass, Southwestern	<i>Eriochloa acuminata</i>	C
Cupgrass, woolly	<i>Eriochloa villosa</i>	PC <sup>1</sup>
Foxtail, bristly	<i>Setaria verticillata</i>	C
Foxtail, giant	<i>Setaria faberi</i>	C
Foxtail, green	<i>Setaria viridis</i>	C
Foxtail, millet	<i>Setaria italica</i>	C
Foxtail, yellow	<i>Setaria pumila</i>	C
Goosegrass	<i>Eleusine indica</i>	C
Johnsongrass (seedling)	<i>Sorghum halepense</i>	PC
Millet, wild-proso	<i>Panicum millaceum</i>	PC <sup>1</sup>
Panicum, fall	<i>Panicum dichotomiflorum</i>	C
Panicum, Texas	<i>Panicum texanum</i>	PC
Rice, red	<i>Oryza punctata</i>	C
Sandbur, field	<i>Cenchrus spinifex</i>	PC
Ryegrass, Italian	<i>Lolium multiflorum</i>	C
Sandbur, Southern	<i>Cenchrus spinifex</i>	PC
Shattercane	<i>Sorghum bicolor</i>	PC
Signalgrass, broadleaf	<i>Urochloa platyphylla</i>	C
Sorghum (Volunteer)	<i>Sorghum bicolor</i>	PC
Witchgrass	<i>Panicum capillare</i>	C
<b>BROADLEAF WEEDS</b>		
Amaranth, Palmer	<i>Amaranthus palmeri</i>	C
Amaranth, Powell	<i>Amaranthus powellii</i>	C
Beggarweed, Florida	<i>Desmodium illinoense</i>	PC
Carpetweed	<i>Mollugo verticillata</i>	C
Eclipta	<i>Eclipta prostrata</i>	PC
Galinsooga, hairy	<i>Galinsooga quadriradiata</i>	C
Galinsooga, smallflower	<i>Galinsooga parviflora</i>	C
Nightsahde, Eastern black	<i>Solanum ptychanthum</i>	C
Nightshade, hairy	<i>Solanum physalifolium</i>	PC
Pigweed, prostrate	<i>Amaranthus blitoides</i>	C
Pigweed, redroot	<i>Amaranthus retroflexus</i>	C
Pigweed, smooth	<i>Amaranthus hybridus</i>	C
Pigweed, tumble	<i>Amaranthus albus</i>	C
Purslane, common	<i>Portulaca oleracea</i>	PC
Pusley, Florida	<i>Richardia scabra</i>	C
Spiderwort, tropical	<i>Commelina benghalensis</i>	C
Waterhemp, common	<i>Amaranthus rudis</i>	C
Waterhemp, tall	<i>Amaranthus tuberculatus</i>	C

(cont'd next page)

Common Name	Scientific Name	Control (C) Partial Control(PC)
<b>SEDGES</b>		
Nutsedge, yellow	<i>Cyperus esculentus</i>	C
1 Refer to the corn section of this label for additional directions.		

## REPLANT AND ROTATIONAL CROPS SECTION

### Replanted Crop Directions

This section covers replant crops that may be planted following a lost crop that has had an application of *VISOR II XRT*.

If a crop treated with *VISOR II XRT* is lost, any crop on this label, or on an EPA-approved supplemental label, may be replanted immediately provided that the rate of this product applied to the previous crop was not greater than the labeled rate for the crop to be replanted. If the first application was banded and the replant crop is planted in the center of the untreated bands, a second banded treatment may be applied at the rate for the use pattern for the replant crop, provided the application does not overlap the first application band.

### Rotational Crop Directions

**DO NOT rotate to food or feed crops other than those listed below.**

Crop	Crop Rotational Interval (Months)
Alfalfa	4
Barley, oats, rye and wheat	4.5
Clover (seeded)	9
Tobacco, buckwheat, and rice	Spring following the application
All crops not listed	12

### Restrictions

- To avoid injury to rotational alfalfa or clover:
  - DO NOT** apply more than 2.7 pints (1.9 lb ai) per acre of this product per acre in the previous crop.
  - DO NOT** make lay-by or other postemergent applications of this product in the previous crop.

Below in the rotational crop subsections A through C is a listing of rotational crop options that are made possible through S-metolachlor tolerances which were established by the EPA as crop groupings.

### Precautions

- Rotating to crops within these crop groupings at less than 60 days may result in crop injury.
- If the rate of this product applied in the previous crops was greater than the rate listed here (Sections A-C below), these crops cannot be planted until the following spring.

- A. If not more than 1.75 pints (1.25 lb ai) per acre of *VISOR II XRT* was applied to the field, the following crops (as well as those listed under subsections B or C below) may be planted 60 days after the last application. A second application of an S-metolachlor-containing product to the following crops is prohibited within 60 days of the original application.**

**Crop Subgroup 1B - Root Vegetables:** garden beet, edible burdock, carrot, celeriac, turnip-rooted chervil, chicory, ginseng, horseradish, turnip-rooted parsley, parsnip, radish, oriental radish, rutabaga, salsify, black salsify, Spanish salsify, skirret, and turnip.

**Crop Subgroup 3 - Bulb Vegetables (if to be harvested green)** - garlic, great-headed garlic, leek, green onion, Welsh onion, shallot.

Winter squash (including pumpkins).

- B. If not more than 2.2 pints (1.57 lb ai) per acre of *VISOR II XRT* was applied to the field, the following crops (as well as any listed under subsection C below) may be planted 60 days after the last application. A second application of an S-metolachlor-containing product to the following crops is prohibited within 60 days of the original application.**

**Crop Group 8 - Fruiting Vegetables, except Cucurbits and Tabasco Peppers:** eggplant, groundcherry (*Physalis* spp.), pepino, peppers (bell, chili, cooking, pimiento, and sweet), tomato, tomatillo, and tomato.

- C. If not more than 2.7 pints (1.9 lb ai) per acre of *VISOR II XRT* was applied to the field, the following crops may be planted 60 days after the last application. A second application of an S-metolachlor-containing product to the following crops is prohibited within 60 days of the original application.**

**Crop Subgroup 1C - Tuberous and Corn Vegetables:** arracacha; arrowroot; Chinese artichoke; Jerusalem artichoke; edible canna; bitter and sweet cassava; chayote (root); chufa; dasheen (taro); ginger; leren; potato; sweet potato; taniar; turmeric; yam bean; and yam, true.

**Crop Group 3 - Bulb Vegetables (if to be harvested dry)** - garlic, great-headed garlic, leek, dry bulb and green onion, Welsh onion, shallot.

**Crop Subgroup 4B - Leaf Petiole Vegetables:** cardoon, celery, Chinese celery, celtuce, Florence fennel, rhubarb, and Swiss chard.

**Crop Subgroup 5A - Head and Stem Brassica Vegetables:** broccoli, Chinese broccoli, Brussels sprouts, cabbage, Chinese (Napa) cabbage, Chinese mustard, cauliflower, cavalo broccolo, and kohlrabi.

## APPLICATION PROCEDURES

### APPLICATION TIMING

*VISOR II XRT* alone or in some tank mixtures with other labeled herbicides may be applied for weed control in certain crops at various times. Refer to specific crop use directions to determine if application timings listed below are allowed.

**Preplant Surface-Applied:** For minimum-tillage or no-tillage systems only, *VISOR II XRT* alone and some *VISOR II XRT* tank mixtures may be applied up to 45 days before planting certain crops. Use only split applications for treatments made 30 to 45 days before planting, with 2/3 the labeled broadcast rate for the crop and soil texture applied initially and the remaining 1/3 at planting. Treatments less than 30 days before planting may be made either as a split or a single application. Refer to individual crop section on this label to determine if early preplant surface application is indicated. If weeds are present at the time of treatment, apply in a tank mixture combination with a contact herbicide (for example Glyphosate or Paraquat). Observe directions for use, precautions, and restrictions on the label of the contact herbicide. To the extent possible, **DO NOT** move treated soil out of the row or move untreated soil to the surface during planting, or weed control will be diminished.

**Preplant Incorporated:** Apply *VISOR II XRT* to the soil and incorporate into the top 2 inches of soil within 14 days before planting, using a finishing disk, harrow, rolling cultivator, or similar implement capable of providing uniform 2-inch incorporation. Use a preplant incorporated application if furrow irrigation is used or when a period of dry weather after application is expected. If crop will be planted on beds, apply and incorporate this product after bed formation, unless specified otherwise.

**Preemergence:** Apply *VISOR II XRT* during planting (behind the planter) or after planting, but before weeds or crops emerge.

**Postemergence:** *VISOR II XRT* will not control emerged weeds so it must be applied to a weed-free soil surface or in tank mixture with products that provide postemergence control of weeds present at the time of application. Refer to the individual crop section of this label if a postemergence application is listed.

### SPECIAL APPLICATION PROCEDURES

**CA Only (Beans, Peas, and Lentils; Corn; Safflowers): Preplant Incorporated:** Broadcast *VISOR II XRT* alone or with tank mix partners listed on this label to the soil and thoroughly incorporate with a disk or similar implement set to till 4 to 6 inches deep. For more thorough incorporation, till the soil in 2 different directions (cross-till). Crops may be planted on flat surface or on beds. Use caution when forming the beds that only soil from the *VISOR II XRT* treated zone is used (i.e., untreated soil should not be brought to soil surface). If the application is made to preformed beds, incorporate this product with a tillage implement set to till 2 to 4 inches deep. Care should be taken during tilling to keep the tilled (*VISOR II XRT*-treated) soil on the beds.

**Preemergence:** Apply *VISOR II XRT* after planting. Water with sprinkler or flood irrigation within 7 to 10 days.

**Fall Application for Spring Weed Control (Only in IA, MN, ND, SD, WI, and portions of NE and IL - See specific instructions in the Beans, Peas, and Lentils; Corn; and Soybeans sections of this label for timing of application and other information):** Use on medium and fine soils with greater than 2.5% organic matter that will be planted to corn or soybeans the next spring. Ground may be tilled before or after application. **DO NOT** exceed a 2 to 3-inch incorporation depth if tilled after treatment.

#### Restrictions

- **DO NOT** apply to frozen ground.
- If a spring application is made, the total rate of the fall plus spring applications must not exceed the maximum total rate for the specific crop.

**Fall Application for Italian Ryegrass Control (Corn, Cotton, Grain and Forage Sorghum, and Soybean Only - See specific instructions in the Corn, Cotton, Grain and Forage Sorghum, and Soybean sections of this label for timing of application and other information):** *VISOR II XRT* may be applied in the fall (September 1 to December 1) for residual control of glyphosate-resistant Italian ryegrass (*Lolium multiflorum*). A tillage operation may precede the application. **DO NOT** incorporate to a depth greater than 2 to 3 inches if tillage follows the application of this product. All crops on the label for this product may be planted the following spring after application. Refer to the crop sections on this label for specific directions.

#### Restrictions

- **DO NOT** apply this product to frozen ground.
- If a spring application is made, the combined total amount of this product applied in the fall plus the spring must not exceed the maximum seasonal S-metolachlor rate for the specific crop/planted.

**Ground Application:** Apply *VISOR II XRT* alone or in tank mixtures by ground equipment in a minimum of 10 gallons of spray mixture per acre, unless otherwise specified.

Use sprayers that provide accurate and uniform application. For *VISOR II XRT* tank mixtures with wettable powder or dry flowable formulations, Use screens and strainers no finer than 50-mesh. Rinse sprayer thoroughly with clean water immediately after use.

Calculate the amount of herbicide needed for band treatment by the formula:

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast rate per acre} = \text{Amount needed per acre of field}$$

For information on applying in lower volumes of carrier, see **Low Carrier Application** section.

For application by air or through center pivot systems, see **Aerial Drift Management and Aerial Drift Reduction Advisory Information** sections.

For information on impregnating dry fertilizer, see **Dry Bulk Granular Fertilizer** section.

For information on application using variable-rate technologies, see **Variable-Rate Application** section.

### SPRAY EQUIPMENT

#### LOW CARRIER APPLICATION

##### For Broadcast Ground Application Only

Use sprayers, such as Ag-Chem RoGator<sup>®</sup>, Hagie, John Deere Hi-Cycle TM, Melroe Spracoupe, Tyler Patriot<sup>™</sup>, or Willmar Air Ride<sup>™</sup>, that provide accurate and uniform application. **Only water may be used as a carrier.** Screens in suction and in-line strainers should be 50-mesh. Manufacturers may require that tip screens as fine as 100-mesh be used with some nozzles. Use a pump with capacity to: (1) maintain up to 35 to 40 psi at the nozzles and (2) provide sufficient agitation in tank to keep mixture in suspension. Use a minimum of 5.0 gallons of spray mixture per acre. Maximum recommended sprayer speed is 15 mph. Rinse sprayer thoroughly with clean water immediately after each use.

#### Precautions

- Use low pressure nozzles to reduce drift and increase application accuracy.
- Take care when using automatic rate controlling devices to spray the material within the rated working pressure and flow ranges of the nozzles selected.

- Use nozzle screens when specified by the manufacturer.
- Place all nozzles on 20-inch centers, except flooding types which should be placed on 40-inch centers. When Flat Fan-type nozzles are used, angles of 80° or 110° are recommended.
- Always read and follow the manufacturer's directions for optimum setup and performance of their nozzles or tips.

#### AERIAL APPLICATION

Apply *VISOR II XRT* in water alone or in tank mixtures with Atrazine, Linuron or Metribuzin in a minimum total volume of 2.0 gallons per acre by aircraft. This product may also be applied by air in combination with Bentfluralin, Pendimethalin or Trifluralin. Avoid application under conditions where uniform coverage cannot be obtained or where excessive spray drift may occur. In order to assure that spray will be controllable within the target area when used according to label directions, make applications at a maximum height of 10 feet, using low-drift nozzles at a maximum pressure of 40 psi, and restrict application to periods when wind speed does not exceed 10 mph. To assure that spray will not adversely affect adjacent sensitive nontarget plants, apply *VISOR II XRT* alone or *VISOR II XRT* + Atrazine by aircraft at a minimum upwind distance of 400 feet from sensitive plants, or apply *VISOR II XRT* + Linuron or Metribuzin at a minimum upwind distance of 300 feet from sensitive plants.

#### Aerial Drift Management

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment- and weather-related factors determines the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops. These requirements **DO NOT** apply to forestry applications, public health uses, or to applications using dry formulations.

1. The distance of the outermost nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downward more than 45 degrees.
3. Where states have more stringent regulations, they must be observed.

The applicator should be familiar with and take into account the information covered in the **Aerial Drift Reduction Advisory Information** section below.

#### Aerial Drift Reduction Advisory Information Information on Droplet Size

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see **Wind, Temperature and Humidity, and Temperature Inversions**).

#### Controlling Droplet Size

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** - **DO NOT** exceed the nozzle manufacturer's recommended pressures. For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- **Number of Nozzles** - Use the minimum number of nozzles that provide uniform coverage.
- **Nozzle Orientation** - Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.
- **Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

#### Application Height

Applications should not be made at a height greater than 10 feet above the top of the largest plants, unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

## Swath Adjustment

When applications are made with a crosswind, the swath will be displaced downward. Therefore, on the upwind and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase with increasing drift potential (higher wind, smaller drops, etc.).

## Wind

Drift potential is lowest between wind speeds of 2 to 10 mph. However, many factors, including droplet size and equipment type, determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. **Note:** Local terrain can influence wind patterns.

## Temperature and Humidity

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is greatest when conditions are both hot and dry.

## Temperature Inversions

Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

## Sensitive Areas

The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, nontarget crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

Avoid application to humans or animals. Flagmen and loaders should avoid inhalation of spray mist and prolonged contact with skin.

## CENTER PIVOT IRRIGATION APPLICATION

This product alone or in tank mixture with other herbicides on this label, which are registered for center pivot application, may be applied in irrigation water preemergence (after planting, but before weeds or crop emerge) at rates specified on this label. This product also may be applied postemergence to the crop and preemergence to weeds in crops where postemergence applications are allowed on this label. Follow all restrictions (height, timing, rate, etc.) to avoid illegal residues. Apply this product only through a center pivot irrigation system. **DO NOT** apply this product through any other type of irrigation system. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water. If you have questions about calibration, you should contact State Extension specialists, equipment manufacturers, or other experts. **DO NOT** connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system, unless the pesticide label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

## Operating Instructions

1. The system must contain a functional check-valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent watersource contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quickclosing check-valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump or piston pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. **DO NOT** apply when wind speed favors drift beyond the area intended for treatment.
8. Prepare a mixture with a minimum of 1 part water to 1 part herbicide(s) and inject this mixture into the center pivot system. Injecting a larger volume of a more dilute mixture per hour will usually provide more accurate calibration of metering equipment. Maintain sufficient agitation to keep the herbicide in suspension.
9. Meter into irrigation water during entire period of water application.
10. Apply in 1/2 to 1 inch of water. Use the lower water volume (1/2 inch) on coarse textured soils and the higher volume (1 inch) on fine-textured soils. More than 1 inch of water at application may reduce weed control by moving the herbicide below the effective zone in the soil.

## Precaution for Center Pivot Applications

- Where sprinkler distribution patterns **DO NOT** overlap sufficiently, unacceptable weed control may result. Where sprinkler distribution patterns overlap excessively, crop injury may result.

## DRY BULK GRANULAR FERTILIZERS

Many dry bulk granular fertilizers may be impregnated or coated with **VISOR II XRT** alone or selected **VISOR II XRT** tank mixtures which are registered for preplant incorporated or preplant surface applications which are used to control weeds in crops on the label for this product and are not prohibited from use on dry bulk granular fertilizers.

When applying **VISOR II XRT** or **VISOR II XRT** mixtures with dry bulk granular fertilizers, follow all directions for use and precautions on the respective product labels, regarding target crops, rates per acre, soil texture, application methods (including timing of application), and rotational crops.

All individual state regulations relating to dry bulk granular fertilizer blending, registration, labeling, and application are the responsibility of the individual and/or company selling the herbicide/fertilizer mixture.

Prepare the herbicide/fertilizer mixtures by using any closed drum, belt, ribbon, or other commonly used dry bulk fertilizer blender. Nozzles used to spray **VISOR II XRT** and **VISOR II XRT** mixtures onto the fertilizer must be placed to provide uniform spray coverage. Care should be taken to aim the spray directly onto the fertilizer only and to avoid spraying the walls of the blender.

If the herbicide/fertilizer mixture is too wet, add a highly absorptive material, such as Agsorb® or Celatemp MP-79®, or similar granular clay or diatomaceous earth materials, to obtain a dry, free-flowing mixture. Add absorptive materials only after the herbicide has been thoroughly blended into the fertilizer mixture. Best application results will be obtained by using a granule of 6/30 particle size or of a size similar to that of the fertilizer material being used. Generally, less than 2% by weight of absorptive material will be needed. Avoid using more than 5% absorptive material by weight.

Calculate amounts of **VISOR II XRT**, Atrazine, Atrazine + Simazine, Ethafluralin, Linuron, Metribuzin or Simazine by the following formula:

$$\frac{2000}{\text{pounds of fertilizer per acre}} \times \frac{\text{pints per acre of liquid or flowable product}}{\text{pounds of fertilizer per acre}} = \frac{\text{pints of liquid or flowable product per ton of fertilizer}}{\text{pounds of fertilizer per acre}}$$
$$\frac{2000}{\text{pounds of fertilizer per acre}} \times \frac{\text{pounds per acre of dry product}}{\text{pounds of fertilizer per acre}} = \frac{\text{pounds of dry product per ton of fertilizer}}{\text{pounds of fertilizer per acre}}$$

## Pneumatic (Compressed Air) Application (**VISOR II XRT** Alone)

High humidity, high urea concentrations, low fertilizer use rates, and dusty fertilizer may cause fertilizer mixture to build up or plug the distributor head, air tubes, or nozzle deflector plates. To minimize buildup, premix **VISOR II XRT** with Exxon Aromatic 200 at a rate of 1.0 to 4.0 pints per gallon of this product. Aromatic 200 is a noncombustible/nonflammable petroleum product. Aromatic 200 may be used in either a fertilizer blender or through direct injection systems. Drying agents should not be used when using Aromatic 200.

## Precautions

- Use mixtures of this product and Aromatic 200 on dry fertilizer only. Poor results or crop injury may result if a mixture of this product and Aromatic 200 are used in water or liquid fertilizer solutions for spraying applications.
- When impregnating this product in a blender before application, a drier mixture can be attained by substituting a drying agent for Aromatic 200. The use of Agsorb FG or drying agents of 6/30 particle size are recommended.

## Restrictions

- To avoid potential for explosion, **DO NOT** impregnate this product or mixtures of this product on ammonium nitrate, potassium nitrate, or sodium nitrate, either alone or in blends with other fertilizers.
- **DO NOT** use this product or mixtures of this product on straight limestone, since absorption will not be achieved. Fertilizer blends containing limestone can be impregnated.
- **DO NOT** use drying agents with the On-The-Go impregnation equipment.

## Application

Apply 200 to 700 pounds of the herbicide/fertilizer mixture per acre. For best results, apply the mixture uniformly to the soil with properly calibrated equipment immediately after blending. Uniform application of the herbicide/fertilizer mixture is essential to prevent possible crop injury. Nonuniform application may also result in unsatisfactory weed control. In areas where conventional tillage is practiced, a shallow incorporation of the mixture into the soil may improve weed control. On fine- or medium-textured soils in areas where soil incorporation is not planned, i.e., reduced tillage situations or in some conventional till situations, make applications approximately 30 days before planting to allow moisture to move the herbicide/fertilizer mixture into the soil. On coarse-textured soils, make applications approximately 14 days prior to planting.

## Precaution

- To avoid crop injury, **DO NOT** use the herbicide/fertilizer mixture on crops where bedding occurs.

## MIXING INSTRUCTIONS

### VISOR II XRT Alone

Mix **VISOR II XRT** with water or fluid fertilizer and apply as a spray. Fill the spray tank 1/2 to 3/4 full with water or fluid fertilizer, add the proper amount of this product, and then add the rest of the water or fluid fertilizer. Provide sufficient agitation during mixing and application to maintain a uniform emulsion.

### VISOR II XRT in Tank Mixtures

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

### Tank Mix Instructions

For each mixture, check compatibility as described below before mixing in spray tank. When adding ingredients to the mixture, allow time for each ingredient to be thoroughly mixed before adding the next. Be sure to agitate during mixing and application to maintain a uniform suspension.

1. Fill the spray tank 1/4 full with water, and start agitation
2. Add the following herbicides and allow it to become dispersed
  - a. 2,4-D, 2,4-DB, Atrazine, Atrazine + Dicamba, Atrazine + Simazine, Benfluralin, Bentazon, Clomazone, Dicamba, EPTC, Ethalfluralin, Fluometuron, Glufosinate, Glufosinate + Atrazine, Imazethapyr, Isoxaflutole, Linuron, Metribuzin, Metribuzin + Chlorimuron, MSMA, Pendimethalin, Prometryn, Simazine or Trifluralin
3. Then add this product
4. Then add Paraquat, Glyphosate or Glyphosate + 2,4-D if these products are being used
5. Add the rest of the water

For tank mixtures with Atrazine, Atrazine + Dicamba, Atrazine + Simazine, Clomazone, Dicamba, EPTC, Ethalfluralin, Fluometuron\*, Imazaquin, Imazethapyr, Isoxaflutole, Linuron, Metribuzin, Metribuzin + Chlorimuron Ethyl, Pendimethalin\*, Prometryn, Simazine or Trifluralin, fluid fertilizers may replace all or part of the water as carrier, except in the Atrazine postemergence and the Dicamba postemergence tank mixes. For tank mixtures with Atrazine, see additional mixing instructions on the Atrazine label. For each mixture, check compatibility with fluid fertilizer, as described below, before mixing in spray tank. For all tank mixtures, agitate during mixing and application to maintain a uniform suspension.

\*See **Special Mixing Instructions** for tank mixtures with Fluometuron and with Atrazine or Simazine + Pendimethalin under the appropriate tank mixture section.

For directions on how to conduct a compatibility test, see the **Compatibility Test** section below.

## COMPATIBILITY TEST

A jar test is recommended before tank mixing to ensure compatibility of **VISOR II XRT** with other pesticides. The following test assumes a spray volume of 25 gallons per acre. For other spray volumes, make appropriate changes in the ingredients.

Nitrogen solutions or complete fluid fertilizers may replace all or part of the water in the spray. Because liquid fertilizers vary, even within the same analysis, **always check compatibility with pesticide(s) before use**. Incompatibility of tank mixtures is more common with suspensions of fertilizer and pesticides.

## Test Procedure

1. Add 1.0 pint of carrier (fertilizer or water) to each of 2 one quart jars with tight lids.
  - a. **Restrictions:** Use the same source of water that will be used for the tank mix and conduct the test at the temperature the tank mix will be applied.
2. To one of the jars, add 1/4 tsp. or 1.2 milliliters of a compatibility agent approved for this use, such as Inmix Envelop (1/4 tsp. is equivalent to 2.0 pints per 100 gallons spray). Shake or stir gently to mix.
3. To both jars, add the appropriate amount of pesticide(s) in their relative proportions based on specified label rates. If more than one pesticide is used, add them separately with dry pesticides first, flowables next, and emulsifiable concentrates last. After each addition, shake or stir gently to thoroughly mix.
4. After adding all ingredients, put lids on and tighten, and invert each jar ten times to mix. Let the mixtures stand 15 to 30 minutes and then look for separation, large flakes, precipitates, gels, heavy oily film on the jar, or other signs of incompatibility. Determine if the compatibility agent is needed in the spray mixture by comparing the two jars. If either mixture separates, but can be remixed readily, the mixture can be sprayed as long as good agitation is used. If the mixtures are incompatible, test the following methods of improving compatibility: (a) Slurry the dry pesticide(s) in water before addition, or (b) add 1/2 the compatibility agent to the fertilizer or water and the other 1/2 to the emulsifiable concentrate or flowable pesticide before addition to the mixture. If incompatibility is still observed, **DO NOT** use the mixture.
5. After compatibility testing is complete, dispose of any pesticide wastes in accordance with the **Storage and Disposal** section in this label.

## CROP USE DIRECTIONS

### CORN (ALL TYPES) - VISOR II XRT ALONE

Apply **VISOR II XRT**, either preplant surface, preplant incorporated, preemergence, postemergence, or lay-by, using the appropriate rate specified below.

### PREPLANT SURFACE-APPLIED

Refer to instructions for use of **VISOR II XRT** alone under **Application Procedures**.

### Fall Application for Spring Weed Control:

1. Apply after September 30 in ND, SD, MN, WI, and north of Route 30 in IA.
2. Apply after October 15 north of Route 91 in NE and south of Route 30 in IA.
3. Apply after October 31 north of Route 136 in IL.

In all locations, apply to crop stubble after harvest when the sustained soil temperature at a 4-inch depth is less than 55°F and falling. In minimum-till or no-tillage systems on soils having greater than 2.5% organic matter, use 2.2 to 2.7 pints (1.57 to 1.9 lb ai) per acre on medium-textured and 2.7 pints (1.9 lb ai) per acre on fine-textured soils. A tillage operation may precede the application. A fall and/or a spring tillage may follow application, but **DO NOT** exceed an incorporation depth greater than 2 to 3 inches. Minimize furrow and ridge formation in the tillage operations.

## Restrictions

- **DO NOT** apply to frozen ground.
- If a spring application is made, the total rate of the fall plus spring applications must not exceed the maximum total rate for corn 5.2 pints (3.7 lb ai) per acre depending on soil texture).



**Fall Application for Italian Ryegrass Control:** *VISOR II XRT* may be applied for residual control of glyphosate-resistant Italian ryegrass (*Lolium multiflorum*). Apply this product at 1.75 to 2.2 pints (1.25 to 1.57 lb ai) per acre in the fall (September 1 to December 1) after harvest of the previous crop and prior to Italian ryegrass emergence. Use the lower rate of this product for coarse-textured soils and the higher rate for fine-textured soils. A tillage operation may precede the application. If tillage follows the application of this product, avoid incorporating to a depth greater than 2 to 3 inches. For fall applications after emergence of glyphosate-resistant Italian ryegrass, Paraquat can be tank mixed with this product to control emerged ryegrass. Refer to the Paraquat label for specific rates, application instructions, and restrictions. Other registered herbicides may be tank mixed with this product for control or improved control of other weeds present at the time of application.

#### Restrictions

- **DO NOT** apply this product to frozen ground.
- If a spring application is made, the combined total amount of this product applied in the fall plus spring must not exceed the maximum seasonal S-metolachlor rate for corn 5.2 pints (3.7 lb ai) per acre depending on soil texture).

**Fall Application for Control or Suppression of Yellow Nutsedge (ID, OR, and WA only):** For preemergent control or suppression of yellow nutsedge the following spring, apply 1.75 pints (1.25 lb ai) per acre of *VISOR II XRT* in the fall after the harvest of the previous crop but before freeze-up. Fall applications of this product can be surface-applied or incorporated.

#### Restrictions

- Make no more than one fall application per crop.
- Apply no more than 1.75 pints (1.25 lb ai) per acre in a single fall preplant application.
- **DO NOT** apply to frozen ground.
- If a spring application is made, the combined total amount of this product applied in the fall plus spring must not exceed the maximum seasonal S-metolachlor rate for corn 5.2 pints (3.7 lb ai) per acre depending on soil texture).

#### EARLY PREPLANT APPLICATIONS

Use on medium- and fine-textured soils with minimum-tillage or no-tillage systems in CO, IA, IL, IN, KS, KY, MN, MO, MT, ND, NE, SD, TN, WI, and WY. Apply 2/3 the specified rate of *VISOR II XRT* 2.2 pints (1.57 lb ai) per acre on medium soils and 2.7 pints (1.9 lb ai) per acre on fine soils) as a split treatment 30 to 45 days before planting and the remainder at planting. Applications made less than 30 days prior to planting may be as either a split or single treatment. Apply 1.75 pints (1.25 lb ai) per acre on coarse soils not more than 2 weeks before planting.

#### Restrictions

- If a spring application is made, the total rate of the fall plus spring applications must not exceed the maximum total rate for corn 5.2 pints (3.7 lb ai) per acre depending on soil texture).

On medium- and fine-textured soils with minimum- or no-tillage systems in CT, DE, MA, MD, ME, MI, NH, NY, OH, PA, RI, VA, VT, and WV, preplant surface applications may be applied following the directions for use above. If the amount of rainfall results in unsatisfactory length of weed control following the earlier treatment, a postemergence application of an appropriately labeled broadleaf and/or grass weed herbicide may be used, i.e., Atrazine, Atrazine + Metolachlor/S-metolachlor, Bentazon, Bromoxynil, Dicamba, Nicosulfuron, Primisulfuron, Primisulfuron + Proflusulfuron or 2,4-D. If the postemergence treatment includes the herbicide used preplant surface-applied, **DO NOT** exceed the total labeled rate for corn on a given soil texture. Observe all directions for use, precautions and limitations on the label of the postemergent herbicide.

#### PREPLANT INCORPORATED OR PREEMERGENCE

Follow instructions for use of *VISOR II XRT* alone under **Application Procedures**. On coarse soils, apply 1.3 to 1.75 pints (0.93 to 1.25 lb ai) per acre of this product if organic matter content is less than 3%, or 1.75 pints (1.25 lb ai) per acre if organic matter content is 3% or greater. On medium soils, apply 1.75 to 2.2 pints (1.25 to 1.57 lb ai) per acre of this product. On fine soils, apply 1.75 to 2.2 pints (1.25 to 1.57 lb ai) per acre of this product if organic matter content is less than 3%, or 2.2 pints to 2.7 pints (1.57 to 1.9 lb ai) per acre if organic matter content is 3% or greater.

#### POSTEMERGENCE OR LAY-BY

To extend the duration of weed control in corn, a maximum rate of 2.7 lb ai (1.9 lb ai) per acre of *VISOR II XRT* may be applied after corn emergence until the corn plants reach 40 inches in height, following any preplant surface-applied, preplant incorporated, or preemergence herbicide application, including this product. For best results, applications should be made to soil free of emerged weeds and directed toward the base of corn plants in excess of 5 inches tall.

#### Restrictions for all applications to corn

- **DO NOT** graze or feed forage from treated areas for 30 days following application.
- The total rate of this product applied on corn during any one crop year should not exceed 5.2 pints (3.7 lb ai) per acre, depending on soil texture.
- **Preharvest Interval (PHI): DO NOT** harvest sweet corn ears from treated areas for 30 days following application.

#### PROBLEM WEED CONTROL DIRECTIONS

**Shattercane, Wild Proso Millet, Woolly Cupgrass, and Eclipta - Partial Control:** For more consistent partial control of shattercane, wild proso millet, woolly cupgrass, or eclipta, apply 1.3 to 1.75 pints (0.93 to 1.25 lb ai) per acre of *VISOR II XRT* preplant incorporated followed by 1.3 to 1.75 pints (0.93 to 1.25 lb ai) per acre of this product preemergence. Make the preemergence application during or after planting, but before weeds and corn emerge. Apply the 1.75 pints (0.93 to 1.25 lb ai) per acre rate of this product when a heavy infestation of shattercane, wild proso millet, woolly cupgrass, or eclipta is expected. A shallow cultivation may be needed to control any late emerging weeds.

**Woolly Cupgrass and Wild Proso Millet Control Program:** For control of these species, use the following 3-step program:

1. Apply *VISOR II XRT* early preplant, preplant incorporated, or preemergence at 2.2 pints (1.57 lb ai) per acre on medium soils and 2.7 lb ai (1.9 lb ai) per acre on fine-textured soils, up to the maximum label rate. Lightly incorporate with a rotary hoe if rainfall does not occur within 5 to 7 days.
2. Apply a postemergence tank mix of labeled rates of Primisulfuron-methyl or Proflusulfuron + Primisulfuron-methyl plus Nicosulfuron plus 1.0 quart of crop oil concentrate plus 1.0 gallon per acre of 28% nitrogen, or the equivalent amount of ammonium sulfate, when grasses are 2 to 3 inches tall and the corn is at least 4 inches tall.
3. Cultivate 14 to 21 days after the postemergence application.

#### Restrictions

- **DO NOT** apply more than the labeled application rate for a given soil texture per year, either as a single or split treatment applications.
- In corn, use up to 3.3 pints (2.35 lb ai) per acre of this product preplant surface, preplant incorporated or preemergence treatment on soils having an organic matter contents between 6% and 20%.
- In the event of escape of annual weeds following a preplant surface, preplant incorporated, or preemergence treatment of this product, follow with a postemergence application of an appropriately labeled broadleaf and/or grass weed herbicide, e.g., Atrazine, Atrazine + Metolachlor/S-metolachlor, Bentazon, Bromoxynil, Dicamba, Nicosulfuron, Primisulfuron, Primisulfuron + Proflusulfuron, or 2,4-D. If the postemergence treatment includes the herbicide used in the earlier treatment, **DO NOT** exceed the total labeled rate for corn on a given soil texture.
- Bromoxynil may be applied postemergence alone or in tank mix combination with Atrazine. **DO NOT** exceed 1.2 lbs. a.i. per acre of Atrazine in tank mix combination with Bromoxynil postemergence.
- **DO NOT** use this product on peat or muck soils.

#### CORN - *VISOR II XRT* COMBINATIONS

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

*VISOR II XRT* in any tank mixture for corn may be applied in water or fluid fertilizer before corn emerges. Use only water as a carrier when this product is applied after corn emergence.

#### Restrictions for all applications to corn

- **DO NOT** graze or feed forage from treated areas for 30 days following application.
- **Preharvest Interval (PHI): DO NOT** harvest sweet corn ears from treated areas for 30 days following application.

**Important: For Tank Mixtures with Atrazine** — If applying this product in tank mixture with Atrazine, all the restrictions and rate limitations on the Atrazine label must be followed, if more restrictive/protective than those on this label. In addition, if Atrazine is/must be applied at rates lower than those specified on this label, broadleaf weed control may be affected. Refer to the Atrazine label for weeds controlled at the reduced rates.

**Table 2: VISOR II XRT Tank Mixtures for Corn - Additional Weeds Controlled and Special Instructions**

	<b>VISOR II XRT + Atrazine and/or Simazine (Preplant Surface, PPI, PRE)</b>	<b>VISOR II XRT + Atrazine (Post)</b>	<b>VISOR II XRT + Dicamba (Field Corn)</b>	<b>VISOR II XRT + Atrazine + Linuron</b>	<b>VISOR II XRT + Atrazine or Simazine + Pendimethalin</b>	<b>VISOR II XRT + Atrazine + Dicamba</b>	<b>VISOR II XRT + Isoxaflutole</b>
<b>Special Mixing Instructions</b>					1		
<b>Comments</b>	2, 3, 4, 5	2, 3		2, 3	2, 3	4	4
Browntop panicum	X			X	X		X
Cocklebur	X	0	0	X	X		0-X
Common purslane	X			X	X	X	X
Hairy nightshade	X			X	X		X
Jimsonweed		X	0			X	X
Kochia		X				X	X
Lambsquarters	X	X	X	X	X	X	X
Morningglory	X	0	0	X	X		X
Mustard		X				X	X
Pigweed				X	X	X	X
Prickly sida		X					
Ragweed	X	X	X	X	X	X	X
Smartweed	X	X	X	X	X	X	X
Velvetleaf	X	X	0	X	X	0-X	0-X

X = control; 0 = partial control; 0-X = partial to full control depending on ratio of products used or on weed population

#### Instructions - Table 1: VISOR II XRT in Tank Mixtures

##### 1. Special Mixing Instructions for VISOR II XRT + Atrazine or Simazine and Pendimethalin.

- Fill the spray tank 1/4 full with water or fluid fertilizer and start agitation.
- To aid compatibility, add a compatibility agent, at 4 pints per 100 gallons of spray mixture.
- Then add the Atrazine or Simazine and allow it to become dispersed.
- Then add this product and Pendimethalin.
- Add the rest of the water.

##### 2. See additional mixing instructions on the Atrazine label.

3. **DO NOT** apply more than a total of 2.5 lbs. a.i. of Atrazine per acre per year. However, certain states may have established rate limitations for Atrazine within specific geographical areas. Consult your state lead pesticide control agency for additional information. It is a violation of this label to deviate from state use regulations.

4. In Minimum-Tillage and No-Tillage systems, mix with Paraquat for control of most emerged annual weeds and suppression of perennial weeds; or with Glyphosate + 2,4-D for suppression of emerged Field bindweed and control or suppression of annual weeds; or with Glyphosate for control of most emerged annual and perennial weeds.

5. Refer to **VISOR II XRT Combination – Tank Mixture with Atrazine; or Atrazine + 2,4-D; or Atrazine + 2,4-D + Dicamba for Minimum-Tillage or No-Tillage Systems** sections for specific directions for 2,4-D or Dicamba burndown combinations in Minimum-Tillage and No-Tillage systems.

**VISOR II XRT** in any tank mixture for corn may be applied in water or fluid fertilizer, except as noted. Refer to **Corn (All Types) – VISOR II XRT** Alone for specified sequential postemergence treatments if escape weeds develop.

#### Restrictions

- For all applications to corn, **DO NOT** graze or feed forage from treated areas for 30 days following application.
- Preharvest Interval (PHI): DO NOT** harvest sweet corn ears from treated areas for 30 days following application.
- When applying this product in tank mixture with Atrazine, **DO NOT** exceed a total of 2.5 lbs. a.i. of atrazine per acre per year.

#### TANK MIXTURE WITH ATRAZINE OR SIMAZINE, OR ATRAZINE + SIMAZINE - PREPLANT SURFACE, PREPLANT INCORPORATED, OR PREEMERGENCE

In addition to the weeds controlled by this product alone, this product + Atrazine or Simazine, or this product + Atrazine + Simazine, applied preplant surface, preplant incorporated, or preemergence, also controls the following weeds: browntop panicum, cocklebur, common purslane, hairy nightshade, lambsquarters, morningglory, ragweed, smartweed, and velvetleaf. Apply this product + Atrazine or Simazine, or this product + Atrazine + Simazine either preplant surface, preplant incorporated, or preemergence.

**Preplant Surface-Applied:** Follow instructions for use of this product alone under **Application Procedures** and under application instructions for this product alone on corn. Apply labeled rates Atrazine or Simazine, or Atrazine + Simazine combined with 2.2 pints (1.57 lb ai) per acre of this product on medium soils or with 2.2 to 2.7 pints (1.57 to 1.9 lb ai) per acre of this product on fine soils in minimum tillage and no-tillage systems in CO, IA, IL, IN, KS, KY, MN, MO, MT, ND, NE, SD, TN, VI, and WY. Apply the tank mixtures as a split or single treatment in those states and as indicated in the **VISOR II XRT Alone - Preplant Surface-Applied** section of the label for Corn. On coarse soils, apply 1.75 pints (1.25 lb ai) per acre of this product and labeled rates of Atrazine or Simazine, or Atrazine + Simazine combined.

**Preplant Incorporated or Preemergence:** Follow instructions for use of this product alone under **Application Procedures**. This product may be applied in tank mix combination with labeled rates of Atrazine or Simazine or Atrazine + Simazine. Apply this product according to the rates in **Table 3**.

#### Restriction

- **DO NOT** apply more than the labeled rate for a given soil texture per year, either as a split or single treatment.

#### Shattercane and Wild Proso Millet - Partial Control

For more consistent partial control of shattercane or wild proso millet where **VISOR II XRT** is applied in tank mixture or sequentially with other registered corn herbicides, the following applications may be made:

1. Apply 1.3 to 1.75 pints (0.93 to 1.25 lb ai) per acre of this product + 2.0 lbs. a.i. per acre of Atrazine or Simazine preplant incorporated, followed by 1.3 to 1.75 pints (0.93 to 1.25 lb ai) per acre of this product preemergence. Make the preemergence application during or after planting, but before weeds and corn emerge.
2. Apply this product at 1.75 pints (1.25 lb ai) per acre alone or in tank mix combination with up to 2 lbs. a.i. per acre of Atrazine or Simazine, preplant incorporated. **DO NOT** exceed the total rate of triazine herbicide specified in combination with **VISOR II XRT** for corn grown on a given soil texture. Follow with a post-directed application of Ametryn at labeled rates. Refer to the Ametryn label for specific directions for the post-directed application.
3. Apply EPTC or butylate formulations at labeled rates preplant incorporated, followed by a preemergence application of **VISOR II XRT** at 1.3 to 1.75 pints (0.93 to 1.25 lb ai) per acre. **DO NOT** use EPTC or butylate on soils where rapid degradation has been shown to occur. Make the preemergence application during or after planting, but before weeds and corn emerge.

#### Precaution

- When following the application regimes in numbers 1 to 3 above, a shallow cultivation may be needed after the preemergence or postemergence application to help control any late emerging shattercane or wild proso millet plants.

#### Restriction

- **DO NOT** apply more than 2.7 lb ai (1.9 lb ai) per acre of this product in the preplant incorporated plus preemergence application on soils with less than 6% organic matter.

**Table 3: *VISOR II XRT* + Atrazine or Simazine, or *VISOR II XRT* + Atrazine + Simazine, Preplant Incorporated or Preemergence - Corn (All Types)**

Soil Texture	Broadcast Rates Per Acre			
	Less than 3% Organic Matter		3% Organic Matter or Greater	
	<i>VISOR II XRT</i> + Atrazine* OR Simazine*	– OR – <i>VISOR II XRT</i> + Atrazine** + Simazine**	<i>VISOR II XRT</i> + Atrazine* OR Simazine*	– OR – <i>VISOR II XRT</i> + Atrazine** + Simazine**
Coarse	1.1 to 1.3 (0.78 to 0.93 lb ai) + See label for application rates	1.1 to 1.3 (0.78 to 0.93 lb ai) + Use 1:1 ratio See label for application rates	1.3 pints (0.93 lb ai) + See label for application rates	1.3 pints (0.93 lb ai) + Use 1:1 ratio See label for application rates
Medium	1.3 to 1.75 pints (0.93 to 1.25 lb ai) + See label for application rates	1.3 to 1.75 pints (0.93 to 1.25 lb ai) + Use 1:1 ratio See label for application rates	1.75 pints (1.25 lb ai) + See label for application rates	1.75 pints (1.25 lb ai) + Use 1:1 ratio See label for application rates

Soil Texture	Broadcast Rates Per Acre			
	Less than 3% Organic Matter		3% Organic Matter or Greater	
	<i>VISOR II XRT</i> + Atrazine* OR Simazine*	– OR – <i>VISOR II XRT</i> + Atrazine** + Simazine**	<i>VISOR II XRT</i> + Atrazine* OR Simazine*	– OR – <i>VISOR II XRT</i> + Atrazine** + Simazine**
Fine	1.75 pints (1.25 lb ai) + See label for application rates	1.75 pints (1.25 lb ai) + Use 1:1 ratio See label for application rates	1.75 to 2.2 pints (1.25 to 1.57 lb ai) + See label for application rates***	1.75 to 2.2 pints (1.25 to 1.57 lb ai) + Use 1:1 ratio See label for application rates
Muck or Peat (soils with more than 20% organic matter)	DO NOT USE			

\* Use Simazine in preference to Atrazine when heavy infestations of crabgrass or fall panicum are expected. On soils having between 6% and 20% organic matter, this product may be used up to 3.1 pints (2.21 lb ai) per acre in tank mix combination with Atrazine. Refer to the Atrazine label for rates, restrictions and limitations.

\*\* When using the tank mixture of this product + Atrazine + Simazine, use a 1:1 ratio of each when heavy broadleaf weed infestations are expected. When heavy infestations of crabgrass or fall panicum are expected, use a 1:2 ratio of Atrazine + Simazine.

\*\*\* For cocklebur, yellow nutsedge, and velvetleaf control on fine-textured soils above 3% organic matter, apply 1.75 to 2.2 pints (1.25 to 1.57 lb ai) per acre of this product with labeled rate of Atrazine, or the same total amount of Atrazine + Simazine.

#### TANK MIXTURE WITH ATRAZINE - POSTEMERGENCE

##### Weeds Controlled

Barleygrass (watergrass)	Lambsquarters
Crabgrass	Mustard
Crowfootgrass	Pigweed
Fall panicum	Prickly sida
Giant foxtail	Purslane
Yellow foxtail	Ragweed
Jimsonweed	Smartweed
Kochia	Velvetleaf

##### Weeds Partially Controlled

Cocklebur
Morningglory
Yellow nutsedge

Apply labeled rate of atrazine with 1.3 pints (0.93 lb ai) of this product per acre on coarse soils, with 1.75 pints (1.25 lb ai) of this product per acre on medium soils or with 1.75 to 2.2 pints (1.25 to 1.57 lb ai) of this product per acre on fine soils\*. Apply this tank mixture before grass and broadleaf weeds pass the 2-leaf stage and before corn exceeds 5 inches in height. Application to weeds larger than the 2-leaf stage will generally result in unsatisfactory control.

**Lay-by:** Apply to corn plants not more than 12 inches tall. Applications to corn in excess of 5 inches should be directed to the base of the corn plants; whereas, applications to corn plants less than 5 inches tall may be made over the top. Occasionally, some corn leaf burn may result, but this should not affect later growth or yield. **DO NOT** apply this postemergence tank mixture in fluid fertilizer, or severe corn injury may occur.

\* For better control of cocklebur, morningglory, velvetleaf, and yellow nutsedge on fine-textured soils above 3% organic matter, apply higher labeled rate of Atrazine with 1.75 to 2.2 pints (1.25 to 1.57 lb ai) per acre of **VISOR II XRT**.

Tank mixtures of **VISOR II XRT** + Atrazine may be applied following use of any registered preplant surface-applied, preplant incorporated, or preemergence corn herbicide, including **VISOR II XRT** + Atrazine.

### Restriction

- **DO NOT** apply more than 5.2 pints (3.7 lb ai) of this product or more than 2.5 lbs. a.i. of Atrazine per acre during any one crop year. Refer to the atrazine label for geographic, soil-texture, and rotational restrictions.

### TANK MIXTURE WITH DICAMBA

**Preemergence:** Use this tank mixture only on field corn which is flat-planted (no furrows) in CO, IA, IL, IN, KS, MN, NE, OH, SD, and WI.

In addition to the weeds controlled by *VISOR II XRT* alone, *VISOR II XRT* + Dicamba, applied preemergence, also controls lambsquarters, ragweed, smartweed, cocklebur\*, jimsonweed\*, morningglory\*, and velvetleaf\*.

\* Partially controlled.

Apply *VISOR II XRT* + Dicamba preemergence. Broadcast labeled rate of Dicamba with 1.75 pints (1.25 lb ai) of this product per acre on medium soils, or with 1.75 to 2.2 pints (1.25 to 1.57 lb ai) of this product per acre on fine soils. Apply this tank mixture to the soil surface at planting or after planting, but before corn emerges. Plant corn at least 1.5 inches deep and apply behind planting equipment, avoiding incorporation by the planter wheel or other seed covering device. **DO NOT** incorporate before corn emergence. If it is necessary to rotary hoe to break the soil crust, **DO NOT** disturb the soil more than 1/2 inch deep.

### Precaution

- Avoid drift to sensitive nontarget plants, such as soybeans, during application, or injury may occur.

### Restrictions:

- **DO NOT** apply on coarse soils or on soils with less than 2.5% organic matter.
- **DO NOT** apply by aircraft.

### Postemergence for Control of Pigweed (Mid-Atlantic states, including DE, MD, PA, VA, and WV):

Apply 1.3 to 2 pints (0.93 to 1.42 lb ai) of this product per acre + labeled rate of Dicamba by ground equipment when pigweed plants are less than 3 inches tall and before corn exceeds 5 inches in height in a minimum of 20 gallons of spray per acre. Use the lower rate on coarse-textured and low organic matter soils. Use the higher rate on fine-textured and high organic matter soils.

### Precaution

- Avoid drift to sensitive nontarget plants, such as soybeans, during application, or injury may occur.

### Restriction

- **DO NOT** apply with aircraft.

### TANK MIXTURE WITH ATRAZINE OR SIMAZINE + PENDIMETHALIN FOR PROLONGED CONTROL OF LAMBSQUARTERS AND PIGWEED IN FIELD CORN ONLY (NORTHEAST U.S., INCLUDING MI, IN, KY, AND STATES EAST OF THESE)

For prolonged control of lambsquarters and pigweed, in addition to a broad spectrum of annual broadleaf and grass weeds, *VISOR II XRT* in tank mix combination with labeled rates of Atrazine or Simazine + Pendimethalin may be applied after planting, but before corn or weeds emerge. Apply by ground equipment in a minimum of 10 gallons of water or 20 gallons of liquid fertilizer. Apply by air in a minimum of 5.0 gallons of water. Apply this product according to rates in **Table 3**.

### Mixing Instructions: See Comment No.1 following **Table 2**.

Observe all directions for use, precautions and limitations on the respective product labels when applying these products in tank mix combinations. Refer to the Pendimethalin label for replanting instructions in the event of crop loss.

### TANK MIXTURE WITH ATRAZINE OR SIMAZINE, OR ATRAZINE + SIMAZINE WITH PARAQUAT, GLYPHOSATE + 2,4-D OR GLYPHOSATE FOR MINIMUM-TILLAGE OR NO-TILLAGE SYSTEMS

In minimum-tillage or no-tillage systems where corn is planted directly into a cover crop, stale seedbed, established sod, or previous crop residues, the contact herbicides Paraquat, Glyphosate + 2,4-D or Glyphosate may be tank mixed with *VISOR II XRT* + Atrazine or Simazine or *VISOR II XRT* + Atrazine + Simazine. See Comment No. 4 under **Table 1**.

*VISOR II XRT*, *VISOR II XRT* + Atrazine or Simazine, or *VISOR II XRT* + Atrazine + Simazine portion of the tank mixture provides preemergence control of the weeds listed on this label in the tank mixture section for *VISOR II XRT*, *VISOR II XRT* + Atrazine or Simazine, or *VISOR II XRT* + Atrazine + Simazine - Preplant Surface, Preplant Incorporated, or Preemergence.

**Application:** Apply before, during, or after planting, but before the corn emerges, at rates specified in the Paraquat, Glyphosate + 2,4-D or Glyphosate label.

**Paraquat:** See the Paraquat label for weeds controlled, specified rates and other information concerning use. This treatment will not control weeds taller than 6 inches.

**Glyphosate + 2,4-D:** See the Glyphosate + 2,4-D label for weeds controlled, specified rates for specific weeds, and other information concerning use.

**Glyphosate:** See the Glyphosate label for weeds controlled, specified rates, and other use precaution directions.

Apply in 20 to 60 gallons of water or fluid fertilizer per acre with ground equipment.

Apply labeled rate of Atrazine\* or Simazine\* or Atrazine\*\* + Simazine\*\* with 1.3 pints (0.93 lb ai) of this product per acre on coarse soils, with 1.75 pints (1.25 lb ai) of this product per acre on medium soils or with 1.75 to 2.2 pints (1.25 to 1.57 lb ai) of this product per acre on fine soils\*\*\*.

\* Use Simazine in preference to Atrazine when heavy infestations of crabgrass or fall panicum are expected.

\*\* When using the tank mixture of this product and Atrazine + Simazine, use equal rates of Atrazine and Simazine (1:1 ratio) when heavy broadleaf weed infestations are expected. When heavy infestations of crabgrass or fall panicum are expected, use a 1:2 ratio of Atrazine + Simazine.

\*\*\* For Cocklebur, Yellow nutsedge, and Velvetleaf control on fine-textured soils above 3% organic matter, apply 1.75 to 2.2 pints (1.25 to 1.57 lb ai) per acre of this product with labeled rate of Atrazine, or the same total amount of Atrazine + Simazine.

### Restriction

- **DO NOT** apply combinations containing Paraquat in suspension-type liquid fertilizers because the activity of paraquat will be reduced.

### TANK MIXTURE WITH ATRAZINE- OR ATRAZINE + 2,4-D; OR ATRAZINE + 2,4-D + DICAMBA FOR MINIMUM-TILLAGE OR NO-TILLAGE SYSTEMS

In minimum-tillage or no-tillage systems where corn is planted directly into a cover crop, stale seedbed, established sod, or previous crop residues, *VISOR II XRT* applied in combination with Atrazine will kill most emerged small annual weeds. Apply *VISOR II XRT* + Atrazine before, during, or after planting, but before corn emerges, according to the rates in **Table 3**.

Where heavy crop residues exist, add labeled rate of 2,4-D amine to the spray tank last and apply in a minimum of 25 gallons of carrier per acre.

As carriers, nitrogen solutions and complete liquid fertilizers, applied before corn emergence, enhance burndown of existing weeds, and therefore, are specified instead of water. Add Voyager 90/10 surfactant at 1.0 to 2.0 quarts per 100 gallons of diluted spray, or another appropriate surfactant at its specified rate, or add crop oil concentrate plus 28% liquid nitrogen (or equivalent). Apply before weeds exceed 3 inches in height. If alfalfa is present, add Dicamba to the spray mixture at labeled rates and apply before alfalfa exceeds 6 inches in height.

For fields with existing sod grasses (e.g., bromegrass, orchardgrass, rye, or timothy), when existing weeds exceed 3 inches in height or when very dry conditions exist, add Paraquat at specified rate in place of or in addition to 2,4-D, as indicated above. Observe all directions for use, precautions and restrictions on the respective product labels when applying these products in tank mix combinations.

### Restriction

- **DO NOT** apply Paraquat brands in suspension-type liquid fertilizer.

### TANK MIXTURE WITH ATRAZINE + DICAMBA IN CONSERVATION TILLAGE - FIELD AND SILAGE CORN

In conservation tillage systems where corn is planted directly into a cover crop or previous crop residue, *VISOR II XRT* plus Atrazine + Dicamba will kill most emerged small annual weeds. Apply *VISOR II XRT* + Atrazine + Dicamba before, during, or after planting, but before corn emergence on medium and fine soils with greater than 2.5% organic matter. For fields with

existing vegetation exceeding 3 inches in height or when very dry conditions exist, add Paraquat at its standard rate. **VISOR II XRT** + Atrazine + Dicamba may be applied postemergence to corn less than 3 inches tall and before weeds grasses exceed the 2-leaf stage.

As carriers, nitrogen solutions and complete liquid fertilizers, applied before corn emergence, enhance burndown of existing weeds.

Refer to the Atrazine + Dicamba label and follow all directions, limitations, precautions and restrictions regarding application and use in corn.

#### Restriction

- **DO NOT** apply Paraquat in suspension-type liquid fertilizer or use on emerged corn.

#### TANK MIXTURE WITH ISOXAFLOLUTE - FIELD CORN ONLY

This product and Isoxaflofute have a complementary response and weed control profile which allows various tank mix rate combinations to be considered. The addition of Isoxaflofute will improve the control of certain problem weeds including Texas panicum, woolly cupgrass and wild proso millet. **VISOR II XRT** improves both the duration and spectrum of annual grass and small-seeded broadleaf weed control, in particular foxtails (yellow foxtail), witchgrass and yellow nutsedge.

The rates of this product and Isoxaflofute are inversely proportional. Higher rates of this product are used with lower rates of Isoxaflofute and vice versa. The appropriate rate is selected based on your specific conditions, the weeds being targeted and your level of acceptance for potential crop damage.

**Examples:** (1) Where Texas panicum, woolly cupgrass or wild proso millet is a primary target weed, use a tank mix combination with a higher Isoxaflofute rate for the given soil type. (2) If your acceptance of an adverse crop response risk is low and/or a more general weed spectrum is targeted (especially yellow foxtail, witchgrass or yellow nutsedge), use a tank mix combination with a higher rate of this product for the given soil type. (3) If target weed is listed as controlled on both product labels, a tank mix consisting of intermediate rates of both products may be used.

#### Precautions

- To minimize risk to crops, refer to the Isoxaflofute label for specific restrictions regarding soil textures, pH, percent organic matter and other important considerations.
- Where a target weed is listed as controlled on only one of these two products, **DO NOT** apply less than the application rate for that weed as specified in the individual product label or unacceptable control may result. The most restrictive combination of instructions, limitations and other restrictions from all the products being applied must be followed.

**For coarse-textured soils:** Apply labeled rate of Isoxaflofute with 1.3 to 1.75 pints (0.93 to 1.25 lb ai) of this product per acre.

#### Restriction

- **DO NOT** use Isoxaflofute on coarse-textured soils with less than 1.5% organic matter.

**For medium-textured soils:** Where lower labeled rate of Isoxaflofute is used, rates as low as 1.75 pints (1.25 lb ai) per acre of this product may be applied. Where higher labeled rate of Isoxaflofute is used, rates as low as 1.3 pints (0.93 lb ai) per acre of this product may be applied. This product can be used in combination with Isoxaflofute at rates up to 2.2 pints (1.57 lb ai) per acre on medium-textured soils.

**For fine-textured soils:** Where lower labeled rate of Isoxaflofute is used, rates as low as 1.75 pints (1.25 lb ai) per acre of this product may be applied if the soil organic matter is less than 3%. If the soil organic matter is 3% or greater, 2.2 pints (1.57 lb ai) per acre of this product should be applied. Where mid-range labeled rates of Isoxaflofute are used, rates as low as 1.75 pints (1.25 lb ai) per acre of this product may be applied. Where higher labeled rate of Isoxaflofute is used, rates as low as 1.3 pints (0.93 lb ai) per acre of this product may be applied. This product can be used in combinations with Isoxaflofute at rates up to 2.7 lb ai (1.9 lb ai) per acre on fine-textured soils if the organic matter is 3% or greater.

#### TANK MIXTURES FOR POSTEMERGENCE SALVAGE WEED CONTROL IN FIELD CORN ONLY

For postemergence control of weeds in specific types of field corn, the **VISOR II XRT** combinations listed below may be used. Full season weed control from early preplant, preplant incorporated or preemergence treatments can lead to maximum yield potential under competition-free conditions. However, if control of emerged weeds is needed, a

postemergence program listed below can be applied to provide residual control for the remainder of the season.

Follow all label directions, instructions, precautions, and limitations for each product used. For each tank mixture with this product, apply only to the specific field corn type specified on the tank mix product label.

#### Precautions

- In-row weed control may be reduced because of lack of coverage when applied to corn over 4 inches tall.
- **DO NOT** use fluid fertilizer with these mixtures or corn injury may occur.

#### **VISOR II XRT + Glufosinate: Postemergence Use in Glufosinate-Resistant Corn**

These tank mixtures can be applied postemergence to weeds and corn from seed designated as Glufosinate-resistant. Glufosinate provides postemergence control of a broad spectrum of grass and broad leaf weeds and the **VISOR II XRT** provides residual control of grasses and certain broad leaf weeds listed in the label section **VISOR II XRT Applied Alone - Weeds Controlled**. Refer to section **VISOR II XRT Alone - Preplant Incorporated or Preemergence** and use the minimum rate per soil texture and organic matter classification for season-long residual control from this tank mix combination with Glufosinate. Refer to the Glufosinate label for the postemergence application rates according to weed species and their maximum height at the time of postemergence application. Where multiple weed species are present, use the highest specified rate to control the species and growth stages present.

Follow all applicable use directions, limitations, precautions, and information regarding application to corn on the **VISOR II XRT** and Glufosinate labels. Where difficult species and/or severe weed populations are expected, use the maximum rate where rate ranges are listed.

#### **VISOR II XRT + Glyphosate for Postemergence Application to Glyphosate-Resistant Corn**

The tank mixture of **VISOR II XRT** + Glyphosate can be applied postemergence to weeds and to corn designated as glyphosate-resistant. Application may be applied postemergence to glyphosate-resistant corn from emergence until corn reaches 30 inches tall or the V8 stage (8 leaves with collars), whichever comes first. This mixture will provide postemergence control of weed species on the Glyphosate label and residual control of weed species on the **VISOR II XRT** label. Use the minimum rate of this product postemergence with Glyphosate in glyphosate-tolerant corn as specified in the **Corn - VISOR II XRT Alone - Preplant Incorporated or Preemergence** section of this label according to soil texture and organic matter. Refer to the Glyphosate label and follow appropriate use directions, application procedures, precautions, and limitations. Refer to the Glyphosate label for directions for control of problem species. Where difficult species and/or severe weed populations are expected, use the maximum rate where rate ranges are listed.

#### **VISOR II XRT + Glyphosate + Atrazine for Postemergence Application to Glyphosate-Resistant Corn**

The tank mixture of **VISOR II XRT** + Atrazine + Glyphosate can be applied postemergence to weeds and to corn designated as glyphosate-resistant. Application may be applied postemergence to glyphosate-tolerant corn from emergence up to 12 inches in height. This mixture will provide postemergence control of weed species on the Glyphosate label and residual control of weed species on the **VISOR II XRT** + Atrazine label. Use the minimum **VISOR II XRT** + Atrazine rate postemergence with Glyphosate in glyphosate-resistant corn as specified in the **Corn - VISOR II XRT Combinations - Tank Mixture with Atrazine or Simazine, or Atrazine + Simazine - Preplant Incorporated or Preemergence** section and **Table 3** of this label according to soil texture and organic matter.

Follow all applicable use directions, limitations, precautions, and information regarding application to corn on the **VISOR II XRT**, Atrazine and Glyphosate labels for application to glyphosate-resistant corn. Where difficult species and/or severe weed populations are expected, use the maximum rate where rate ranges are listed.

#### **COTTON - VISOR II XRT ALONE**

**Application:** Apply **VISOR II XRT** postemergence to cotton and preemergence to weeds at 0.65 to 1.75 pints (0.46 to 1.25 lb ai) per acre, according to the state limitations in the following Postemergence section.

**AR, KS, LA, MS, TN, and Bootheel of MO:** Apply **VISOR II XRT** preemergence only at the rate of 0.65 to 1.3 pints (0.46 to 0.93 lb ai) per acre on sandy loams, 0.9 to 1.75 (0.64 to 1.25 lb ai) per acre on medium soils, or 1.3 to 1.75 pints (0.93 to 1.25 lb ai) per acre on fine soils.

**NM, OK, and TX:** Apply *VISOR II XRT* preplant incorporated or preemergence at 1.3 pints (0.93 lb ai) per acre on sandy loams, 1.3 to 1.75 pints (0.93 to 1.25 lb ai) per acre on medium soils, or 1.75 pints (1.25 lb ai) per acre on fine soils.

**Fall Application for Italian Ryegrass Control:** *VISOR II XRT* may be applied for residual control of glyphosate-resistant Italian ryegrass (*Lolium multiflorum*). Apply this product at 1.75 to 2.25 pints (1.25 to 1.57 lb ai) per acre in the fall (September 1 to December 1) after harvest of the previous crop and prior to Italian ryegrass emergence. Use the lower rate of this product for coarse-textured soils and the higher rate for fine-textured soils. A tillage operation may precede the application. **DO NOT** incorporate to a depth greater than 2 to 3 inches if tillage follows the application of this product. For fall applications after emergence of glyphosate-resistant Italian ryegrass, Paraquat can be tank mixed with *VISOR II XRT* to control emerged ryegrass. Refer to the Paraquat label for specific rates, application instructions, and restrictions. Other registered herbicides may be tank mixed with this product for control or improved control of other weeds present at the time of application.

#### Restrictions

- **DO NOT** apply this product to frozen ground.
- If a spring application is made, the combined total amount of this product applied in the fall plus spring must not exceed the maximum seasonal S-metolachlor rate for cotton (3.45 pints (2.45 lb ai) per acre, depending on soil texture).

**Preplant Incorporated (NM, OK, and TX Only):** Apply to the soil and incorporate into the top inch of soil immediately before planting, at planting, or after planting, but before crop or weeds emerge. Use a rolling cultivator or similar implement to uniformly incorporate not more than 1 inch deep. Use a preplant incorporated application if furrow irrigation is used or when a period of dry weather after application is expected. Where furrow irrigation is used, wet the top of the bed for best results. If the crop is to be planted on beds, apply and incorporate after bed formation. Cotton should be planted below the zone of incorporation; i.e., at least 1 inch on fine soils and 1.5 inches on coarse and medium soils. If incorporated prior to planting, use a planter that will result in a minimum of soil disturbance.

For best control of yellow nutsedge and suppression of seedling johnsongrass, apply this product preplant incorporated at the maximum rate for the soil texture, whether applied alone or mixed with Prometryn.

**Preemergence:** Apply to the soil surface at planting or after planting, but before weeds or crop emerge.

**Postemergence:** Apply *VISOR II XRT* broadcast over the top or directed to the soil surface according to the rate restrictions listed below by state. Over-the-top postemergence application may be made not later than 100 days before harvest, and directed-postemergence application may be made not later than 80 days before harvest. Application before weeds emerge or after clean cultivation to remove existing weeds is necessary since this product will not control emerged weeds. *VISOR II XRT* postemergence may be applied over any previous registered herbicide treatment. In sprinkler-irrigated areas, sprinkler irrigate after application with 1/2 to 1 inch of water (1/2 inch on coarse-textured soils to 1 inch on fine-textured soils) to incorporate this product. In furrow-irrigated areas, apply this product, incorporate with a rolling cultivator or similar implement that provides uniform shallow incorporation (2 inches or less), and then irrigate. In non-irrigated areas, if at least 1/2 inch of rainfall does not occur within 10 days after application, cultivate with a rolling cultivator or similar implement that provides uniform shallow incorporation of this product.

**VA, NC, SC, GA, FL, and AL:** Apply this product postemergence at 1.3 to 1.75 pints (0.93 to 1.25 lb ai) per acre.

**TN, AR, KS, MS, MO, and LA:** Apply this product postemergence at 0.65 to 1.75 pints (0.46 to 1.25 lb ai) per acre.

**TX, OK, NM, AZ, CA, and Clay Soils in AR:** Apply this product postemergence at 1.3 to 1.75 pints (0.93 to 1.25 lb ai) per acre before August 1.

**Multiple Applications:** Where weed pressure is heavy, difficult-to-control species are expected, or re-infestation may occur, and a weed control program is used, multiple applications of *VISOR II XRT* are effective when used as part of the weed control program. Apply as a preplant incorporated or preemergence treatment and follow with an application postemergence to cotton before weeds emerge or after clean cultivation to remove existing weeds, since this product will not control emerged weeds. Apply this product postemergence over a previous preplant or preemergence *VISOR II XRT* application as shown in the following table.

**Table 4. Multiple *VISOR II XRT* Applications to Cotton**

States	Preplant Incorporated or Preemergence (pints per acre) + Postemergence (pints per acre)
MS, LA, TN, AR, KS, MO	0.65 to 1.75 (0.46 to 1.25 lb ai) (Preemergence Only) + 0.65 to 1.75 (0.46 to 1.25 lb ai)
TX, OK, NM	1.3 to 1.75 (0.93 to 1.25 lb ai) + 1.3 to 1.75 (0.93 to 1.25 lb ai) before August 1
NC, VA	1.3 to 1.75 (0.93 to 1.25 lb ai) (Preemergence Only) + 1.3 to 1.75 (0.93 to 1.25 lb ai)

In sprinkler-irrigated areas, sprinkler irrigate after application with 1/2 to 1 inch of water (1/2 inch on coarse-textured soils to 1 inch on fine-textured soils) to incorporate *VISOR II XRT*. In furrow-irrigated areas, apply this product, incorporate with a rolling cultivator or similar implement that provides uniform shallow incorporation (2 inches or less), and then irrigate. In nonirrigated areas, if at least 1/2 inch of rainfall does not occur within 10 days after application, cultivate with a rolling cultivator or similar implement that provides uniform shallow incorporation of this product.

For best control of yellow nutsedge and suppression of seedling johnsongrass, apply *VISOR II XRT* preplant incorporated, preemergence, or postemergence to cotton and preemergence to weeds at the maximum rate for the soil texture, whether applied alone or in combinations. These treatments may be applied over previous registered herbicide treatments.

#### Precautions

- To avoid concentration in the seed furrow, **DO NOT** make broadcast applications of this product to cotton planted in furrows more than 2 inches deep. Band applications may be made to cotton planted in furrows deeper than 2 inches, but band width should not exceed the width of the bottom of the furrow.
- In furrow-planted cotton, to avoid concentration in the furrow and potential injury, **DO NOT** apply this product postemergence until after first "knifing" or cultivation to level soil surface.
- **DO NOT** apply over the top in fluid fertilizer or any other adjuvant, surfactant, oil, or other pesticide not specified in the cotton section of this label, or injury may occur.

#### Restrictions

- **DO NOT** apply more than a total of 2.7 lb ai (1.9 lb ai) of this product per acre on coarse soils or 3.45 pints (2.45 lb ai) of this product per acre on medium and fine soils during a growing season.
- **DO NOT** apply this product on sand or loamy sand soils, or in areas where water is likely to "pond" over the bed.
- **DO NOT** graze or feed forage or fodder from cotton to livestock.
- **DO NOT** apply on Taloka silt loam.
- **DO NOT** use in Gaines County, TX.

#### COTTON - *VISOR II XRT* COMBINATIONS

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

#### Tank Mixture with Prometryn

*VISOR II XRT* tank mixtures with Prometryn may be applied preplant incorporated or preemergence in water or fluid fertilizer. When fluid fertilizer is used as a carrier for this product, either alone or in combination with Prometryn, mix only the amount that will be sprayed in one operation. These mixtures should not be allowed to stand without agitation. Only use water as a carrier for postemergence -directed application.

In addition to those weeds controlled by *VISOR II XRT* alone, *VISOR II XRT* + Prometryn, applied preplant incorporated or preemergence, also controls the following weeds: junglerice, wild oats, annual morningglory, groundcherry, hairy nightshade, lambquarters, malva, mustard, prickly sida (teaweed), purslane, ragweed, and shallow-germinating seedlings of

cocklebur and coffeeweed. As a postemergence-directed application, Prometryn provides postemergence control and residual control of weeds on its label, while this product provides residual control of weed species on its label. **VISOR II XRT** will not control emerged weeds.

**Preplant Incorporated or Preemergence:** Apply **VISOR II XRT** + labeled rate of Prometryn, either preplant incorporated or preemergence using the appropriate rate from **Table 5**. Plant cotton below the zone of incorporation, i.e., at least 1.0 inch on fine soils and 1.5 inches on coarse and medium soils. If incorporated before planting, use a planter that will result in a minimum of soil disturbance.

**Table 5: VISOR II XRT + Prometryn - Cotton (NM, OK, TX)**

Use Areas	Soil Texture	Broadcast Rates Per Acre	
		VISOR II XRT	Prometryn
All	Sand, loamy sand	<b>DO NOT USE</b>	
OK, and Blacklands and Gulf Coast of TX	Loams	1.1 to 1.75 pints (0.78 to 1.25 lb ai)	See Prometryn label of product being used for application rates
	Clays	1.75 pints (1.25 lb ai)	
Rio Grande Valley of TX	Loams	1.1 to 1.75 pints (0.78 to 1.25 lb ai)	
	Clays	1.75 pints (1.25 lb ai)	
NM; High Plains, Rolling Plains, Edwards Plateau of TX; and Southwest TX	Sandy loam	1.1 to 1.3 (0.78 to 0.93 lb ai)	
	Loams	1.1 to 1.75 pints (0.78 to 1.25 lb ai)	
	Sandy clay loams Other clay soils	1.75 pints (1.25 lb ai)	

**Postemergence Directed (AR, AZ, CA, LA, MS, NM, OK, TN, TX, and MO):** **VISOR II XRT** may be tank mixed with Prometryn in water and applied postemergence directed in cotton for control of emerged weeds listed on the Prometryn label and residual preemergence control of weeds controlled by **VISOR II XRT** and Prometryn, or application may be made after cultivation for residual preemergence control. These treatments may be applied over previous registered treatments, including **VISOR II XRT**, provided the maximum label rate of any product is not exceeded.

Apply **VISOR II XRT** + Prometryn in a minimum of 20 gallons of spray volume per acre. Follow the directions, limitations, and precautions on the Prometryn label when Prometryn is applied as a postemergence-directed application. Refer to the directions, limitations, and precautions for use of this product under the **Cotton - VISOR II XRT Alone - Postemergence** section.

#### Precautions

- Do avoid concentration in the seed furrow, **DO NOT** make broadcast applications of this product + Prometryn to cotton planted in furrows more than 2 inches deep. Band applications may be made to cotton planted in furrows deeper than 2 inches, but band width should not exceed the width of the bottom of the furrow.
- **DO NOT** apply over the top of cotton or injury may occur.

#### Restrictions:

- **DO NOT** apply on sand or loamy sand soils or in areas where water is likely to "pond" over the bed.
- **DO NOT** apply in cut areas of newly leveled fields, or in areas of excess salt.
- **DO NOT** graze or feed forage or fodder from cotton to livestock.
- **DO NOT** apply to glandless cotton varieties.
- **DO NOT** apply on Taloka silt loam.
- **DO NOT** use in Gaines County, TX

Refer to the Prometryn label for further instructions and limitations.

#### Tank Mixture with Fluometuron

Apply **VISOR II XRT** in tank mixture with Fluometuron preemergence for control of those weeds controlled by this product alone and those as listed on the Fluometuron label. This combination will also control spotted spurge, hyssop spurge, nodding spurge, and prostrate spurge. Apply to the soil surface at planting or after planting, but before weeds or crops emerge, using the appropriate rates from **Table 6**. Apply the tank mixture postemergence to cotton but preemergence to weeds, or apply postemergence to both cotton and broad leaf weeds for control of weeds on the Fluometuron label. Apply as a directed, semi-directed, or over-the-top spray. **VISOR II XRT** will not control emerged weeds but will provide preemergence control of species on its label. Where rate ranges are given for Fluometuron, use the higher rate when applying postemergence to weeds that are 2 inches or less. These treatments may be applied over previous registered treatments, including this product, provided the maximum label rate of any product is not exceeded.

**Mixing Instructions:** Incompatibility may occur when tank mixing **VISOR II XRT** and Fluometuron. To help overcome this condition, fill the spray tank 1/4 full with water or fluid fertilizer and start agitation, add the Fluometuron and allow it to become dispersed. Add Voyager 90/10 at 0.5% volume/volume final spray (4.0 pints per 100 gallons), then add this product and finally the rest of the water or fluid fertilizer. Agitate during mixing and application to maintain a uniform suspension. **DO NOT** use fluid fertilizer as a carrier for postemergence applications.

**Table 6: VISOR II XRT + Fluometuron - Cotton**

Soil Texture	Broadcast Rates Per Acre		
	VISOR II XRT (pints)		Fluometruon
	Area 1*	Area 2**	
Sand, loamy sand	DO NOT USE		
Sandy loam	0.65 to 1.3 (0.46 to 0.93 lb ai)	1.1 to 1.3 (0.78 to 0.93 lb ai)	See Fluometuron label of product being used for application rates
Loam, silt loam, silt	0.9 to 1.75 (0.64 to 1.25 lb ai)	1.3 to 1.75 (0.93 to 1.25 lb ai)	
Fine soil	1.3 to 1.75 (0.93 to 1.25 lb ai)	1.75 (1.25 lb ai)	
* Area 1 = AR, LA, MS, Bootheel of MO and TN			
** Area 2 = Eastern OK, Gulf Coast, Rio Grande Valley, and Eastern TX			

#### Precautions

- **DO NOT** apply this product + Fluometuron on sand or loamy sand soils, or in areas where water is likely to "pond" over the bed, or crop injury may occur.
- To avoid concentration in the seed furrow, **DO NOT** make broadcast applications of this product + Fluometuron to cotton planted in furrows more than 2 inches deep. Band applications may be made to cotton planted in furrows deeper than 2 inches, but band width should not exceed the width of the bottom of the furrow.
- The use of Fluometuron following the use of a systemic insecticide at planting may result in crop injury.

#### Restrictions

- **DO NOT** feed treated forage or gin trash to livestock, or graze treated areas.
- **DO NOT** apply on Taloka silt loam.
- **DO NOT** use in Gaines County, TX.

Refer to the Fluometuron labels for further instructions, precautions, and limitations.

#### TANK MIXTURE OF VISOR II XRT OR VISOR II XRT + FLUOMETURON WITH PARAQUAT OR GLYPHOSATE FOR MINIMUM-TILLAGE OR NO-TILLAGE SYSTEMS

In minimum-tillage or no-tillage systems where cotton is planted directly into a cover crop, stale seedbed, or previous crop residues, the contact herbicides Paraquat or Glyphosate may be added to a tank mix of either **VISOR II XRT** or **VISOR II XRT** + Fluometuron. When used as directed, the Paraquat portion of the tank mixture controls most emerged weeds and suppresses many perennial weeds. Glyphosate combinations will control emerged annual and perennial weeds when applied as directed on the Glyphosate label. The **VISOR II XRT** and **VISOR II XRT** + Fluometuron portion of the tank mixture provides preemergence control of the weeds listed on this label in the **VISOR II XRT** and **VISOR II XRT** + Fluometuron sections, respectively.

Refer to the label of each product used in combination and observe the planting details, information regarding application, geographical restrictions, and all other precautions and limitations. Refer to **Mixing Instructions** under **Tank Mixture with Fluometuron** section.

**Application:** Apply before, during, or after planting, but before the cotton emerges. Apply *VISOR II XRT* at 1.1 to 1.3 (0.76 to 0.93 lb ai) per acre on sandy loams, medium-, and fine-textured soils. Refer to Table 6 for Fluometuron rates.

**Paraquat:** Apply as directed on the product label. This treatment will not control weeds taller than 6 inches.

**Glyphosate:** See the Glyphosate label for weeds controlled, specified rates, and other use directions.

Apply in 20 to 60 gallons of water or fluid fertilizer per acre with ground equipment.

#### Precautions

- If heavy rain occurs soon after application, crop injury may result, especially in poorly drained areas where water stands for several days, or where the seeding slit has not been properly closed.
- Refer to the Fluometuron labels and the **Tank Mixture with Fluometuron** section of this label for further instructions, precautions, and limitations.

#### Restriction

- **DO NOT** apply combinations containing Paraquat in suspension-type liquid fertilizers, as the activity of paraquat will be reduced.
- **DO NOT** apply this product + Fluometuron + Glyphosate in tank mixture because of compatibility problems.
- **DO NOT** use in Gaines County, TX

**TANK MIXTURE WITH MSMA, MSMA + FLUOMETURON, OR MSMA + FLUOMETURON**  
*VISOR II XRT* may be tank mixed with MSMA in water and applied postemergence directed for control of emerged weeds listed on the MSMA product label and residual preemergence control of weeds controlled by this product. The addition of Fluometuron or Prometryn will add control of weed species on their respective labels.

**Postemergence Directed (AL, AR, AZ, CA, FL, GA, LA, MS, NC, NM, OK, SC, TN, TX, VA, and Bootheel of MO):** Apply *VISOR II XRT* + MSMA postemergence directed to cotton at least 3 inches tall according to the directions, limitations, and precautions on the MSMA product label, as well as the directions, limitations, and precautions for use of this product in the section for **Cotton - *VISOR II XRT* Alone - Postemergence**. These treatments may be applied over previous registered treatments, including *VISOR II XRT*, provided the maximum label rate of any product is not exceeded. Fluometuron or Prometryn may be added to the *VISOR II XRT* + MSMA tank mixture according to the respective label directions for application to cotton at least 3 inches tall. When these mixtures are used, follow the mixing instructions for *VISOR II XRT* + Fluometuron or Prometryn and then add the MSMA product.

#### Restrictions

- **DO NOT** apply after first cotton bloom.
- **DO NOT** use this product in tank mix with premixes of MSMA plus herbicides other than those registered for use in tank mixture with this product on cotton.

#### TANK MIXTURE WITH TRIFLURALIN FOR POST-DIRECTED FOLLOWED BY SOIL INCORPORATION APPLICATIONS

*VISOR II XRT* may be applied as a tank mixture with Trifluralin in cotton for improved late-season weed control when used as an incorporated lay-by-type application. This combination may be applied after the cotton is at least 3 inches tall and has reached the 4 true-leaf stage. Make the application directed to the soil surface and away from the crop foliage. Incorporate using a sweep or rolling type cultivator to provide uniform and shallow mixing into the top 2 inches of soil. Refer to each product label for the appropriate application rates by soil type and for this application timing, and follow all product use limitations and restrictions.

#### TANK MIXTURE WITH GLYPHOSATE FOR USE ON ROUNDUP READY OR GLYPHOSATE-RESISTANT COTTON ONLY

Apply *VISOR II XRT* as a tank mixture with Glyphosate in water postemergence over-the-top or postemergence directed for control of emerged weeds listed on the Glyphosate label and for residual preemergence control of weeds listed on the label of this product. See the **Cotton - *VISOR II XRT* Alone Postemergence** section of this label for rates and timings of *VISOR II XRT* and follow the Glyphosate label for their respective rates, application methods, and application timing restrictions. Refer to the Glyphosate label and follow appropriate use

directions, application procedures, precautions, and limitations.

#### Precautions

- Postemergence over-the-top applications of this tank mixture may cause temporary injury in the form of necrotic spotting to exposed cotton leaves, which will not affect normal plant development.
- **DO NOT** add additional spray adjuvants, surfactants, fertilizer additives, or pesticides to this tank mixture if applied postemergence over-the top or unacceptable injury may occur.

#### Restrictions:

- **DO NOT** apply Glyphosate postemergence over-the-top to cotton past the growth stage limit specified on their respective labels.
- **DO NOT** apply this tank mixture postemergence to any cotton variety unless it is designated Roundup Ready or glyphosate-resistant and unless the Glyphosate formulation being used is registered for postemergence use in Roundup Ready or glyphosate-resistant Cotton.
- **DO NOT** use on sand or loamy sand soils in Gaines County, TX.

#### SOYBEAN, IMMATURE SEED

*VISOR II XRT* may be applied preplant or preemergence for the control or suppression of grass and small-seeded weeds in immature-seed soybean or other foodgrade soybeans. For specific rates, see the rate table listed below.

**Preplant Surface-Applied:** For minimum-tillage or no-tillage systems only, *VISOR II XRT* alone may be applied up to 45 days before planting. Use only split applications for treatments made 30 to 45 days before planting, with 2/3 the specified broadcast rate for the crop and soil texture applied initially and the remaining 1/3 applied at planting. Treatments less than 30 days before planting may be made either as a split or a single application. If weeds are present at the time of treatment, apply in a tank mixture combination with a contact herbicide (e.g., Paraquat or Glyphosate). Observe directions for use, precautions, and restrictions on the label of the contact herbicide. To the extent possible, **DO NOT** move treated soil out of the row or move untreated soil to the surface during planting, or weed control will be diminished.

**Preplant Incorporated:** Apply *VISOR II XRT* to the soil and incorporate into the top 2 inches of soil within 14 days before planting, using a finishing disk, harrow, rolling cultivator, or similar implement capable of providing uniform 2-inch incorporation. Use a preplant incorporated application if furrow irrigation is used or when a period of dry weather after application is expected. If crop will be planted on beds, apply and incorporate this product after bed formation, unless specified otherwise.

**Preemergence:** Apply *VISOR II XRT* during planting (behind the planter) or after planting, but before weeds emerge.

#### *VISOR II XRT* Broadcast Rates Per Acre

Soil Texture	Percent Organic Matter in Soil	
	< 3%	≥ 3%
Coarse	1.3 to 1.75 pints (0.93 to 1.25 lb ai)	1.75 pints (1.25 lb ai)
Medium	1.75 to 2.2 pints (1.25 to 1.57 lb ai)	1.75 to 2.2 pints (1.25 to 1.57 lb ai)
Fine		2.2 to 2.7 pints (1.57 to 1.9 lb ai)

#### Precaution

- This product will not control emerged weeds.

#### Restrictions

- **DO NOT** cut for hay within 120 days following an application of this product.
- **DO NOT** use for forage within 60 days following an application of this product.
- **DO NOT** apply more than 2.7 lb ai (1.9 lb ai) of this product per acre during anyone crop year.

#### GRASSES GROWN FOR SEED (ID, OR, WA) - *VISOR II XRT* APPLIED ALONE

To control weeds and volunteer grasses in established grasses grown for seed, apply *VISOR II XRT* to established stands of tall fescue, orchardgrass, perennial ryegrass, fine fescue,



bentgrass, and Kentucky bluegrass just before, during, or immediately following the first fall rains or just before or during a late summer or early fall irrigation, but before target grasses emerge. The seed crop must have had one seed harvest or been established at least one year. The post-harvest residue (straw) should be evenly spread, removed, or burned before applying this product. Rainfall or irrigation is required after application and before weed emergence for best control. This product will provide preemergence control/suppression of volunteer seedlings of perennial ryegrass, fine fescue species, tall fescue, orchardgrass, bentgrass, and Kentucky bluegrass. This product will control those weed species listed in the **VISOR II XRT Alone** section of this label and will suppress or control rattle fescue, annual bluegrass, Italian ryegrass, California brome, downy brome, and roughstalk bluegrass.

Apply **VISOR II XRT** by ground equipment in a minimum of 10 gallons of water per acre using the rate listed below according to grass species.

Established Grass Crop Grown for Seed	Pints per Acre
Fine fescue species	1.3
Perennial ryegrass	(0.93 lb ai)
Bentgrass	
Kentucky bluegrass	1.3 to 1.75
Orchardgrass	(0.93 to 1.25 lb ai)
Tall fescue	

#### Precautions

- **DO NOT** apply after the November 15 or poor control may result.
- Tank mixtures with other pesticides, or the addition of an adjuvant, can increase the risk of crop injury.
- Application to perennial ryegrass and fine fescue stands under stress may cause crop injury.
- If weed escapes occur following an application of this product, an application of a postemergence herbicide may be necessary to control escapes. When making such an application, follow all directions, precautions, and limitations on the label of the postemergence herbicide.
- Control may be decreased if excessive straw from the previous harvest is present at application and/or insufficient rainfall/irrigation occurs.

#### Restrictions

- Apply this product only once per crop year.
- **DO NOT** graze forage regrowth for 60 days following application west of the Cascades.
- In areas east of the Cascades, **DO NOT** graze forage regrowth for 150 days following application.
- Hay may be harvested anytime between seed harvest and the next application of S-metolachlor.

#### HORSERADISH

Apply a single application of **VISOR II XRT** at a broadcast rate of 1.3 to 1.75 pints (0.93 to 1.25 lb ai) per acre to the soil surface after planting, but before weed or crop emergence (i.e., preemergence). Use lower rates on soils relatively coarse-textured and higher rates on fine-textured soils. A band application may also be used, applying proportionally less spray mixture on the area actually treated. This product will not control emerged weeds. Control emerged weeds with an appropriate registered foliar herbicide or by mechanical means. Harvest horseradish at normal timing.

#### Restrictions

- Make only one application of this product per crop.
- **DO NOT** apply more than 1.75 pints (1.25 lb ai) of this product per acre per crop.

#### PEANUTS - VISOR II XRT ALONE

Apply **VISOR II XRT** either preplant incorporated, postplant incorporated, or preemergence, using the appropriate rate specified below.

**Preplant Incorporated or Preemergence:** Follow instructions for use of **VISOR II XRT Alone** under **Application Procedures**.

**Postplant Incorporated:** Apply and shallowly incorporate **VISOR II XRT** into the soil after planting, but before peanut germination. Incorporation depth and incorporating implements must be kept above the seed, or seed will be damaged.

Apply **VISOR II XRT** alone, preplant incorporated, postplant incorporated, or preemergence, at a broadcast rate of 1.3 to 1.75 pints (0.93 to 1.25 lb ai) per acre in the Southeast\* and 1.1 to 1.75 pints (0.78 to 1.25 lb ai) per acre in NM, OK, and TX.

\* In the Southeast, use 1.75 to 2.7 pints (1.25 to 1.9 lb ai) per acre and apply preemergence for partial control of Florida beggarweed.

This product alone may be applied as directed after any of the following preplant incorporated herbicides when used according to their label: Benfluralin, Ethafluralin, Imazethapyr, Pendimethalin or Trifluralin.

#### Restrictions

- **DO NOT** graze or feed peanut forage or fodder to livestock for 30 days following application.
- **Preharvest Interval (PHI): DO NOT** apply within 90 days of harvest.

#### PEANUTS - VISOR II XRT COMBINATIONS

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

#### Tank Mixture with Benfluralin

**VISOR II XRT** + Benfluralin tank mixture applied preplant incorporated controls those weeds listed under **VISOR II XRT Applied Alone** and those weeds as listed on the Benfluralin label.

Apply 1.3 to 1.75 pints (0.93 to 1.25 lb ai) per acre of **VISOR II XRT** + labeled rate of Benfluralin in a minimum of 10 gallons of spray volume per acre for ground application or in a minimum of 5.0 gallons of spray volume per acre for aerial application. Follow the procedures for Benfluralin on the Benfluralin label for soil preparation and incorporation of this tank mix. Apply and incorporate **VISOR II XRT** + Benfluralin up to 14 days prior to planting. Follow all restrictions and precautions on the Benfluralin label.

#### Tank Mixture or Sequentially with Imazethapyr

The tank mixture or sequential treatment of **VISOR II XRT** and Imazethapyr controls all weeds controlled by this product alone and by Imazethapyr alone. Refer to the **VISOR II XRT Applied Alone** section for weeds controlled by this product and to the Imazethapyr label for weeds controlled by Imazethapyr.

Refer to the respective labels for application methods, timing, rates, restrictions, and precautions; and use in accordance with the more restrictive label. **DO NOT** exceed the label rate of either product. **VISOR II XRT** will not control emerged weeds.

#### Tank Mixture with Ethafluralin

The tank mixture controls all weeds controlled by **VISOR II XRT** alone and by Ethafluralin alone. Refer to the **VISOR II XRT Applied Alone** section for weeds controlled by this product and to the Ethafluralin label for weeds controlled by Ethafluralin.

Apply **VISOR II XRT** + Ethafluralin preplant incorporated, using the appropriate rate from **Table 7**. Follow specified soil preparation procedures for Ethafluralin.

**Table 7: VISOR II XRT + Ethafluralin – Peanuts**

Broadcast Rates Per Acre				
Soil Texture	Southeast		NM, OK, TX	
	VISOR II XRT	Ethafuralin	VISOR II XRT	Ethafuralin
Coarse	1.3 to 1.75 pints (0.93 to 1.25 lb ai)	See Ethafuralin label of product being used for application rates	1.1 to 1.75 pints (0.78 to 1.25 lb ai)	See Ethafuralin label of product being used for application rates
Medium				
Fine				
Follow all use directions, limitations, use precautions and information regarding application to Peanuts on respective product labels.				

#### Tank Mixture with Pendimethalin

**VISOR II XRT** + Pendimethalin applied preplant incorporated controls all weeds controlled by this product alone plus Texas panicum, field sand bur, johnsongrass from seed,

lambsquarters, kochia, annual spurge, and other species on the Pendimethalin label. Apply **VISOR II XRT** + Pendimethalin by ground or by aerial equipment within 14 days before planting. Incorporate into the top 1 to 2 inches of soil before planting and within 7 days of application, using a finishing disk or similar implement capable of providing uniform incorporation. If peanuts will be planted on beds, apply and incorporate after bed formation. Refer to the **Incorporation** instructions of the respective labels for additional directions.

Apply **VISOR II XRT** + Pendimethalin preplant incorporated, using the appropriate rates from Table 8.

**Table 8: *VISOR II XRT* + Pendimethalin – Peanuts**

Soil Texture	Broadcast Rates Per Acre			
	NM, OK, TX		Other Peanut Growing States	
	<i>VISOR II XRT</i>	Pendimethalin	<i>VISOR II XRT</i>	Pendimethalin
Sand, Loamy sand	1.1 (0.78 lb ai)	See Pendimethalin label of product being used for application rates	1.3 to 1.75 pints (0.93 to 1.25 lb ai)	See Pendimethalin label of product being used for application rates
Sandy loam	1.1 to 1.3 (0.78 to 0.93 lb ai)			
Fine soil	1.75 pints (1.25 lb ai)		1.75 pint (1.25 lb ai)	

Follow all use directions, limitations, use precautions and information regarding application to Peanuts on respective product labels.

#### Tank Mixture with Paraquat

**VISOR II XRT** + Paraquat applied at ground cracking will control or suppress small (1 to 6 inch) emerged annual grass and broadleaf weeds and provide residual control of weed species listed in the **VISOR II XRT Applied Alone** section of this label. Apply Paraquat plus the appropriate **VISOR II XRT** rate from the **Peanuts - *VISOR II XRT Alone*** section in a minimum spray volume of 20 gallons per acre with ground equipment. Refer to the Paraquat label and follow all directions, limitations, and restrictions.

#### Tank Mixture or Sequentially with Paraquat + Bentazon

The addition of Bentazon to the **VISOR II XRT** + Paraquat mixture will result in improved control of such problem broadleaf weeds as prickly sida, cocklebur, smartweed, and bristly starbur. **VISOR II XRT** + Paraquat + Bentazon applied at ground cracking will control or suppress small (1 to 6 inch) emerged annual grass and broadleaf weeds and provide residual control of weed species listed in the **VISOR II XRT Applied Alone** section of this label. Apply Bentazon + Paraquat with the appropriate **VISOR II XRT** rate from the **Peanuts - *VISOR II XRT Alone*** section in a minimum spray volume of 20-gallons per acre with ground equipment. Refer to the Paraquat and Bentazon labels and follow all directions, limitations, and restrictions.

#### Tank Mixture with Paraquat + 2,4-DB

The addition of 2,4-DB to the **VISOR II XRT** + Paraquat mixture will result in improved control of such problem broadleaf weeds as sicklepod, morningglory, and cocklebur. **VISOR II XRT** + Paraquat + 2,4-DB applied at ground cracking will control or suppress small (1-6 inch) emerged annual grass and broadleaf weeds and provide residual control of weed species listed in the **VISOR II XRT Applied Alone** section of this label. Apply Paraquat + 2,4-DB with the appropriate **VISOR II XRT** rate from the **Peanuts - *VISOR II XRT Alone*** section in a minimum spray volume of 20 gallons per acre with ground equipment. Refer to the Paraquat and 2,4-DB labels and follow all directions, limitations and restrictions.

#### Tank Mixture with Bentazon

**VISOR II XRT** + Bentazon applied at ground cracking will control species on the Bentazon label and provide residual control of species listed in the **VISOR II XRT Applied Alone** section of this label. Apply the labeled rate of Bentazon in 20 gallons per acre, depending on weed species and stage of growth as specified on the Bentazon label, with the appropriate rate of this product from the **Peanuts - *VISOR II XRT Alone*** section. A second Bentazon application may be made in a peanut-growing areas, if needed. Refer to the respective labels and follow all directions, limitations, and restrictions for each product.

#### Tank Mixture with Bentazon + 2,4-DB

**VISOR II XRT** + Bentazon + 2,4-DB applied at ground cracking will control species on the Bentazon and 2,4-DB labels, especially morning glories. Apply labeled rates of Bentazon + 2,4-DB in 20 gallons per acre depending on weed species and stage of growth as specified on the Bentazon label, with the appropriate rate of this product from the **Peanuts - *VISOR II XRT Alone*** section. A second Bentazon + 2,4-DB application may be made in all peanut-growing areas, if needed. Refer to the respective labels and follow all directions, limitations, and restrictions for each product.

#### Sequentially with Bentazon + Acifluorfen

Apply **VISOR II XRT** according to the directions for **VISOR II XRT Alone** and follow with a postemergence treatment of Bentazon + Acifluorfen as specified on its label for the control of weeds listed on this label and on the Bentazon + Acifluorfen label. Refer to the **VISOR II XRT - Peanuts - Alone** section and to the Bentazon + Acifluorfen label and follow all directions, limitations, and restrictions for each product.

#### MULTIPLE APPLICATIONS

Where weed pressure is heavy or where species difficult to control are expected, **VISOR II XRT** is most effective when used as follows:

##### 1. Southeast Only (AL, FL, GA, NC, SC, VA)

**Preplant Incorporated:** Apply **VISOR II XRT** preplant incorporated as directed under **Peanuts - *VISOR II XRT Alone*** or apply **VISOR II XRT** - Benfluralin preplant incorporated as directed previously in this section. Refer to the respective section for weeds controlled. **OR**

**Preemergence to before "ground cracking":** Apply **VISOR II XRT** any time from preemergence to before "ground cracking" at 1.3 to 2.7 pints (0.93 to 1.9 lb ai) per acre for extended control of weeds not yet emerged. **DO NOT use *VISOR II XRT* after peanut emergence.** If peanuts have emerged, use **VISOR S-MOC HERBICIDE** according to its label; **Peanuts - Combinations - Multiple Applications.**

Follow the PPI or PRE application by:

**Lay-by: DO NOT use *VISOR II XRT*.** Apply **VISOR S-MOC HERBICIDE** at lay-by as directed under the **Peanuts - Alone** section of the **VISOR S-MOC HERBICIDE** label.

##### 2. Southwest Only (NM, OK, TX)

**1st Application:** Apply **VISOR II XRT** preplant incorporated or preemergence or before "ground cracking" as directed under **Peanuts - *VISOR II XRT Alone*** or apply **VISOR II XRT** + Bentazon preplant incorporated as directed previously in this section. **DO NOT use *VISOR II XRT* after peanut emergence.** If peanuts have emerged, use **VISOR S-MOC HERBICIDE** according to its label.

**2nd Application: DO NOT use *VISOR II XRT*.** Apply **VISOR S-MOC HERBICIDE** at lay-by as directed under **Peanuts - Alone** section of the **VISOR S-MOC HERBICIDE** label. Use only when late germinating weeds are expected to be a problem. Refer to the product **Applied Alone** section for a list of weeds controlled.

#### Restrictions for Multiple Applications

- **DO NOT** apply more than the equivalent of 2.67 lbs. S-metolachlor per acre during any one year. If **VISOR S-MOC HERBICIDE** is used as a sequential treatment, the pounds of active ingredient for all S-metolachlor applications must not exceed 2.67 lbs a.i. per acre.
- **DO NOT use this product or *VISOR S-MOC HERBICIDE* after peanuts have emerged.**
- **DO NOT** graze or feed peanut forage or fodder to livestock for 30 days following application.
- **Preharvest Interval (PHI): DO NOT** apply within 90 days of harvest.

#### BEANS, PEAS, AND LENTILS - *VISOR II XRT ALONE*

Beans, peas, and lentils, including garbanzo, great northern beans, kidney beans, lima beans, mung beans, navy beans, peas (English), southern peas, such as blackeye, pinkeye, crowder, etc.), pinto beans, snap beans (green, wax, string), lentils, and lupines (sweet, white, white sweet, and grain).

#### Fall Application:

1. Apply after September 30 in ND, SD, MN, WI, and north of Route 30 in IA.
2. Apply after October 15 north of Route 91 in NE and south of Route 30 in IA.
3. Apply after October 31 north of Route 136 in IL.

In all locations, apply to crop stubble after harvest when the sustained soil temperature at a 4-inch depth is less than 55°F and falling. In minimum-till or no-tillage systems on soils having greater than 2.5% organic matter, use 2.2 to 2.7 pints (1.57 to 1.9 lb ai) per acre on medium-textured and 2.7 lb ai (1.9 lb ai) per acre on fine-textured soils. A tillage operation may precede the application. A fall and/or a spring tillage may follow application, but **DO NOT** exceed an incorporation depth greater than 2 to 3 inches. Minimize furrow and ridge formation in the tillage operations.

#### Restrictions

- If a spring application is made, the total rate of the fall plus spring applications must not exceed the maximum total rate for beans, peas, and lentils.
- **DO NOT** apply to frozen ground.

#### Spring Application:

Apply **VISOR II XRT**, either preplant incorporated or preemergence, using the appropriate rate specified below. **Preplant Incorporated or Preemergence:** Follow instructions for use of this product alone under **Application Procedures**. On coarse soils with less than 3% organic matter, apply 1.3 to 1.75 pints (0.93 to 1.25 lb ai) of this product per acre or 1.75 pints (1.25 lb ai) per acre if organic matter is 3% or greater. On medium soils, apply 1.75 to 2.2 pints (1.25 to 1.57 lb ai) of this product per acre. On fine soils, apply 1.75 to 2.2 pints (1.25 to 1.57 lb ai) of this product per acre if organic matter content is less than 3%, or 2.2 to 2.7 pints (1.57 to 1.9 lb ai) per acre if organic matter content is 3% or greater.

\*On English peas, use only preemergence applications. If soils are cold and wet during pea germination and emergence, the use of this product may delay maturity and/or reduce yields.

#### Restrictions

- **DO NOT** cut for hay within 120 days following an application of this product.
- **DO NOT** use for forage within 60 days following an application of this product.
- **DO NOT** apply more than 2.7 lb ai (1.9 lb ai) of this product per acre during any one crop year.

#### BEANS, PEAS, AND LENTILS - **VISOR II XRT** COMBINATIONS

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

#### Restriction

- When applying this product in combination on beans, peas, and lentils, **DO NOT** cut for hay within 120 days following application.

#### Tank Mixture and Sequential Applications with EPTC- Beans (Green or Dry)

This mixture controls all weeds controlled by **VISOR II XRT** alone and by EPTC alone. Refer to the **VISOR II XRT Applied Alone** section of this label for weeds controlled by this product alone and to the EPTC label for weeds controlled by EPTC.

**Preplant Incorporated:** Follow instructions for use of **VISOR II XRT** alone under **Application Procedures**.

**Sequential:** Apply EPTC alone preplant incorporated, as specified on that label. Follow with a preemergence application of **VISOR II XRT**, at rates specified for this product alone, during planting (behind the planter) or after planting, but before the weeds or crop emerge.

Refer to the **Product Information** section of this label and to the EPTC label for weather, cultural practices, and all other precautions and limitations that affect performance of these products.

Apply labeled rate of EPTC\* with **VISOR II XRT** as specified below:

On coarse soils, apply 1.1 pints (0.78 lb ai) of this product per acre if organic matter content is less than 3%, or 1.3 pints (0.93 lb ai) per acre if organic matter content is 3% or greater. On medium soils, apply 1.3 pints (0.93 lb ai) of this product per acre if organic matter content is less than 3%, or 1.75 pints (1.25 lb ai) per acre if organic matter content is 3% or greater. On fine soils, apply 1.75 pints (1.25 lb ai) of this product per acre if organic matter content is less than 3%, or 1.75 to 2.2 pints (1.25 to 1.57 lb ai) per acre if organic matter is 3% or greater.

\*Refer to the EPTC label for rate limitations depending on geographical area, and for species and varietal restrictions.

#### Restriction

- **DO NOT** exceed 3 lbs. a.i. per acre of EPTC on small white beans or green beans grown on coarse-textured soils.

#### Tank Mixture with Trifluralin - Beans (Dry - Kidney, Navy, Pinto, Etc.; Lima; And Snap)

**VISOR II XRT** + Trifluralin tank mix applied preplant incorporated controls those weeds listed under **VISOR II XRT Applied Alone** and those weeds listed for Trifluralin alone on the Trifluralin label. **VISOR II XRT** + Trifluralin may be applied by ground or by aerial equipment and incorporated up to 14 days prior to planting. Follow the procedures on this label and on the respective Trifluralin label using equipment that provides uniform 2-inch incorporation.

Apply **VISOR II XRT** + Trifluralin tank mix using the appropriate **VISOR II XRT** rate specified for this product alone, and the Trifluralin rate from the Dry Beans, and the Lima and Snap Beans sections of the respective Trifluralin label. Choose the product rate for the specific soil texture/organic matter classification and weed species expected.

Follow all restrictions and precautions on the respective Trifluralin label and in the **Beans, Peas, and Lentils - **VISOR II XRT** Alone** section of this label.

#### POTATOES - **VISOR II XRT** ALONE

Apply **VISOR II XRT**, either incorporated, preemergence, or postemergence to potatoes after hilling/lay-by, according to directions specified below for control of weeds listed under the **Product Information** section. Within a rate range, use the lower rate on soils relatively coarse-textured or low in organic matter; use the higher rate on soils relatively fine-textured or high in organic matter. Effectiveness will be reduced if later cultural practices expose untreated soil. For application by center pivot irrigation, see the **Center Pivot Irrigation Application** section of this label.

**Incorporated:** Apply **VISOR II XRT** at 1.3 to 2.7 pints (0.93 to 1.9 lb ai) per acre to the soil and incorporate into the top 3 inches before planting, using a finishing sick, harrow, rolling cultivator, or similar implement. Planting and later cultural practices should not bring untreated soil to the surface. Postplant incorporated application may be made any time after planting to drag-off, but before potato emergence. Use an implement that evenly distributes this product in the top 2 inches of soil. **DO NOT** damage potato seed pieces or sprouts with incorporation equipment.

**Preemergence:** Apply **VISOR II XRT** at 1.3 to 2.7 pints (0.93 to 1.9 lb ai) per acre, either after planting as a preemergence, delayed preemergence, after drag-off or hilling treatment, but before weeds emerge. Up to 3.3 pints (2.35 lb ai) per acre of this product alone may be used where soil organic matter is between 6% and 20%.

**Postemergence After Hilling/Lay-by:** Apply 2.2 pints (1.57 lb ai) per acre of **VISOR II XRT** postemergence to potatoes through after hilling/at lay-by to control **VISOR II XRT** sensitive species for remainder of the growing season. This application will not control emerged weeds. It may be applied over a previous **VISOR II XRT** application, but **DO NOT** apply more than 4.85 pints (3.45 lb ai) per acre of this product in a single crop season.

#### Precaution

- If cool, wet soil conditions occur after application, this product may delay maturity and/or reduce yield of Superior and other early maturing potato varieties.
- These directions for use **DO NOT** apply to sweet potatoes or yams.

#### Restrictions

- **DO NOT** use on muck or peat soils.
- **DO NOT** apply both as a preemergence and an incorporated treatment.
- **Preharvest Interval (PHI):** **DO NOT** harvest potatoes within 60 days after the at-planting to drag-off application.

#### POTATOES - **VISOR II XRT** COMBINATIONS

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

#### Tank Mixture with Metribuzin

In addition to those weeds controlled by **VISOR II XRT** alone, **VISOR II XRT** applied in tank mix combination with, or sequentially with, any of the registered Metribuzin formulations

also controls the following broad leaf weeds: cocklebur\*, hairy nightshade\*, hemp sesbania, jimsonweed\*, lambsquarters, prickly sida, ragweed, smartweed, velvetleaf, Venice mallow, and wild mustard.

\* Partially controlled.

**VISOR II XRT** at 1.3 to 2.7 pints (0.93 to 1.9 lb ai) per acre plus the labeled Metribuzin use rate may be used preemergence or postemergence to potatoes through after last hilling. Apply 1.3 to 1.75 pints (0.93 to 1.25 lb ai) per acre of this product on coarse soils and 1.75 to 2.7 pints (1.25 to 1.9 lb ai) per acre on other soil textures. Within this rate range, use the lower rate on soils relatively coarse-textured or low in organic matter; use the higher rate on soils relatively fine-textured or high in organic matter. Effectiveness will be reduced if later cultural practices expose untreated soil. This product will not control emerged weeds.

Refer to the Metribuzin label for precautionary statements, restrictions, application information, center pivot irrigation application, weeds controlled, and varietal limitations.

#### Precautions

- Postemergence applications to potatoes, except center pivot, must be made only as a directed or semi-directed spray to avoid chlorosis, minor necrosis, or leaf distortion.
- These directions for use **DO NOT** apply to sweet potatoes or yams.

#### Restrictions

- **DO NOT** use this tank mixture on muck or peat soils.
- **Preharvest Interval (PHI): DO NOT** harvest potatoes treated with this product in a tank mix with Metribuzin within 60 days after application.
- **Preharvest Interval (PHI): DO NOT** harvest potatoes within 40 days after a lay-by application of this product.

#### **VISOR II XRT + Linuron Tank Mixture (East of Rocky Mountains)**

**VISOR II XRT** may be applied in a tank mix combination with any of the registered Linuron formulations as a preemergence broadcast application to potatoes. Apply to the soil surface after planting and before emergence of the crop or after final drag-off according to the rates specified in **Table 9**.

Refer to the **Product Information** section of this label and to the Linuron label for precautionary statements, restrictions, application information, and weeds controlled.

**Table 9: VISOR II XRT + Linuron – Potatoes (East of Rocky Mountains)**

Soil Texture	Broadcast Rates Per Acre			
	1% to Less Than 3% Organic Matter		3 to 5% Organic Matter	
	<b>VISOR II XRT</b>	Linuron	<b>VISOR II XRT</b>	Linuron
<b>Coarse</b> Sandy loam	1.3 pints (0.93 lb ai)	See Linuron label of product being used for application rates	1.75 pints (1.25 lb ai)	See Linuron label of product being used for application rates
<b>Medium</b> Loam, Silt loam, Silt	1.75 pints (1.25 lb ai)		2.2 to 2.7 pints (1.57 to 1.9 lb ai)	

#### Restrictions

- **DO NOT** use on sands or loamy sands
- **DO NOT** incorporate or spray over the top of emerged potatoes.

#### **Tank Mixture with Pendimethalin**

In addition to the weeds controlled by **VISOR II XRT** alone, this tank mixture with Pendimethalin controls such problem species as kochia, lambsquarters, purslane, annual spurge, stinging nettle, and others specified on the Pendimethalin label. Apply **VISOR II XRT** + Pendimethalin preemergence, preemergence incorporated, or early postemergence, according to the specific directions on the Pendimethalin label, using the rates in **Table 10**.

Refer to the label for this product and Pendimethalin and observe all directions, timings, limitations, precautions, and restrictions concerning the use of these products on potatoes and follow the most restrictive

**Table 10: VISOR II XRT + Pendimethalin – Potatoes**

Soil Texture	Broadcast Rates Per Acre			
	Less Than 3% Organic Matter		More Than 3% Organic Matter	
	<b>VISOR II XRT</b>	Pendimethalin	<b>VISOR II XRT</b>	Pendimethalin
<b>Coarse</b>	1.3 to 1.75 pints (0.93 to 1.25 lb ai)	See Pendimethalin label of product being used for application rates	1.3 to 1.75 pints (0.93 to 1.25 lb ai)	See Pendimethalin label of product being used for application rates
<b>Medium</b>	1.75 pints (1.25 lb ai)		1.75 to 2.2 pints (1.25 to 1.57 lb ai)	
<b>Fine</b>	1.75 to 2.2 pints (1.25 to 1.57 lb ai)		2.2 to 2.7 pints (1.57 to 1.9 lb ai)	

#### **Tank Mixture with Pendimethalin + EPTC**

In addition to the weeds controlled by **VISOR II XRT** alone, this tank mixture will control those species on the Pendimethalin and EPTC labels. Refer to the **VISOR II XRT** + Pendimethalin labels for rates of those products and add labeled rate of EPTC per acre depending on geographical area. Refer to the respective labels and observe all directions, limitations, precautions, and restrictions concerning the use of these products on potatoes and follow the most restrictive.

#### **PUMPKIN - VISOR II XRT ALONE**

##### **Preemergence**

Apply **VISOR II XRT** preemergence (before the weeds have emerged) at 1.3 to 1.75 pints (0.93 to 1.25 lb ai) per acre as an inter-row or inter-hill application in pumpkin. Leave 1 foot of untreated area over the row, or 6 inches to each side of the planted hill and/or any emerged pumpkin foliage (inter-row or inter-hill means not directly over the planted seed or young pumpkin plants). Use the lower rate of this product on soils light in texture (loamy sand or lighter) and low in soil organic matter (less than 3%). **VISOR II XRT** applied as a broadcast spray over the planted row or hill, or applications made directly to crop foliage, will increase the risk of injury (e.g., stand loss, delayed maturity, and loss of yield) to the pumpkin crop.

**VISOR II XRT** will not control emerged weeds, and thus should be applied before the weeds emerge. Weeds that are present should be controlled by other means, e.g., by mechanical means or by another herbicide.

#### Restriction

- **Preharvest Interval (PHI): DO NOT** apply within 30 days before pumpkin harvest.

#### **RHUBARB - VISOR II XRT ALONE**

Apply **VISOR II XRT** at a broadcast rate of 0.9 to 1.75 pints (0.64 to 1.25 lb ai) per acre to the soil surface in early spring, prior to crop emergence. Use lower rates on soils relatively coarse-textured and higher rates on fine-textured soils. A band application may also be used, applying proportionally less spray mixture on the area actually treated. This product will not control emerged weeds. Control emerged weeds with an appropriate registered foliar herbicide or by mechanical or physical means.

#### Restrictions:

- Make only one application of this product per crop.
- **DO NOT** apply more than 1.75 pints (1.25 lb ai) of this product per acre per crop.
- **Preharvest Interval (PHI): DO NOT** harvest rhubarb within 62 days of an application of this product.

#### **SAFFLOWERS - VISOR II XRT**

**Preplant Incorporated or Preemergence:** Follow instructions for use of **VISOR II XRT** alone under **Application Procedures**.

On coarse soils, apply 1.3 to 1.75 pints (0.93 to 1.25 lb ai) of this product per acre if organic matter content is less than 3%, or 1.75 pints (1.25 lb ai) per acre if organic matter is 3% or greater. On medium soils, apply 1.75 to 2.2 pints (1.25 to 1.57 lb ai) of this product per acre. On fine soils, apply 1.75 to 2.2 pints (1.25 to 1.57 lb ai) of this product per acre if organic matter content is less than 3%, or 2.2 to 2.7 pints (1.57 to 1.9 lb ai) per acre if organic matter content is 3% or greater.

**SORGHUM (GRAIN OR FORAGE) SEED TREATED WITH A SEED SAFENER (e.g. CONCEPT®) THAT PROVIDES TOLERANCE TO S-METOLACHLOR – VISOR II XRT ALONE**  
Apply **VISOR II XRT**, either preplant surface, preplant incorporated, preemergence, or postemergence using the appropriate rate specified below. Apply this product alone only when the sorghum seed has been properly treated with a seed safener that provides tolerance to S-metolachlor. Preplant or preemergence applications of this product to sorghum not treated with a seed safener that provides tolerance to S-metolachlor will result in crop death.

**Fall Application for Italian Ryegrass Control:** **VISOR II XRT** may be applied for residual control of glyphosate-resistant Italian ryegrass (*Lolium multiflorum*). Apply this product at 1.75 to 2.2 pints (1.25 to 1.57 lb ai) per acre in the fall (September 1 to December 1) after harvest of the previous crop and prior to Italian ryegrass emergence. Use the lower rate of this product for coarse-textured soils and the higher rate for fine-textured soils. A tillage operation may precede the application. **DO NOT** incorporate to a depth greater than 2 to 3 inches if tillage follows the application of this product. For fall applications after emergence of glyphosate-resistant Italian ryegrass, Paraquat can be tank mixed with this product to control emerged ryegrass. Refer to the Paraquat label for specific rates, application instructions, and restrictions. Other registered herbicides may be tank mixed with **VISOR II XRT** for control or improved control of other weeds present at the time of application.

#### Restrictions

- **DO NOT** apply this product to frozen ground.
- If a spring application is made, **DO NOT** apply this product or any other product containing S-metolachlor the following spring to grain or forage sorghum.

**Preplant Surface-Applied:** Refer to instructions for use of **VISOR II XRT** under **Application Procedures** section on this label. For minimum-tillage or no-tillage systems only, apply this product up to 45 days before planting in CO, IA, IL, KS, MO, NE, and SD. Use only split applications for treatments made 30 to 45 days prior to planting, with 2/3 of the broadcast rate applied initially and the remaining 1/3 at planting. Apply 2 pints (1.42 lb ai) per acre of this product on medium soils or 2.2 pints (1.57 lb ai) per acre on fine soils. Treatments less than 30 days prior to planting may be made either as a split or single application. Apply 1.75 pints (1.25 lb ai) of this product per acre on coarse soils not more than 2 weeks prior to planting. Under dry conditions, irrigation after application is needed to move **VISOR II XRT** into the soil.

**Preplant Incorporated or Preemergence:** Refer to instructions for use of **VISOR II XRT** under **Application Procedures** section on this label. Broadcast 1.3 to 1.75 pints (0.93 to 1.25 lb ai) of this product per acre on coarse soils, 1.75 to 2 (1.25 to 1.42 lb ai) per acre on medium soils, or 1.75 to 2.2 pints (1.25 to 1.57 lb ai) per acre on fine soils.

**Postemergence:** Refer to instructions for use of **VISOR II XRT** under **Application Procedures** section on this label. This product may be applied broadcast postemergence at 1.3 to 1.75 pints (0.93 to 1.25 lb ai) per acre on coarse soils, 1.75 to 2 (1.25 to 1.42 lb ai) per acre on medium soils, or 1.75 to 2.2 pints (1.25 to 1.57 lb ai) per acre on fine soils. This product will not control emerged weeds. Therefore, emerged weeds must be controlled by cultural or chemical means. When applied alone, **VISOR II XRT** will be safe to emerged sorghum. The risk of sorghum injury increases when adjuvants (e.g., non-ionic, crop oil), nitrogen sources (e.g., AMS, UAN), or fertilizers are applied with this product.

#### Precautions

- If sorghum seed is not properly treated with a seed safener that provides tolerance to S-metolachlor, preplant and preemergence applications of this product will severely injure the crop.
- Under high soil moisture conditions prior to sorghum emergence, injury may occur following preplant and preemergence application of this product. The crop will normally outgrow this effect.
- Application of this product on sorghum grown under dry mulch tillage, may result in crop injury.

#### Restrictions

- Except for the split preplant surface treatment, **DO NOT** make more than one application per year.
- **Preharvest Interval (PHI):** **DO NOT** apply this product postemergence within 75 days of harvest.

**SORGHUM (GRAIN OR FORAGE) SEED TREATED WITH A SEED SAFENER (e.g. CONCEPT®) THAT PROVIDES TOLERANCE TO S-METOLACHLOR - VISOR II XRT TANK MIXTURES**

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

**VISOR II XRT** preplant or preemergence (prior to sorghum emergence) tank mixtures with Atrazine may be applied in water or fluid fertilizer. Apply this product preplant or preemergence in tank mixtures only when the sorghum seed has been properly treated with a seed safener that provides tolerance to S-metolachlor. Preplant or preemergence applications of **VISOR II XRT** to sorghum not treated with a seed safener that provides tolerance to S-metolachlor will result in crop death.

**Important: For tank mixtures with atrazine –** If applying this product in tank mixture with Atrazine, all the restrictions and rate limitations on the Atrazine label must be followed if more restrictive/protective than those on this label. In addition, if Atrazine is/must be applied at lower rates, broadleaf weed control may be affected. Refer to the Atrazine label for weeds controlled at the reduced rates.

#### Precautions

- Applications of this product + Atrazine on highly alkaline soils or on eroded areas where calcareous subsoils are exposed may cause sorghum injury.
- If sorghum seed is not properly treated with a seed safener that provides tolerance to S-metolachlor, this product + Atrazine may severely injure the crop.
- Under high soil moisture conditions prior to sorghum emergence, injury may occur following preplant and preemergence application of this product + Atrazine. The crop will normally outgrow this effect.
- Use of this product + Atrazine on sorghum grown under dry mulch tillage, may cause crop injury.

#### Restriction

- Except for the split preplant surface treatment, **DO NOT** make more than one application per year.

#### Tank Mixture with Atrazine

In addition to the weeds controlled by **VISOR II XRT** alone, **VISOR II XRT** + Atrazine also controls the following broadleaf weeds when applied either preplant surface, preplant incorporated, or preemergence: cocklebur, common purslane, hairy nightshade, lambsquarters, morningglory, ragweed, smartweed, and velvetleaf.

**Preplant Surface-Applied:** Refer to the instructions for use of this product under the section **Application Procedures**. For minimum-tillage or no-tillage systems only, this product + Atrazine may be applied up to 45 days prior to planting in IA, IL, Eastern KS, MO, NE and SD. Use only split applications for treatments made 30 to 45 days prior to planting, with 2/3 of the broadcast rate applied initially and the remaining 1/3 applied at planting. Apply labeled rate of Atrazine with 2 pints (1.42 lb ai) of this product per acre on medium soils with 1.5% organic matter or greater or with 2 pints (1.42 lb ai) per acre of this product on fine soils with less than 1.5% organic matter or 2.2 pints (1.57 lb ai) of this product per acre on fine soils with 1.5% organic matter or greater. Treatments less than 30 days prior to planting may be made either as a split or single application. Under dry conditions, irrigate after application is needed to move this product + Atrazine into the soil.

#### Precautions

- Use of this product on coarse or medium soils with less than 1.5% organic matter may cause crop injury.

**Preplant Incorporated or Preemergence:** Refer to instructions for use of **VISOR II XRT** under **Application Procedures** on this label. Apply labeled rate of Atrazine with 1.3 pints (0.93 lb ai) of this product per acre on medium soils with 1.5% organic matter or greater, with 1.3 pints (0.93 lb ai) of this product per acre on fine soils with less than 1.5% organic matter or with 1.6 to 1.75 pints (1.14 to 1.25 lb ai) of this product per acre of this product on fine soils with 1.5% organic matter or greater.

## Precautions

To avoid crop injury:

- **DO NOT** use on coarse soils.
- **DO NOT** use on medium soils with less than 1.5% organic matter.
- **DO NOT** use in NM, OK, or TX, except in northeast OK and the TX Gulf Coast and Blacklands areas.
- **DO NOT** apply preplant incorporated in AZ or the Imperial Valley of CA.

## Tank Mixture of VISOR II XRT or VISOR II XRT + Atrazine with Paraquat, Glyphosate + 2,4-D or Glyphosate for Minimum-Tillage or No-Tillage Systems

In minimum-tillage or no-tillage systems where sorghum (seed treated with a seed safener that provides tolerance to S-metolachlor) is planted directly into a cover crop, stale seedbed, established soil, or previous crop residues, the contact herbicides Paraquat, Glyphosate + 2,4-D or Glyphosate may be tank mixed with **VISOR II XRT** or **VISOR II XRT + Atrazine**. See Comment No. 4 following **Table 2**. The **VISOR II XRT** or **VISOR II XRT + Atrazine** portion of the tank mixture provides preemergence control of the weeds listed on this label under the respective sections.

Refer to the label of each product used in combination and observe the planting details, restrictions, and all other precautions and limitations.

**Application:** Apply before, during, or after planting, but before sorghum emerges. Add Paraquat, Glyphosate + 2,4-D or Glyphosate and apply as directed on the product labels.

**Paraquat:** Apply as directed on the Paraquat product label. This treatment will not control weeds taller than 6 inches.

**Glyphosate + 2,4-D:** See the Glyphosate + 2,4-D label for weeds controlled, specified rates for specific weeds and other information concerning use.

**Glyphosate:** See the Glyphosate label for weeds controlled, specified rates and other use directions.

## SWEET SORGHUM (SEED TREATED WITH A SEED SAFENER (e.g. CONCEPT®) THAT PROVIDES TOLERANCE TO S-METOLACHLOR)

Apply **VISOR II XRT** preplant surface, preplant incorporated, preemergence, or postemergence using the appropriate rate specified below. Apply this product only when the sweet sorghum seed has been properly treated with a seed safener that provides tolerance to S-metolachlor (e.g. Concept). Preplant or preemergence applications of this product to sweet sorghum not treated with a seed safener that provides tolerance to S-metolachlor will result in crop death.

**Soil-Applied:** Apply **VISOR II XRT** up to 45 days before planting. Use only split applications for treatments made 30 to 45 days prior to planting, with 2/3 of the broadcast rate applied initially and the remaining 1/3 at planting. Treatments less than 30 days prior to planting may be made either as a split or single application. Under dry conditions, irrigation after application is necessary to move **VISOR II XRT** into the soil.

## VISOR II XRT Rates for Soil Applications to Sweet Sorghum

Soil Type	30 to 45 Days Prior to Planting <sup>1</sup>	<30 Days Prior to Planting	At Planting <sup>2</sup>
Coarse	DO NOT USE	1.75 pints/acre (1.25 lb ai)	1.3 to 1.75 pints/acre (0.93 to 1.25 lb ai)
Medium	2 pints/acre (1.42 lb ai)	2 pints/acre (1.42 lb ai)	1.75 to 2 pints/acre (1.25 to 1.42 lb ai)
Fine	2.2 pints/acre (1.57 lb ai)	2.2 pints/acre (1.57 lb ai)	1.75 to 2.2 pints/acre (1.25 to 1.57 lb ai)

<sup>1</sup> Use only as a split application with 2/3 of the broadcast rate applied initially and the remaining 1/3 applied at planting.

<sup>2</sup> Preplant incorporated or preemergence

**Post-Applied:** **VISOR II XRT** may be applied postemergence to sweet sorghum for residual control of grasses and small-seeded broad leaf weeds. Postemergence application to sweet sorghum may be made to crop up to 5 inches in height. This product will not control emerged weeds. Therefore, emerged weeds must be controlled by cultural or other chemical methods. When applied alone this product will be safe to emerged sweet sorghum. Use of adjuvants is prohibited on sweet sorghum.

## VISOR II XRT Rates for Postemergence Applications to Sweet Sorghum

Soil Type	Postemergence Rate Per Acre
Coarse	1.3 to 1.75 pints (0.93 to 1.25 lb ai)
Medium	1.75 pints (1.25 lb ai)
Fine	

## Precautions

- If sweet sorghum seed is not properly treated with a seed safener that provides tolerance to S-metolachlor (e.g. Concept®), soil applications of this product prior to sorghum emergence will severely injure the crop.
- Under high soil moisture conditions prior to sweet sorghum emergence, injury may occur following soil applications of this product. The crop will normally outgrow this effect.
- To avoid crop injury, **DO NOT** use this product on sorghum grown under dry mulch tillage.

## Restrictions

- Only one application per season is allowed. This product may be applied either as a soil-applied treatment or a postemergence treatment, but not both.
- **Preharvest Interval (PHI): DO NOT** apply this product postemergence within 90 days of harvest.

## SOYBEANS - VISOR II XRT ALONE

Apply **VISOR II XRT** preplant surface-applied, preplant incorporated, preemergence, or postemergence using the appropriate rate specified below. Follow instructions for use of this product alone under **Application Procedures** section of this label.

## Fall Application for Spring Weed Control:

1. Apply after September 30 in ND, SD, MN, WI, and north of Route 30 in IA.
2. Apply after October 15 north of Route 91 in NE and south of Route 30 in IA.
3. Apply after October 31 north of Route 136 in IL.

In all locations, apply to crop stubble after harvest when the sustained soil temperature at a 4-inch depth is less than 55°F and falling. In minimum-till or no-tillage systems on soils having greater than 2.5% organic matter, use 2.2 to 2.7 pints (1.57 to 1.9 lb ai) of this product per acre on medium-textured and 2.7 lb ai (1.9 lb ai) of this product per acre on fine-textured soils. A tillage operation may precede the application. A fall and/or a spring tillage may follow application, but **DO NOT** exceed an incorporation depth greater than 2 to 3 inches. Minimize furrow and ridge formation in the tillage operations.

## Restrictions

- **DO NOT** apply to frozen ground.
- If a spring application is made, the total rate of the fall plus spring applications must not exceed the maximum total rate for soybeans of 3.3 pints (2.35 lb ai) per acre depending on soil texture.

**Fall Application for Italian Ryegrass Control:** **VISOR II XRT** may be applied for residual control of glyphosate-resistant Italian ryegrass (*Lolium multiflorum*). Apply this product at 1.75 to 2.2 pints (1.25 to 1.57 lb ai) per acre in the fall (September 1 to December 1) after harvest of the previous crop and prior to Italian ryegrass emergence. Use the lower rate of this product for coarse-textured soils and the higher rate for fine-textured soils. A tillage operation may precede the application. **DO NOT** incorporate to a depth greater than 2 to 3 inches if tillage follows the application of this product. For fall applications after emergence of glyphosate-resistant Italian ryegrass, Paraquat can be tank mixed with this product for control emerged ryegrass. Refer to the Paraquat label for specific rates, application instructions, and restrictions. Other registered herbicides may be tank mixed with this product for control of improved control of other weeds present at the time of application.

## Restrictions

- **DO NOT** apply this product to frozen ground.
- If a spring application is made, the combined total amount of this product applied in the fall plus the spring must not exceed the maximum seasonal S-metolachlor rate for soybeans 3.3 pints (2.35 lb ai) per acre, depending on soil texture).

**Preplant Surface - Spring Application:** Use on medium and fine soils with minimum tillage or no-tillage systems in CO, CT, DE, IA, IL, IN, KS, KY, MA, MD, ME, MI, MN, MO, MT,

ND, NE, NH, NY, OH, PA, RI, SD, TN, VA, VT, WI, WV, and WY. Apply 2/3 the specified rate of this product (2.2 pints (1.57 lb ai) per acre on medium soils and 2.7 lb ai (1.9 lb ai) per acre on fine soils) as a split treatment 30 to 45 days prior to planting and the remainder at planting. Applications made less than 30 days before planting may be as either a split or single treatment. Apply 1.75 pints (1.25 lb ai) per acre of this product on coarse soils not more than 2 weeks prior to planting.

#### Restrictions

- On soybeans, use up to 3.3 pints (2.35 lb ai) per acre of this product preplant incorporated or preemergence treatment on soils having an organic matter content between 6% and 20%.
- DO NOT** apply more than 3.3 pints (2.35 lb ai) of this product per acre during any one crop season.
- DO NOT** graze or feed treated soybean forage, hay, or straw to livestock 30 days following treatment.

**Preplant Incorporated or Preemergence:** On coarse soils, apply 1.3 to 1.75 pints (0.93 to 1.25 lb ai) of this product per acre if organic matter content is less than 3%, or 1.75 pints (1.25 lb ai) per acre if organic matter content is 3% or greater. On medium soils, apply 1.75 to 2.2 pints (1.25 to 1.57 lb ai) of this product per acre. On fine soils, apply 1.75 to 2.2 pints (1.25 to 1.57 lb ai) of this product per acre if organic matter content is less than 3%, or 2.2 to 2.7 pints (1.57 to 1.9 lb ai) per acre if organic matter content is 3% or greater.

#### Restrictions

- On soybeans, use up to 3.3 pints (2.35 lb ai) per acre of this product preplant incorporated or preemergence treatment on soils having an organic matter content between 6% and 20%.
- The total rate of this product applied to soybeans during any one crop should not exceed 3.3 pints (2.35 lb ai) per acre.
- DO NOT** graze or feed treated soybean forage, hay, or straw to livestock 30 days following treatment.

**Postemergence:** Apply 1.3 to 1.75 pints (0.93 to 1.25 lb ai) of this product per acre as a postemergence treatment to soybeans from emergence up through the third trifoliate leaf stage. This product will not control emerged weeds so it must be applied to a weed-free soil surface or in a tank mixture with products that provide postemergence control of weeds present at the time of application.

**VISOR II XRT** can also be applied as part of a sequential soybean weed control program. If this product was applied as a preplant surface, preplant incorporated, or a preemergence treatment, a second treatment of this product can be applied postemergence provided that the total **VISOR II XRT** rate during any one crop does not 3.3 pints (2.35 lb ai) per acre.

#### Restrictions

- DO NOT** apply more than 1.75 pints (1.25 lb ai) of this product per acre postemergence.
- The total rate of this product rate applied preplant, preemergence or postemergence to soybeans during any one crop year should not exceed 3.3 pints (2.35 lb ai) of this product per acre.
- Preharvest Interval (PHI):** Make postemergence application at least 90 days before harvest.
- DO NOT** graze or feed treated forage or hay from soybeans to livestock following a postemergence application of this product.

#### SOYBEANS - VISOR II XRT COMBINATIONS

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Water or fluid fertilizer may be used as carrier for **VISOR II XRT** in combination with Clomazone, Ethalfuralin, Imazaquin, Imazethapyr, Linuron, Metribuzin or Metribuzin + Chlorimuron.

#### Restrictions

- For all of the following combinations, on soybeans use up to 3.3 pints (2.35 lb ai) of this product per acre preplant incorporated or preemergence treatment on soils having an organic matter content between 6% and 20%.
- The total rate of this product rate applied to soybeans during any one crop year should not exceed 3.3 pints (2.35 lb ai) per acre.

#### Tank Mixture with Metribuzin

In addition to those weeds controlled by **VISOR II XRT** alone, **VISOR II XRT** + Metribuzin, when applied as directed, also controls the following broadleaf weeds: cocklebur\*, hairy nightshade, hemp sesbania, jimsonweed\*, lambsquarters, prickly sida, ragweed, smartweed, velvetleaf, Venice mallow, and wild mustard.  
\*Partially controlled.

Apply **VISOR II XRT** and Metribuzin preplant incorporated or preemergence, using the appropriate rates from **Table 11**.

**Preplant Incorporated or Preemergence:** Follow instructions for use of **VISOR II XRT** alone under **Application Procedures**.

**Sequential:** Apply **VISOR II XRT** alone **Preplant Incorporated**, as specified in **Table 11** for this tank mixture. Follow with a preemergence application of Metribuzin during planting (behind the planter) or after planting, but before weeds or soybeans emerge.

Refer to the Metribuzin label for planting details and soybean variety restrictions. Follow most restrictive limitations and precautions on the **VISOR II XRT - Soybeans Alone** section of this product label and the Soybean directions on the Metribuzin label.

**Table 11: VISOR II XRT + Metribuzin – Soybeans**

	Broadcast Rates Per Acre			
Soil Texture*	0.5 to < 3% Organic Matter		≥ 3% Organic Matter	
	VISOR II XRT	Metribuzin	VISOR II XRT	Metribuzin
<b>Coarse</b> Loamy sand (over 2% organic matter), Sandy loam	1.1 to 1.3 (0.78 to 0.93 lb ai)	See Metribuzin label of product being used for application rates	1.3 pints (0.93 lb ai)	See Metribuzin label of product being used for application rates
<b>Medium</b>	1.3 to 1.75 pints (0.93 to 1.25 lb ai)		1.75 pints (1.25 lb ai)	
<b>Fine</b>	1.75 pints (1.25 lb ai)		1.75 to 2.2 pints (1.25 to 1.57 lb ai)	
<b>Mississippi Delta Only</b> Silty clay, Clay				
<b>Muck or Peat</b> (soils with more than 20% organic)	<b>DO NOT USE</b>			

\*On all sand and on loamy sand with less than 2% organic matter, **DO NOT USE** this tank mixture preemergence or the sequential treatment. **DO NOT USE** the tank mixture preplant incorporated on any sand, loamy sand or sandy loam or crop injury may occur.

\*On all sand and on loamy sand with less than 2% organic matter, **DO NOT** use this tank mixture preemergence or the sequential treatment. **DO NOT** use the tank mixture preplant incorporated on any sand, loamy sand or sandy loam or crop injury may occur.

#### Precautions

- DO NOT** use the tank mix or sequential application on soil with less than 0.5% organic matter or on alkaline soil with a pH over 7.4, or crop injury may occur.
- If heavy rain occurs soon after application, crop injury may result, especially in poorly drained areas where water stands for several days.

#### Tank Mixture with Linuron

In addition to those weeds controlled by **VISOR II XRT** alone, **VISOR II XRT** + Linuron applied preemergence, also controls the following broadleaf weeds: cocklebur\*, jimsonweed\*, lambsquarters, morningglory\*, prickly sida, ragweed, smartweed, velvetleaf\*, Venice mallow, and wild mustard.  
\*Partially controlled.

**Preemergence:** Apply during planting (behind planter) or after planting, but before weeds or soybeans emerge. Refer to the Linuron label for planting details. Apply the appropriate rates from **Table 12**.

**Table 12: *VISOR II XRT* + Linuron – Soybeans**

Soil Texture*	Broadcast Rates Per Acre			
	0.5 to Less Than 3% Organic Matter		3% Organic Matter or Greater	
	VISOR II XRT	Linuron	VISOR II XRT	Linuron
Coarse**	1.1 pints (0.78)	See Linuron label of product being used for application rates	1.3 pints (0.93 lb ai)	See Linuron label of product being used for application rates
Medium	1.3 pints (0.93 lb ai)		1.75 pints (1.25 lb ai)	
Fine	1.75 pints (1.25 lb ai)		1.75 to 2.2 pints (1.25 to 1.57 lb ai)	
Muck or Peat (soils with more than 20% organic	DO NOT USE			
* DO NOT use on sand, gravelly soils, or exposed subsoils. ** DO NOT use on loamy sand, except in the Northeastern U.S. on loamy sand with over 1% organic matter.				

**Precaution**

- To avoid crop injury, **DO NOT** use on soil with less than 0.5% organic matter.

**Tank Mixture with Trifluralin**

*VISOR II XRT* + Trifluralin tank mix applied preplant incorporated controls those weeds listed under *VISOR II XRT Applied Alone* and those weeds listed for Trifluralin Alone on the Trifluralin label. *VISOR II XRT* + Trifluralin may be applied by ground or aerial equipment and incorporated up to 14 days before planting. Follow the specified procedures on the Trifluralin and *VISOR II XRT* labels, using equipment that provides uniform 2-inch incorporation.

Apply *VISOR II XRT* + Trifluralin tank mix using the appropriate rate from the **Soybeans – *VISOR II XRT Alone*** section of this label and the Trifluralin Alone section of the Trifluralin label for the specific soil texture/organic matter classification and weed species expected.

To control Dinitroaniline-resistant (DNA-resistant) Goosegrass\* and other species on the respective labels where the soil organic matter is 3% or less, apply the rate in **Table 13**. Follow the most restrictive limitations and precautions on the **Soybeans – *VISOR II XRT Alone*** section of the *VISOR II XRT* label and the Soybean directions on the Trifluralin labels.

**Table 13: *VISOR II XRT* + Trifluralin - Organic Matter Content Less Than 3%**

Soil Texture	Broadcast Rates Per Acre		
	<i>VISOR II XRT</i>	Trifluralin	
	Less Than 3% Organic Matter	Less than 2% Organic Matter	2 to 3% Organic Matter
Coarse*	1.1 to 1.3 (0.78 to 0.93 lb ai)	See Trifluralin label of product being used for application rates	
Medium	1.3 pints (0.93 lb ai)		
Fine	1.75 pints (1.25 lb ai)		

\* Where a range of rates is given for this product use the minimum rate where DNA-resistant Goosegrass is the predominant species.

**Tank Mixture with Imazaquin**

This tank mixture controls all weeds controlled by *VISOR II XRT* alone and by Imazaquin alone. Refer to the *VISOR II XRT Applied Alone* section for weeds controlled by this product and to the Imazaquin label for weeds controlled by Imazaquin. Refer to the Imazaquin label for geographical locations where this tank mixture may be applied.

Apply *VISOR II XRT* + Imazaquin preplant incorporated or preemergence, using rates in **Table 14**. Follow use directions under **Application Instructions** on the Imazaquin label. For preplant incorporated applications, apply and incorporate within 30 days before planting. Observe all other precautions and limitations on the Imazaquin labels.

Follow the most restrictive limitations and precautions on the *VISOR II XRT - Soybeans Alone* section of the *VISOR II XRT* label and the Soybean directions on the Imazaquin label.

**Table 14: *VISOR II XRT* + Imazaquin – Soybeans**

Soil Texture	Broadcast Rates Per Acre			
	Less Than 3% Organic Matter		3% or More Organic Matter	
	<i>VISOR II XRT</i>	Imazaquin	<i>VISOR II XRT</i>	Imazaquin
Coarse	1.1 pints (0.78 lb ai)	See Imazaquin label of product being used for application rates	1.3 pints (0.93 lb ai)	See Imazaquin label of product being used for application rates
Medium	1.3 pints (0.93 lb ai)		1.75 pints (1.25 lb ai)	
Fine	1.75 pints (1.25 lb ai)		1.75 to 2.2* pints (1.25 to 1.57 lb ai)	
Muck or Peat (soils with more than 20% organic	DO NOT USE			
* Use the higher rate of this product if heavy weed infestations are expected.				

**Tank Mixture with Metribuzin + Chlorimuron Ethyl**

This tank mixture controls all weeds controlled by *VISOR II XRT* alone and by Metribuzin + Chlorimuron Ethyl alone. Refer to the *VISOR II XRT Applied Alone* section for weeds controlled by this product and to the Metribuzin + Chlorimuron Ethyl label for weeds controlled by Metribuzin + Chlorimuron Ethyl.

Apply preplant incorporated or preemergence, using the appropriate rates from **Table 15**.

**Preplant Incorporated:** Apply within 2 weeks of planting. Uniformly incorporate into the top 1 to 2 inches of soil before planting soybeans.

**Preemergence:** Apply after planting, but before soybeans emerge. Follow all use directions, varietal restrictions, limitations, use precautions and information regarding application to Soybeans and rotational restrictions on the *VISOR II XRT* and Metribuzin + Chlorimuron labels.

**Table 15: *VISOR II XRT* + Metribuzin + Chlorimuron Ethyl – Soybeans**

Soil Texture	Broadcast Rates Per Acre		
	VISOR II XRT		Metribuzin + Chlorimuron
	<Less Than 3% Organic Matter	3% or More Organic Matter	
Coarse	1.1 pints (0.78 lb ai)	1.3 pints (0.93 lb ai)	*
Medium	1.3 pints (0.93 lb ai)	1.75 pints (1.25 lb ai)	
Fine	1.75 pints (1.25 lb ai)	1.75 to 2.2 pints (1.25 to 1.57 lb ai)	
* Refer to the Metribuzin + Chlorimuron label for appropriate rate according to geographical location, soil and organic matter classification and pH limitations.			

**Restriction**

- DO NOT** apply to sand, or to any soil with less than 0.5% organic matter, or to any soil with pH greater than 7.0, except as noted on the Metribuzin + Chlorimuron Ethyl label.



### Tank Mixture with Clomazone\*

This tank mixture controls all weeds controlled by **VISOR II XRT** alone and by Clomazone alone. Refer to the **VISOR II XRT Applied Alone** section for weeds controlled by this product and to the Clomazone label for weeds controlled by Clomazone.

Apply **VISOR II XRT** + Clomazone preplant incorporated, using rates in **Table 16**. Follow all Clomazone application instructions as to incorporation interval, geographical location, equipment operation, soil moisture conditions, etc.

\* Follow the most restrictive limitations and precautions on the **VISOR II XRT – Soybeans** **Alone** section of this label and the Soybean directions on the Clomazone label, including rotational restrictions.

**Table 16: VISOR II XRT + Clomazone – Soybeans**

Soil Texture	Broadcast Rates Per Acre			
	<b>VISOR II XRT</b>		<b>Clomazone</b>	
	0.5 - 3% Organic Matter	Greater Than 3% Organic Matter	Northern Area	Southern Area
Coarse	1.1 pints (0.78 lb ai)	1.3 pints (0.93 lb ai)	See Clomazone label of product being used for application rates	
Medium	1.3 pints (0.93 lb ai)	1.75 pints (1.25 lb ai)		
Fine	1.75 pints (1.25 lb ai)	1.75 to 2.2 pints (1.25 to 1.57 lb ai)		

### Tank Mixture with Ethafluralin

This tank mixture controls all weeds controlled by **VISOR II XRT** alone and by Ethafluralin alone. Refer to the **VISOR II XRT Applied Alone** section for weeds controlled by this product and to the Ethafluralin label for weeds controlled by Ethafluralin.

Apply **VISOR II XRT** and Ethafluralin preplant incorporated, using the appropriate rates from **Table 17**.

**Preplant Incorporated:** Follow specified soil preparation procedures for Ethafluralin. Refer to the Ethafluralin label for incorporation directions.

**Sequential:** Apply Ethafluralin alone preplant incorporated as specified on the Ethafluralin label. Follow with a preemergence application of **VISOR II XRT** during planting (behind the planter) or after planting, but before weeds or soybeans emerge.

Follow the most restrictive limitations and precautions on the **VISOR II XRT – Soybeans** **Alone** section of the **VISOR II XRT** label and the Soybean directions on the Ethafluralin label.

**Table 17: VISOR II XRT + Ethafluralin – Soybeans**

Soil Texture	Broadcast Rates Per Acre			
	Less Than 3% Organic Matter		3% or More Organic Matter	
	VISOR II XRT	Ethafuralin	VISOR II XRT	Ethafuralin
Coarse	1.1 pints (0.78 lb ai)	See Ethafuralin label of product being used for application rates	1.75 pints (1.25 lb ai)	See Ethafuralin label of product being used for application rates
Medium *	1.75 to 2.2 pints (1.25 to 1.57 lb ai)		1.75 to 2.2 pints (1.25 to 1.57 lb ai)	
Fine *			2.2 to 2.7 pints	
Muck or Peat (soils with more than 20% organic	DO NOT USE			

\*For Eastern black nightshade on these soils, apply Ethafuralin at labeled rate and follow with 2 incorporation passes.

\*For Eastern black nightshade on these soils, apply Ethafluralin at labeled rate and follow with 2 incorporation passes.

### Tank Mixture with Imazethapyr

This tank mixture controls all weeds controlled by **VISOR II XRT** alone and by Imazethapyr alone. Refer to the **VISOR II XRT Applied Alone** section for weeds controlled by this product and to the Imazethapyr label for weeds controlled by Imazethapyr. Refer to the Imazethapyr label for geographical locations where this tank mixture may be applied.

Apply **VISOR II XRT** + Imazethapyr early preplant, preplant incorporated, or preemergence after planting, using rates in **Table 18**. Application can be made in water or liquid fertilizer. Follow all use directions under Soil Applications on the Imazethapyr label. For early preplant and preplant incorporated applications, apply within 30 days before planting.

Follow the most restrictive limitations and precautions on the **VISOR II XRT – Soybeans** **Alone** section of this label and the Soybean directions on the Imazethapyr label, including rotational restrictions.

**Table 18: VISOR II XRT + Imazethapyr – Soybeans**

Soil Texture	Broadcast Rates Per Acre		
	Less Than 3% Organic Matter	3% or More Organic Matter	Less Than 3% - 3% or More Organic Matter
	<b>VISOR II XRT</b>		
Coarse	1.1 pints (0.78 lb ai)	1.3 pints (0.93 lb ai)	See Imazethapyr label of product being used for application rates
Medium	1.3 pints (0.93 lb ai)	1.75 pints (1.25 lb ai)	
Fine	1.75 pints (1.25 lb ai)	1.75 to 2.2 pints (1.25 to 1.57 lb ai)	

**Sequential:** Apply **VISOR II XRT** early preplant, preplant incorporated, or preemergence after planting at 1.1 pints (0.78 lb ai) per acre on coarse soils and 1.3 pints (0.93 lb ai) per acre on medium- and fine-textured soils. Follow with a sequential postemergence application of Imazethapyr to control emerged weeds according to the Imazethapyr label. **VISOR II XRT** will improve the consistency and level of control from Imazethapyr on most grass species. Refer to the Imazethapyr postemergence label for a listing of weeds controlled, application rate, and growth stage limitations.

### Tank Mixture with Metribuzin, Imazaquin, Linuron, Metribuzin + Chlorimuron or Imazethapyr Plus Paraquat or Glyphosate for Minimum-Tillage or No-Tillage Systems

In minimum-tillage or no-tillage systems where Soybeans are planted directly into a cover crop, stale seedbed, established sod or previous crop residues, the contact herbicides Paraquat or Glyphosate may be added to a tank mix of this product plus one of the following:

Imazaquin  
Imazethapyr  
Linuron

Metribuzin  
Metribuzin + Chlorimuron

When used as directed, the Paraquat portion of the tank mixture controls most emerged weeds and suppresses many perennial weeds. Glyphosate combinations will control emerged annual and perennial weeds when applied as directed on the Glyphosate label. This product + tank mix herbicides listed above portion of the tank mixture provides preemergence control of the weeds listed on this label in the tank mixture section for this product + each respective herbicide.

Refer to the label of each product used in combination and observe the planting details, Soybean variety restrictions, information regarding application to Soybeans, geographical restrictions, and all other use precautions and limitations.

**Application:** Apply before, during, or after planting, but before the soybeans emerge. Add Paraquat or Glyphosate and apply as directed on the product labels.

**Paraquat:** Apply as directed on the Paraquat label. This treatment will not control weeds taller than 6 inches.

**Glyphosate:** See the Glyphosate label for weeds controlled, specified rates, and other use directions.

Apply in 20 to 60 gallons of water or fluid fertilizer per acre with ground equipment.

## Restriction

- **DO NOT** apply combinations containing Paraquat in suspension type liquid fertilizers, as the activity of paraquat will be reduced.

## **VISOR II XRT + Metribuzin + Paraquat or Glyphosate**

Apply labeled rate of Metribuzin with 1.3 pints (0.93 lb ai) of this product per acre on loamy sand with over 2% organic matter, with 1.75 pints (1.25 lb ai) of this product per acre on medium soils, or with 1.75 to 2.2 pints (1.25 to 1.57 lb ai) of this product per acre on fine soils. Refer to the Metribuzin label for appropriate rate according to geographical location, soil and organic matter classification, pH limitations and all other use directions.

## Precautions

- To avoid crop injury, **DO NOT** use this tank mixture on soil with less than 0.5% organic matter, on alkaline soil with a pH over 7.4, or on all sand and on loamy sand with less than 2% organic matter.
- If heavy rain occurs soon after application, crop injury may result, especially in poorly drained areas where water stands for several days, or where the seeding slit has not been properly closed.

## **VISOR II XRT + Imazaquin + Paraquat or Glyphosate**

Apply labeled rate of Imazaquin with 1.3 pints (0.93 lb ai) of this product per acre on coarse soils, with 1.75 pints (1.25 lb ai) of this product per acre on medium soils or with 2.2 pints (1.57 lb ai) of this product per acre on fine soils. Refer to the Imazaquin label for appropriate rate according to geographical location, soil and organic matter classification, pH limitations and all other use directions.

## Restrictions

- **Preharvest Interval (PHI):** **DO NOT** apply within 90 days of harvest.
- **DO NOT** graze or feed treated soybean forage, hay, or straw to livestock.

## **VISOR II XRT + Linuron + Paraquat or Glyphosate**

Apply labeled rate of Linuron with 1.3 pints (0.93 lb ai) of this product per acre on coarse soils, with 1.75 pints (1.25 lb ai) of this product per acre on medium soils or with 1.75 to 2.2 pints (1.25 to 1.57 lb ai) of this product per acre on fine soils. Refer to the Linuron label for appropriate rate according to geographical location, soil and organic matter classification, pH limitations and all other use directions.

## Precaution

**DO NOT** use on soil with less than 0.5% organic matter or crop injury may occur.

## Restrictions

- **DO NOT** use on loamy sand, except in the Northeastern U.S. on loamy sand with over 1% organic matter or injury may occur.
- **DO NOT** use on sand, gravelly soils or exposed subsoils or injury may occur.

## **VISOR II XRT + Metribuzin + Chlorimuron Ethyl + Paraquat or Glyphosate**

Use only where soils have 0.5 to 5% organic matter. Apply labeled rate of Metribuzin + Chlorimuron with 1.3 pints (0.93 lb ai) of this product per acre on coarse soils (except sand), with 1.75 pints (1.25 lb ai) of this product per acre on medium soils or with 1.75 to 2.2 pints (1.25 to 1.57 lb ai) of this product per acre on fine soils. Refer to the Metribuzin + Chlorimuron label for appropriate rate according to geographical location, soil and organic matter classification, pH limitations and all other use directions.

## Restriction

- **DO NOT** apply to sand, or to any soil with less than 0.5% organic matter, or to any soil with pH greater than 7.0, except as noted on the Metribuzin + Chlorimuron label.

## **VISOR II XRT + Imazethapyr + Paraquat or Glyphosate**

Apply labeled rate of imazethapyr with 1.3 pints (0.93 lb ai) of this product per acre on coarse soils, with 1.75 pints (1.25 lb ai) of this product per acre on medium soils or with 2.2 pints (1.57 lb ai) of this product per acre on fine soils. Refer to the Imazethapyr label for weeds controlled, specified rates and other use directions.

## **POSTEMERGENCE USE ON SOYBEANS - VISOR II XRT TANK MIXTURES**

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

## **Tank Mixture with Glyphosate Products**

**VISOR II XRT** at 1.3 to 1.75 pints (0.93 to 1.25 lb ai) per acre may be tank mixed with glyphosate products at labeled rates and applied from emergence up through the third trifoliate leaf stage of Roundup Ready or glyphosate-resistant soybeans. This product alone will not control emerged weeds. Use this treatment only on soybeans designated for use with glyphosate (e.g., Roundup Ready or glyphosate-resistant soybeans). The glyphosate product must be registered for postemergence use in Roundup Ready or glyphosate-resistant soybeans.

## **Tank Mixture with Imazethapyr**

**VISOR II XRT** at 1.3 to 1.75 pints (0.93 to 1.25 lb ai) per acre may be tank mixed with Imazethapyr at labeled rates and applied from emergence up through the third trifoliate leaf stage of soybeans. This product alone will not control emerged weeds.

## **Tank Mixture with Glufosinate-ammonium Products**

**VISOR II XRT** at 1.3 to 1.75 pints (0.93 to 1.25 lb ai) per acre may be tank mixed with Glufosinate products at labeled rates and applied from emergence up through the third trifoliate leaf stage of soybeans. This product alone will not control emerged weeds. Use this treatment only on soybeans designated for use with glufosinate (e.g. LibertyLink).

## Precaution

- The use of adjuvants such as COC or UAN with this product may result in temporary crop injury. Follow the tank mix product label for adjuvant directions.

## Restrictions

- **DO NOT** apply more than 1.75 pints (1.25 lb ai) per acre postemergence.
- **Preharvest Interval (PHI):** Make postemergence application at least 90 days before harvest.
- **DO NOT** graze or feed treated forage or hay from soybeans to livestock following a postemergence application of this product.

## **SUGAR BEETS - VISOR II XRT ALONE**

### Postemergence Applications

**VISOR II XRT** may be applied postemergence to sugar beets after the sugar beets have reached the first true-leaf stage. However, because this product is primarily a soil-active herbicide, it must be applied prior to weed emergence in order to provide consistent control of listed weeds. As such, weeds that are emerged with or before the crop, or that are present at the time this product is applied, must be controlled with another appropriately labeled herbicide. Apply this product at 1.3 pints (0.93 lb ai) per acre on coarse soils, 1.75 pints (1.25 lb ai) per acre on medium soils or 2.2 pints (1.57 lb ai) per acre on fine soils. More than one postemergence application may be applied, but the total should not exceed 3.45 pints (2.45 lb ai) per acre. Weeds present at the time of application will not be controlled.

## Precaution

- In coarse soils, this product applied before emergence of sugar beets (i.e., other than postemergence) may cause injury.

## Restrictions

- **DO NOT** apply more than 3.45 pints (2.45 lb ai) per acre postemergence.
- **Preharvest Interval (PHI):** **DO NOT** harvest within 60 days after the last application.

## **SUGAR BEETS - VISOR II XRT TANK MIX COMBINATIONS**

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixings. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

**VISOR II XRT** may be tank mixed with Clethodim, Clopyralid, Desmedipham, Phenmedipham + Desmedipham + Ethofumesate, Quizalofop-p-ethyl, Sethoxydim or Triflurosulfuron-methyl and applied to sugar beets. Tank mixtures of these products with **VISOR II XRT** will increase the risk of crop injury over that of either product applied alone, as the **VISOR II XRT** formulation has some adjuvant properties. The addition of a spray adjuvant such as crop oil concentrates (COC's) or methylated seed oils (MSO's) can further increase the risk of crop injury. Injury risk can be reduced by using the lowest effective rate of the tank mix partner(s) and/or adjuvant and by avoiding applications under adverse growing conditions or high soil or air humidity. Refer to the individual product labels and follow all use restrictions and limitations.

## SUNFLOWERS - VISOR II XRT ALONE

### Preplant Incorporated or Preemergence

Within the rate ranges given below, use the higher rate of **VISOR II XRT** if heavy weed infestations are expected. On coarse soils with organic matter less than 3%, apply 1.3 to 1.75 pints (0.93 to 1.25 lb ai) of this product per acre; apply 1.75 pints (1.25 lb ai) per acre if organic matter is 3% or greater. On medium soils, apply 1.75 to 2.2 pints (1.25 to 1.57 lb ai) of this product per acre. On fine soils with organic matter of less than 3%, apply 1.75 to 2.2 pints (1.25 to 1.57 lb ai) of this product per acre; apply 2.2 to 2.7 pints (1.57 to 1.9 lb ai) per acre if organic matter is 3% or greater.

### Restrictions

- **DO NOT** allow livestock to graze or feed in treated area.
- **DO NOT** exceed the maximum label rates for sunflowers for the soil type.

## TOMATOES - VISOR II XRT ALONE

### Transplanted

**VISOR II XRT** may be applied preplant incorporated or preplant before transplanting. If the latter method is used, keep soil disturbance to a minimum during the transplanting operation. Application may also be post directed to transplants after the first settling rain or irrigation. When an application is made post-directed, apply in a minimum of 20 gallons of water per acre and minimize contact with tomato plants. This product will not control emerged weeds. In bedded transplanted tomatoes, apply this product preplant non-incorporated to the top of the pressed bed as the last step prior to laying plastic. This product may also be used to treat row-middles in bedded tomatoes, as long as the total amount of this product does not exceed the maximum allowed per crop.

### Seeded

**VISOR II XRT** may be applied post-directed to direct-seeded tomatoes. Tomato plants must be at least 4 inches tall at the time of application, and the product must be applied in a minimum of 20 gallons of water per acre. Minimize spray contact with tomato plants. This product will not control emerged weeds.

**Tomato Use Rates:** On coarse soils, apply 1.3 to 1.75 pints (0.93 to 1.25 lb ai) of this product per acre if organic matter is less than 3% or 1.75 pints (1.25 lb ai) per acre if organic matter is 3% or greater. On medium soils, apply 1.75 to 2.2 pints (1.25 to 1.57 lb ai) of this product per acre. On fine soils, apply 1.75 to 2.2 pints (1.25 to 1.57 lb ai) of this product per acre if organic matter is less than 3% or (2.2 to 2.7 pints (1.57 to 1.9 lb ai) per acre if organic matter is 3% or greater.

### Precautions

- This product may damage transplants that have been weakened by any cause. To prevent damage, plant only healthy transplants. **DO NOT** plant when wet, cool, or unfavorable growing conditions exist.
- In transplanted tomatoes, if this product is applied preplant incorporated, incorporate to a depth less than the depth of transplanting, and use the lower end of the rate range for the given soil type, or damage may occur.
- For row-middle applications where tomatoes are grown on sandy soils and where high soil moisture conditions can exist (e.g., low binding and high evaporation conditions), as may be found in the States of Florida, Georgia, Maryland, and Virginia, there is potential for crop injury in the form of leaf epinasty. The risk of this type of injury can be reduced by:
  - incorporating this product immediately following application,
  - applying this product seven or more days before transplanting (but only after the beds have been formed),
  - minimizing the application of this product onto the plastic of the bed, or
  - any combination of the above.

### Restrictions

- **DO NOT** apply to varieties or cultivars with unknown tolerance to this product.
- **DO NOT** exceed the maximum label rate for the soil texture per year.
- Apply only by ground application.
- **90-Day Preharvest Interval (PHI):** If the single application rate of this product is greater than 1.75 pints (1.25 lb ai) per acre (up to 2.7 lb ai (1.9 lb ai) per acre), **DO NOT** harvest tomatoes within 90 days of application.
- **30-Day Preharvest Interval (PHI):** If the application of this product does not exceed 1.75 pints (1.25 lb ai) per acre, **DO NOT** harvest tomatoes within 30 days of application.

**When applying at 1.75 pints (1.25 lb ai) per acre with a 30-day PHI, the following restrictions apply:**

- **DO NOT** exceed two applications per growing season.
- The use of adjuvants is prohibited.
- Applications may be made using ground equipment, in concentrated spray volumes.
- Applications may be made as a foliar broadcast spray to the soil within 1 week of transplanting and again at blooming/fruiting to the row middles as a banded/directed application 38 to 77 days after the first treatment.

## STORAGE AND DISPOSAL

**DO NOT** contaminate water, food, or feed by storage or disposal.

### Pesticide Storage

This product may be stored at temperatures down to 30 degrees below 0°F.

### Pesticide Disposal

Open dumping is prohibited. Wastes resulting from the use of this product are toxic. Improper disposal of unused pesticide, spray mixture, or rinsate is a violation of federal law. Pesticide, spray mixture, or rinsate that cannot be used according to label instructions must be disposed of according to federal, state, or local procedures. For guidance in proper disposal methods, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office.

### Container Handling

**NONREFILLABLE CONTAINER (EQUAL TO OR LESS THAN 5 GALLONS): DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling, if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

**NONREFILLABLE CONTAINER (GREATER THAN 5 GALLONS): DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Offer for recycling, if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

**Pressure rinse as follows (all sizes):** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**REFILLABLE CONTAINER:** Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. After triple rinsing is complete, and the container is not suitable for refilling or reconditioning, offer the container for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

#### **CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY**

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of INNICTIS CROP CARE, LLC or Seller. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW all such risks shall be assumed by Buyer and User and Buyer and User agree to hold INNICTIS CROP CARE, LLC and Seller harmless for any claims relating to such factors.

INNICTIS CROP CARE, LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or INNICTIS CROP CARE, LLC, and TO THE EXTENT CONSISTENT WITH APPLICABLE LAW Buyer and User assume the risk of any such use. To the extent consistent with applicable law INNICTIS CROP CARE, LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, neither INNICTIS CROP CARE, LLC nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF INNICTIS CROP CARE, LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF INNICTIS CROP CARE, LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

INNICTIS CROP CARE, LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of Sale and Limitation of Warranty and Liability which may not be modified except by written agreement signed by a duly authorized representative of INNICTIS CROP CARE, LLC.

All trademarks are the property of their respective owners.