
SUPPLEMENTAL LABELING

Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, North Carolina 27419-8300
SCP 828A-S4 0111

GROUP	9	FUNGICIDE
-------	----------	-----------

Vangard® WG

Fungicide

Active Ingredient:

Cyprodinil: 4-cyclopropyl-6-methyl-N-phenyl-pyrimidinamine* 75.0%

Other Ingredients: 25.0%

Total: 100.0%

*CAS No. 121552-61-2

Vangard WG is a water-dispersible granule.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

EPA Reg. 100-828

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed. Before using Vangard WG as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

syngenta

DIRECTIONS FOR USE

CROP USE DIRECTIONS – TREES, NUTS AND VINES

Crop	Disease	Rate oz./Acre	Remarks
Pome Fruits Apples Crabapples Loquat Mayhaw Pears (see tank mixture instructions) Quince	Prebloom Scab (<i>Venturia</i> spp.)	Vanguard WG alone 5	For pome fruits except pear, begin application at green tip and continue on a 7- to 10-day interval. Under severe disease pressure, use the shorter interval.
		Vanguard WG tank mixtures 3 - 5	For scab control utilizing multiple modes of action, apply Vanguard WG in tank mixture with the label rate of a protectant or systemic fungicide registered on pome fruit. Make applications on a 7- to 10-day interval.
	Pink, bloom, post- bloom Scab (<i>Venturia</i> spp.)	Vanguard WG tank mixtures 3 - 5	Apply Vanguard WG in tank mix combination with the label rate of a protectant fungicide. Use of the label rate of an EBDC fungicide will broaden the disease control spectrum to include cedar apple rust. The addition of the label rate of sulfur or a sterol inhibitor (SI) fungicide to this tank mix will broaden the disease control spectrum to include powdery mildew.
Application Instructions: Application may be made by ground or air. Good coverage is essential for good disease control. Use a minimum of 20 gallons/A spray volume by air. Make no more than two applications by air. Make additional applications by ground.			

Specific Use Restrictions

- When applying to pears, apply Vanguard WG in tank mix only.
- Make no more than two applications by air.
- Do not apply more than 30 oz./A of Vanguard WG (1.4 lb. a.i./A of cyprodinil) per plot of land per year.
- Do not apply within 0 days of harvest (0 day PHI).

Vanguard® and the Syngenta logo are trademarks of a Syngenta Group Company

©2011 Syngenta