
SUPPLEMENTAL LABELING

Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, North Carolina 27419-8300
SCP 828A-S3 0111

GROUP	9	FUNGICIDES
-------	----------	------------

Vangard® WG

Fungicide

Active Ingredient:	
Cyprodinil: 4-cyclopropyl-6-methyl-N-phenyl-pyrimidinamine*	75.0%
Other Ingredients:	25.0%
Total:	100.0%

*CAS No. 121552-61-2

KEEP OUT OF REACH OF CHILDREN.

CAUTION

EPA Reg. No. 100-828

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed. Before using Vangard WG as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DIRECTIONS FOR USE

Crop	Disease	Rate oz./Acre	Remarks
Pistachios	Botrytis (<i>Botrytis</i> spp.)	5.5 - 7	Make the first application during early bloom and repeat applications at 14-day intervals if conditions remain favorable for disease development.
	Alternaria (<i>Alternaria alternata</i>)		
Application Instructions: Application may be made by ground or air. Good coverage is essential for good disease control. Use a minimum of 20 gallons/A spray volume by air. Make no more than two applications by air. Make additional applications by ground.			

Specific Use Restrictions

- Make no more than two applications by air.
- Do not apply more than 28 oz./A of Vanguard WG (1.3 lb. a.i./A of cyprodinil) per plot of land per year.
- Do not apply within 7 days of harvest (7 day PHI).

Vanguard® and the Syngenta logo are trademarks of a Syngenta Group Company

©2011 Syngenta