

RESTRICTED USE PESTICIDE

Due to acute inhalation toxicity to humans. For retail sale to and use by certified applicators or persons under their direct supervision and only for those uses covered by the certified applicator's certification.



FIFRA 2(ee)

FOR END OF SEASON CROP TERMINATION TREATMENT OF: Melon (including: bitter melon, cantaloupe, hybrids and/or cultivars, citron melon, crenshaw melon, golden pershaw melon, mango melon, honeydew melon, muskmelon, Persian melon, pineapple melon, Santa Claus melon, snake melon, watermelon)

FIFRA 2ee RECOMMENDATION

Vapam[®] HL[™]

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF CALIFORNIA.

EPA Reg. No.: 5481-468

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

THIS BULLETIN MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION.

Before applying Vapam HL, read the complete label affixed to the product container. All applicable directions, restrictions and precautions on the EPA-registered label are to be followed. Use of Vapam HL according to this supplemental labeling is subject to the use precautions and limitations imposed by the label on the container.

IMPORTANT: Read the entire Directions for Use and the Limited Warranty and Disclaimer on the EPA-registered Vapam HL label before using this product.

Use Information

End of Season Crop Termination with Drip (Trickle) Chemigation of Beds without Existing Tarp/Plastic Mulch. Follow the instructions listed in the Vapam HL Section 3 label for preparation and application conditions for pre-plant drip chemigation, with the exception of, instructions for soil preparation which are not applicable for this treatment. Use the following additional steps:

- Use existing drip or trickle tape in the bed.
- Ensure that all drip (trickle) tape is functional with all major leaks repaired.
- Apply Vapam HL in beds that are not covered with plastic and have a drip tape buried at a depth of >6 inches.
- Apply Vapam HL soon after the last harvest when the plants are still actively growing.

- Apply Vapam HL at 18-56 gallons/acre (broadcast equivalent) broadcast at a flow rate that allows good lateral movement of water in the bed.
- Contact the local extension specialist for more information about flow rates that provide good fumigant distribution in soil on the crops listed above.
- Use the maximum application rate for fields with hard-to-control weeds, pathogens and nematodes such as nutsedge (*Cyperus* sp.), Fusarium wilt (*Fusarium oxysporum* f. sp. *niveum*) and root knot nematodes (*Meloidogyne* sp.) and Vapam HL application rate per treated acre is not to exceed 75 gallons unless the broadcast equivalent calculation is greater than 56 gal/acre, then use a maximum broadcast equivalent rate of 56 gal/acre.
- Dispose of all crop and/or plant residues following treatment by removal, tillage or other appropriate means.

Restrictions

The terminated crop must not be used for any food or feed purposes after the product has been applied.

Vapam HL is a trademark owned by Amvac Chemical Corporation.

Registrant:

AMVAC Chemical Corporation
4695 MacArthur Court, Suite 1200
Newport Beach, CA U.S.A.
1-888-462-6822