

Product Bulletin



Dow AgroSciences

Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Vastlan™

EPA Reg. No. 62719-687

2(ee) Recommendation †

Control of susceptible weeds pre or post planting of certain annual and perennial grasses

ATTENTION

† This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA.

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- Read the label affixed to the container for Vastlan™ specialty herbicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Except as specified in this product bulletin, use of Vastlan is subject to all use precautions and limitations imposed by the label affixed to the container for Vastlan.

Directions for Use

Vastlan controls only broadleaf plants that are emerged at the time of application. See the main label for a list of susceptible weeds.

PRECAUTIONS

Post emergence applications on newly emerged or established sorghum sudangrass, teff, crabgrass, and pearl millet are not recommended. Unacceptable injury or death of these grasses may occur with Vastlan applications.

Pre-plant applications for control of susceptible emerged weeds

Apply Vastlan at a maximum of 1.5 pts/acre with a non-ionic surfactant to control susceptible weeds prior to planting tall fescue, orchardgrass, timothy, and annual ryegrass, sorghum sudangrass, teff, crabgrass, and pearl millet. Applications can be made the same day of planting.

Post planting applications for control of susceptible emerged weeds

Apply Vastlan at a maximum of 1.5 pts/acre with a non-ionic surfactant to tall fescue, orchardgrass, timothy, and annual ryegrass after seedling plants have at least reached the V3 stage (three collared leaves).

™®Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

R409-009

Issued: 08/14/17

Initial printing.