



Section 2(ee) Recommendation

Date: December 10, 2015
Product: Velista™, EPA Reg. No. 100-1534
Use: Addition of Diseases as Pests Controlled on Turfgrasses
State: Louisiana

Directions for Use

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- The user must refer to the federally approved labels for the above noted product and read and follow all directions for use, restrictions, and precautions.
- The user should have this recommendation in its possession at the time of use.
- Use of this product according to this bulletin has not been reviewed or endorsed by the Office of the Indiana State Chemist.
- This recommendation for use of this product is permitted under Section 2(ee) of FIFRA and has not been submitted to or been approved by EPA.

Disease	Application Site	Rate oz/1,000 sq ft	Application Interval (Days)	Comments
Rapid Blight (<i>Labyrinthula terrestris</i>)	All Turfgrass	0.5	14	Apply when conditions are favorable for disease development. Do not apply more than 2.2 oz product/1,000 sq ft/year.
Fairy Ring (<i>Basidiomycetes</i>)	All Turfgrass	0.5-0.7	14-28	Apply as part of a preventative program. Rotate with Heritage® and Briskway® fungicides. Do not apply more than 2.2 oz product/1,000 sq ft/year.
Leaf and Sheat Spot (<i>Rhizoctonia zeae</i>)	All Turfgrass	0.3-0.5	14	Use as part of a preventative program in tank mixture and/or rotation with Daconil Action™, Heritage and Briskway fungicides. Do not apply more than 2.2 oz product/1,000 sq ft/year.
Spring Dead Spot (<i>Ophiosphaerella</i> spp.)	All Turfgrass	0.7	28	Make two applications on a 28 day interval prior to winter dormancy. Do not apply more than 2.2 oz of product/1,000 sq ft/year.

Disease	Application Site	Rate oz/1,000 sq ft	Application Interval (Days)	Comments
SummerPatch (<i>Magnaporthe poae</i>)	All Turfgrasses	0.3-0.5	14-28	During periods of high disease pressure, tank mix with a fungicide that is registered for the control of summer patch. Apply when conditions are favorable for disease development. Do not apply more than 2.2 oz of product/1,000 sq ft/year.
Brown Ring Patch (<i>Waitea circinata</i> <i>var circinata</i>)	All Turfgrasses	0.5	14	Apply when conditions are favorable for disease development. Do not apply more than 2.2 oz of product/1,000 sq ft/year.
Large Patch (<i>Rhizoctonia solani</i>)	Large Patch (<i>Rhizoctonia solani</i>)	0.5-0.7	14-28	Make 1-2 applications in the fall as a preventative. Apply an early spring application if disease pressure is high. Do not apply more than 2.2 oz product/1,000 sq ft/year.

Velista™, Briskway®, Daconil Action®, Heritage® and the Syngenta logo are trademarks of a Syngenta Group Company.

©2015 Syngenta

2(ee) Registrant:
Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, NC 27419-8300

Expiration Date: December 10, 2019

Label Code: LA1534031AC1215