



Bayer CropScience LP
800 N. Lindbergh Blvd.
St. Louis, Missouri 63167
1-866-99BAYER (1-866-992-2937)

VELUM®

For suppression of Nematodes at a reduced
in-furrow application rate in Corn
EPA Reg. No. 264-1078

FIFRA 2(ee) Recommendation

**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATES OF ALABAMA,
ARKANSAS, FLORIDA, GEORGIA, KANSAS, KENTUCKY, LOUISIANA,
MISSOURI, MISSISSIPPI, NORTH CAROLINA, NEW MEXICO, OKLAHOMA,
SOUTH CAROLINA, TENNESSEE, TEXAS, VIRGINIA**

This recommendation is valid until December 31, 2024

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its label.

VELUM is physically compatible with most commonly used fungicide, herbicide, insecticide, and foliar nutrient products. However, the compatibility of VELUM with all potential tank-mix partners has not been fully investigated. It is highly recommended that users conduct a "jar-test" to make sure that tank mix partners are compatible before mixing. Contact your local Bayer rep for more information.

FIELD CORN; FIELD CORN GROWN FOR SEED; POPCORN

Pest Suppression	Application Rate	Application Instructions
Nematodes	3.0 fl oz/acre	Soil Application Apply specified dosage using any of the following methods: <ul style="list-style-type: none">In-furrow spray during planting directed on or below seed.

Restrictions:

- Do not apply more than 13.7 fl oz of VELUM (0.446 lbs Fluopyram) per acre per year, regardless of formulation or method of application (soil or foliar).
- Apply using ground equipment.
- Do not apply VELUM within 14 days of harvest.
- Do not allow livestock to graze treated area for 14 days and do not harvest for food or feed within 14 days of application.
- To limit the potential for development of disease resistance to this fungicide class, do not make more than 2 sequential applications of VELUM or any Group 7-containing fungicide before rotating with a fungicide from a different Group.
- For soil application, to limit the potential for development of disease resistance to this chemical class, the first foliar fungicide application after VELUM should be a product from a different FRAC group.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937).

*As with any crop-protection product, always read and follow label directions.
For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*

002151_L2EE_01.01