

FIFRA 2(ee) Recommendation

FOR DISTRIBUTION AND USE ONLY WITHIN ALABAMA, ARKANSAS, FLORIDA, GEORGIA, LOUISIANNA, MISSOURI, MISSISSIPPI, NORTH CAROLINA, NEW MEXICO, OKLAHOMA, SOUTH CAROLINA AND TEXAS

This recommendation expires December 31, 2025.

VELUM®

EPA Reg. No. 264-1078

To Allow for Additional Soil Methods for Nematode Suppression in Pecan

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its label.

PECAN		
Pest Suppression	Application Rate	Application Instructions
Nematodes	4.0 – 6.84 fl oz/acre (0.130 – 0.222 lb/acre fluopyram)	In addition to the Soil Application methods listed on the label. Velum may be applied as follows: • Soil application into the planting hole as a drench at planting. Divide the per acre use rate (4.0 to 6.84 fl oz/acre) by the number of trees to be planted in the acre. It is recommended to apply in a high enough gallonage to ensure treatment is distributed through the soil profile. • Root Dip at planting. Soak bareroot trees in a stock solution for 30 minutes. To make 50 gallons of stock solution, add 0.5 fl oz of Velum. It is recommended after the root dip to apply any remaining dip solution into the holes at planting or to the base of the trees after planting. Use no more than a total of 6.84 fl oz of Velum per acre as a root dip. Minimum of 30-day interval between soil applications.

PECAN (continued)

Restrictions:

- Maximum single application rate: 6.84 fl oz/acre of VELUM (0.222 lb/acre fluopyram)
- Maximum annual application rate: 13.7 fl oz of VELUM per acre (0.445 lb/acre fluopyram) per year.
- Maximum number of applications per year: 2 (at 6.84 fl oz/acre of VELUM)
- **DO NOT** apply more than 0.446 lbs Fluopyram per acre per year, regardless of formulation or method of application (soil or foliar).
- Apply using ground or chemigation equipment
- **DO NOT** apply VELUM within 14 days of harvest.
- To limit the potential for development of disease resistance to this fungicide class, DO NOT make more than 2 sequential applications of VELUM or any Group 7-containing fungicide before rotating with a fungicide from a different Group.
- For soil application, to limit the potential for development of disease resistance to this chemical class, it is advised that the first foliar fungicide application after VELUM be a product from a different FRAC group.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937).

As with any crop-protection product, always read and follow label directions. For additional information call toll-free 1-866-99BAYER (1-866-992-2937).

Bayer CropScience LP 800 N. Lindbergh Blvd. St. Louis, MO 63167