



FIFRA 2(ee) Recommendation

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE(S) OF ARIZONA AND CALIFORNIA

VELUM® ONE

EPA Reg. No. 264-1078

For suppression of Fusarium basal rot in Bulb Vegetables (group 3-07)

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its label.

BULB VEGETABLES (group 3-07)

Onion, bulb subgroup: Bulb Daylily; Bulb Fritillaria; Bulb Garlic; Bulb Great-Headed Garlic; Bulb Serpent Garlic; Bulb Lily; Bulb Onion; Bulb Chinese Onion; Pearl Onion; Bulb Potato Onion; Bulb Shallot; cultivars, varieties, and/or hybrids of these.

Onion, green subgroup: Fresh Leaves Chive; Fresh Leaves Chinese Chive; Elegans Hosta; Leaves Fritillaria; Kurrat; Lady's Leek; Leek; Wild Leek; Beltsville Bunching Onion; Fresh Onion; Green Onion; Macrostem Onion; Tree Onion Tops; Welsh Onion Tops; Fresh Leaves Shallot; cultivars, varieties, and/or hybrids of these.

Disease Suppression	Application Rate	Application Instructions
Fusarium basal rot (<i>Fusarium spp.</i>)	6.25 to 6.84 fl oz/acre (0.203 – 0.222 lb/acre fluopyram)	Soil Application Apply specified dosage using any of the following methods: <ul style="list-style-type: none">• Pre-plant banded or broadcast spray directed to the soil.• In-furrow drench/spray during planting directed on or below seed.• Chemigation into the root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment.• Chemigation using overhead irrigation equipment.• Post-plant drench or hill drench. Minimum 14-day interval between applications.

BULB VEGETABLES (group 3-07) - continued**Restrictions:**

- Maximum single application rate: 6.84 fl oz/acre of VELUM ONE (0.222 lb/acre fluopyram)
- Maximum annual application rate: 13.7 fl oz of VELUM ONE per acre (0.445 lb/acre fluopyram) per year.
- Maximum number of applications per year: 2 (at 6.84 fl oz/acre of VELUM ONE)
- Do not apply more than 0.446 lbs Fluopyram per acre per year, regardless of formulation or method of application (soil or foliar).
- Apply using ground or chemigation equipment.
- For soil applications, **DO NOT** apply VELUM ONE within 30 days of harvest.
- To limit the potential for development of disease resistance to this fungicide class, **DO NOT** make more than 2 sequential applications of VELUM ONE or any Group 7-containing fungicide before rotating with a fungicide from a different Group.
- For soil application, to limit the potential for development of disease resistance to this chemical class, it is advised that the first foliar fungicide application after VELUM ONE be a product from a different FRAC group.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937).

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937).

Bayer CropScience LP
800 N. Lindbergh Blvd.
St. Louis, MO 63167