

**TECHNICAL INFORMATION BULLETIN  
FOR THE USE OF VENERATE XC TO CONTROL KATYDID IN STONE FRUIT**

**FIFRA Section 2(ee) Recommendation**

This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the U. S. Environmental Protection Agency.

FOR USE IN ALL STATES WHERE CROP AND PEST(S) EXIST AND WHERE 2(EE) RECOMMENDATIONS ARE RECOGNIZED.

**VENERATE® XC**

**EPA Reg. No. 84059-14**

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

<b>Crop</b>	<b>Target Arthropods</b>	<b>Application Method</b>	<b>Product Use Rate per Application</b>	<b>Application Instructions</b>
Stone Fruit	Fork-Tailed Bush Katydid ( <i>Scudderia furcata</i> ) and Angularwinged Katydid ( <i>Microcentrum retinerve</i> )	Foliar	1 qt-4 qt/Acre	<p>Use calibrated power-operated ground equipment capable of providing uniform coverage of the target crop. Use hollow cone, disc core/hollow cone or twin jet flat fan nozzles suitable for insecticide spraying.</p> <ul style="list-style-type: none"> <li>• Dilute spray application: This application method is based on the premise that all plant parts are thoroughly wetted, to the point of runoff, with spray solution. To determine the number of gallons of dilute spray per acre, contact your extension specialist, state agricultural experiment station, or certified pest control advisor for assistance.</li> <li>• Concentrate spray application: This application method is based on the premise that all plant parts are uniformly covered with spray solution but not to the point of runoff as with a dilute spray. Instead, a lower spray volume is used to deliver the same application rate of product per acre as is used for the dilute spray. Under heavy pest populations, use the higher label rates, shorten the spray interval, and/or apply in tank mixture with another product that has activity on the target pest.</li> </ul>

VENERATE® XC has a pre-harvest interval (PHI) of 0 days.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours without label-required personal protective equipment.

**All applicable directions, restrictions and precautions on the EPA-registered label are to be followed.**

This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and carefully observe the cautionary statements plus all the other information appearing on the product label.

**This recommendation must be in the possession of the user at the time of pesticide application.**

VENERATE® XC is a trademark of Marrone Bio Innovations, Inc.

© Copyright Marrone Bio Innovations, Inc., 2023

Marrone Bio Innovations, Inc.

1530 Drew Ave.,

Davis, CA 95618

Telephone (530) 750-2800

[info@marronebio.com](mailto:info@marronebio.com)

**1530 Drew Avenue • Davis, CA 95618 • Phone: 530-750-2800**