

SUPPLEMENTAL LABEL



PYRAFLUFEN-ETHYL	GROUP	14	HERBICIDE
------------------	-------	-----------	-----------

EPA Reg. No. 71711-25

For Use on Nonbearing Only Citrus Fruit Group

This supplemental label expires March 12, 2023 and must not be used or distributed after this date.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

This labeling and the EPA approved container label must be in the possession of the user at the time of application. Read the label affixed to the container for **VENUE** Herbicide before applying. Use of **VENUE** Herbicide according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for **VENUE** Herbicide

New use directions appear on this supplemental labeling that may be different from those that appear on the container label.

CROP USE DIRECTIONS

Citrus Fruit Group (Crop Group 10-10) Nonbearing Only				
Australian desert lime; Australian finger lime; Australian round lime; Brown River finger lime; calamondin; citron; citrus hybrids; grapefruit; Japanese summer grapefruit; kumquat; lemon; lime; Mediterranean mandarin; Mount White lime; New Guinea wild lime; orange, sour; orange, sweet; pummelo; Russell River lime; satsuma mandarin; sweet lime; tachibana orange; Tahiti lime; tangelo; tangerine (mandarin, clementine); tangor; trifoliolate orange; unig fruit; cultivars, varieties, and/or hybrids of these				
Application	Pest	Rate/Acre	Maximum Applications Per Year	Application Instructions
Dormant	Listed Broadleaf Weeds	3.0 to 4.0 fl oz/acre (0.0040 to 0.0053 lb ai/acre)	DO NOT exceed a combined total of 3 applications per year for this use.	<ul style="list-style-type: none"> • Apply in a minimum of 20 gallons spray solution per acre by ground equipment to target weeds. • Allow a minimum of 30 days between applications for this use. • The addition of a COC adjuvant at a concentration of 1% to 2% is directed for optimum weed control. Use the higher COC rate for larger labeled weed species or in low moisture conditions. • COC adjuvants are advised, though other adjuvants may be used.
In-Season	Listed Broadleaf Weeds	3.0 to 4.0 fl oz/acre (0.0040 to 0.0053 lb ai/acre)	DO NOT exceed a combined total of 3 applications per year for this use.	<ul style="list-style-type: none"> • Avoid spray contact with any foliage or low hanging branches (tree skirts) and with green, uncallused bark of young trees. • Use the higher rate for hard-to-control weeds. • DO NOT apply by air for this use. • DO NOT allow spray to drift onto foliage. • DO NOT exceed 6.8 fl oz/acre (0.0090 lb ai/acre) per year for dormant applications. • DO NOT exceed 12.0 fl oz/acre (0.0159 lb ai/acre) per year for all in-season applications combined.

©2020 Nichino America, Inc. Venue and Nichino America logo are registered trademarks of Nichino America, Inc.



Nichino America, Inc.
4550 Linden Hill Road, Suite 501
Wilmington, DE 19808
888-740-7700